

Clyde River Water Quality Report

Climate Smart Pilots September 7, 2023

www.farmdecisiontech.net.au

Foreword

Funding

This work has been produced by the NSW Primary Industries Climate Change Research Strategy funded by the NSW Climate Change Fund.

NSW Department of Primary Industries Disclaimer

This is a research trial and pilot project, and you should not rely solely on the information or advice provided in these reports.

Feedback and Questions

Please provide feedback and questions to Ben Sefton

Email: ben.sefton@dpi.nsw.gov.au

Ph: 0436 350 759

Contents


List of Figures

1 Salinity

1.1 Weekly

Weekly Minimum and Maximum Salinity Clyde River

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 1: These values represent the highest and lowest salinity reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Salinity Average Trend Clyde River

Average daily salinity at each buoy throughout the Clyde River.

Rocky Point Waterfall Moonlight

| Rocky Point V | Waterfall M | oonlight | | | | | |
|-------------------|-------------|----------|----------|----------|----------|----------|-----------|
| Rocky Point | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Waterfall Creek | 35.5 ppt | 35.7 ppt | 35.7 ppt | 35.8 ppt | 35.8 ppt | 35.9 ppt | 35.9 ppt |
| Rocky Point | 34.8 ppt | 35.0 ppt | 35.1 ppt | 35.2 ppt | 35.2 ppt | 35.3 ppt | 35.3 ppt |
| Angry Man Point | 33.0 ppt | 33.2 ppt | 33.3 ppt | 33.4 ppt | 33.4 ppt | 33.5 ppt | 33.5 ppt |
| Chinamans Point | 31.5 ppt | 31.6 ppt | 31.7 ppt | 31.8 ppt | 31.8 ppt | 31.9 ppt | 31.8 ppt |
| Waterfall | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Buckenbowra Creek | 35.5 ppt | 35.6 ppt | 35.8 ppt | 35.8 ppt | 35.9 ppt | 35.9 ppt | 35.9 ppt |
| Big Island West | 30.5 ppt | 30.8 ppt | 31.0 ppt | 31.1 ppt | 31.2 ppt | 31.2 ppt | 31.2 ppt |
| Double Bay | 26.2 ppt | 26.3 ppt | 26.4 ppt | 26.4 ppt | 26.5 ppt | 26.6 ppt | 26.7 ppt |
| Moonlight | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Snapper Point | 36.1 ppt | 36.2 ppt | 36.3 ppt | 36.3 ppt | 36.4 ppt | 36.4 ppt | 36.4 ppt |
| Wray Street | 33.0 ppt | 33.1 ppt | 33.1 ppt | 33.2 ppt | 33.2 ppt | 33.2 ppt | 33.3 ppt |
| Budd Island North | 32.9 ppt | 33.0 ppt | 33.0 ppt | 33.0 ppt | 33.1 ppt | 33.1 ppt | 33.2 ppt |
| Moonlight | | | | | | | |

Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.


Source: FarmDecisionTECH • Created with Datawrapper

Figure 2: This figure represents the daily average salinity of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week for specific locations within harvest areas.

1.2 Fortnightly

Fortnightly Salinity Average Trend Clyde River

12-hourly average salinity for each harvest area.




Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L. Source: FarmDecisionTECH • Created with Datawrapper

Figure 3: This figure represents the daily average salinity of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week.

Fortnightly Average Salinity Trend Clyde River

Comparison between this weeks average salinity and the average salinity last week. Left pointing arrow means salinity is decreasing, right pointing means salinity is increasing.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 4: This figure demonstrates the average difference in salinity for this week compared with the prior week. It displays a longer term, general trend to see if salinity is increasing, stabilising or decreasing.

2 Water Temperature

2.1 Weekly

Weekly Minimum and Maximum Water Temperature Clyde River

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Source: FarmDecisionTECH • Created with Datawrapper

Figure 5: These values represent the highest and lowest temperature reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Temperature Average Trend Clyde River

Average daily temperature at each buoy throughout the Clyde River.

| Waterfall Moonlight Rocky Point | | | | | | | |
|---------------------------------|----------|---------|----------|---------|---------|---------|-----------|
| Rocky Point | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Waterfall Creek | 16.6 °C | 16.5 °C | 16.4 °C | 16.4 °C | 16.4 °C | 16.5 °C | 16.5 °C |
| Rocky Point | 16.6 °C | 16.5 °C | 16.4°C | 16.4 °C | 16.4 °C | 16.4 °C | 16.4 °C |
| Angry Man Point | 16.4 °C | 16.3 °C | 16.2 °C | 16.2 °C | 16.3 °C | 16.3 °C | 16.3 °C |
| Chinamans Point | 16.4 °C | 16.4 °C | 16.3 °C | 16.2 °C | 16.3 °C | 16.3 °C | 16.3 °C |
| Waterfall | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Buckenbowra Creek | 16.3 °C | 16.3 °C | 16.2 °C | 16.2 °C | 16.2 °C | 16.2 °C | 16.2 °C |
| Big Island West | 16.1 °C | 16.2 °C | 16.1 °C | 16.1 °C | 16.1 °C | 16.2 °C | 16.2 °C |
| Double Bay | 16.1 °C | 16.1 °C | 16.0 °C | 16.1 °C | 16.1 °C | 16.2 °C | 16.2 °C |
| Moonlight | | | | | | | |
| | Thursday | Friday | Saturday | Sunday | Monday | Tuesday | Wednesday |
| Moonlight | 16.7 °C | 16.6 °C | 16.5 °C | 16.5 °C | 16.5 °C | 16.5 °C | 16.5 °C |
| Budd Island North | 16.7 °C | 16.6 °C | 16.5 °C | 16.4 °C | 16.5 °C | 16.5 °C | 16.6 °C |
| Wray Street | 16.6 °C | 16.5 °C | 16.4 °C | 16.4 °C | 16.4 °C | 16.4 °C | 16.4 °C |
| Snapper Point | 16.5 °C | 16.5 °C | 16.3 °C | 16.3 °C | 16.3 °C | 16.4 °C | 16.4 °C |

Temperature is displayed in degrees Celsius (°C)

Source: FarmDecisionTECH • Created with Datawrapper

Figure 6: This figure represents the daily average temperature of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week for specific locations within harvest areas.

2.2 Fortnightly

Fortnightly Average Temperature Trend Clyde River

12-hourly average temperature for each harvest area.




Figure 7: This figure represents the daily average temperature of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week.

Fortnightly Average Temperature Trend Clyde River

Comparison between this weeks average temperature and the average temperature last week.




Figure 8: This figure demonstrates the average difference in temperature for this week compared with the prior week. It displays a longer term, general trend to see if temperature is increasing, stabilising or decreasing.

2.3 Historical

Historical and Current Average Monthly Water Temperature Clyde River

Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.



Figure 9: Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Maximum Monthly Water Temperature Clyde River

Monthly maximum water temperature from historical (1967-73) and recent (2020-23) years. Measured close to the Clyde River entrance.



Figure 10: Monthly maximum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Minimum Monthly Water Temperature Clyde River

Monthly minimum water temperature from historical (1967-73) and recent (2020-22) years. Measured close to the Clyde River entrance.




Figure 11: Monthly minimum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

3 Precipitation

3.1 Weekly

7-Day Clyde River Precipitation Clyde River

Daily total precipitation. Weather station is located on Budd Island.

| Sep 01 | 02 | 03 | 04 | 05 | 06 | 07 |
|-----------|-----|-----|-----|-----|-----|-----|
| | | | | | | |
| | | | | | | |
| | | | | | | |
| 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |


 $Precipitation \ is \ measured \ in \ millimeters \ (mm). \ 1 \ mm \ of \ rainfall \ is \ equal \ to \ 1 \ litre \ of \ water \ per \ meter \ squared.$ Source: FarmDecisionTECH \cdot Created with Datawrapper

Figure 12: Daily total precipitation at Budd Island for the past week.

3.2 Yearly

Year-to-Date Precipitation Clyde River

Accumulation of rainfall since the start of the year. Other years are shown in reference to this year.



Rainfall is displayed in millimetres (mm).

Created with Datawrapper

Figure 13: Compares this years total precipitation against previous years.

4 Flow Rate from Tributaries

4.1 Fortnightly

Brooman Tributary Fortnightly Discharge rate Clyde River

Daily discharge rate totals for Brooman Tributary. This station is roughly 20 km North of the Clyde River.

Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Data: © State of New South Wales through WaterNSW. Quality Codes: 140 = Current rating - may be subject to change, 145 = Telemetry system added point, 255 = Dataset not complete

Source: FarmDecisionTECH • Created with Datawrapper

Figure 14: Fortnightly (daily total) water discharge from Brooman.

Buckenbowra Tributary Fortnightly Discharge rate Clyde River

Daily discharge rate totals for Buckenbowra Tributary. This station is roughly 8 km West of the Clyde River.

Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Quality Codes: 140 = Best available data, 145 = Data under review, 255 = Dataset not complete

Chart: © State of New South Wales through WaterNSW ${\bf \cdot}$ Created with Datawrapper

Figure 15: Fortnightly (daily total) water discharge from Buckenbowra.

This data is released by NSW Department of Primary Industries for evaluation purposes only and should not be relied upon for business decisions, carrying out risk assessments or other uses.

4.2 Yearly

Cumulative Daily Discharge-Rate Brooman Clyde River

Cumulative daily water flow for each year from Brooman into the Clyde River. Units represent mega litres (ML). One ML is equal to 1 million litres.



Data: © State of New South Wales through WaterNSW Chart: FarmDecisionTECH • Created with Datawrapper

Figure 16: Compares this years total water discharge against previous years. Drought years (2018 and 2019) are shown for comparison.