



Department of
Primary Industries

Clyde River Water Quality Report

Climate Smart Pilots
August 12, 2022

www.farmdecisiontech.net.au

Foreword

Funding

This work has been produced by the NSW Primary Industries Climate Change Research Strategy funded by the NSW Climate Change Fund.

NSW Department of Primary Industries Disclaimer

This is a research trial and pilot project, and you should not rely solely on the information or advice provided in these reports.

Feedback and Questions

Please provide feedback and questions to Harvey Bates

Email: harvey.bates@dpi.nsw.gov.au

Ph: 0447 359 557

Contents

1	Salinity	3
1.1	Weekly	3
1.2	Fortnightly	4
2	Water Temperature	6
2.1	Weekly	6
2.2	Fortnightly	7
2.3	Historical	9
3	Precipitation	12
3.1	Weekly	12
3.2	Yearly	13
4	Flow Rate from Tributaries	14
4.1	Fortnightly	14
4.2	Yearly	16

List of Figures


1	Weekly Minimum and Maximum Salinity	3
2	Average Weekly Salinity Table	3
3	Average Fortnightly Salinity Chart	4
4	Average Fortnightly Salinity Difference	5
5	Weekly Minimum and Maximum Temperature	6
6	Average Weekly Temperature Table	6
7	Average Fortnightly Temperature Chart	7
8	Average Fortnightly Temperature Difference	8
9	Historical Average Water Temperature	9
10	Historical Maximum Water Temperature	10
11	Historical Minimum Water Temperature	11
12	Daily Total Precipitation Budd Island	12
13	Yearly Cumulative Precipitation Budd Island	13
14	Fortnightly Discharge From Brooman Tributary	14
15	Fortnightly Discharge Rate Buckenbowra	15
16	Yearly Cumulative Discharge Rate Brooman	16

1 Salinity

1.1 Weekly

Weekly Minimum and Maximum Salinity

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 1: These values represent the highest and lowest salinity reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Salinity Average Trend

Average daily salinity at each buoy throughout the Clyde River.

Waterfall Moonlight Rocky Point

Rocky Point

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Waterfall Creek	25.3 ppt	25.0 ppt	25.2 ppt	26.1 ppt	26.8 ppt	27.4 ppt	27.4 ppt
Rocky Point	24.1 ppt	23.6 ppt	24.3 ppt	25.3 ppt	26.2 ppt	27.3 ppt	27.9 ppt
Chinamans Point	20.6 ppt	20.1 ppt	21.1 ppt	22.1 ppt	22.9 ppt	24.0 ppt	23.9 ppt
Angry Man Point	17.2 ppt	16.8 ppt	17.8 ppt	18.9 ppt	19.7 ppt	20.8 ppt	21.2 ppt

Waterfall

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Buckenbowra Creek	20.6 ppt	19.5 ppt	20.6 ppt	21.4 ppt	22.0 ppt	22.9 ppt	24.1 ppt
Big Island West	16.5 ppt	16.2 ppt	16.9 ppt	18.5 ppt	19.1 ppt	19.8 ppt	19.9 ppt
Double Bay	16.0 ppt	15.0 ppt	15.9 ppt	17.2 ppt	17.8 ppt	18.2 ppt	18.6 ppt

Moonlight

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Moonlight	25.4 ppt	24.5 ppt	26.0 ppt	27.1 ppt	27.8 ppt	28.7 ppt	29.2 ppt
Snapper Point	25.9 ppt	24.7 ppt	26.0 ppt	27.2 ppt	28.7 ppt	29.7 ppt	30.4 ppt
Budd Island North	23.6 ppt	23.3 ppt		25.2 ppt	26.0 ppt	26.8 ppt	27.2 ppt

Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.


Source: FarmDecisionTECH • Created with Datawrapper

Figure 2: This figure represents the daily average salinity of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week for specific locations within harvest areas.

1.2 Fortnightly

Fortnightly Salinity Average Trend

12-hourly average salinity for each harvest area.




Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 3: This figure represents the daily average salinity of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week.

Fortnightly Average Salinity Trend

Comparison between this weeks average salinity and the average salinity last week. Left pointing arrow means salinity is decreasing, right pointing means salinity is increasing.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.
Source: FarmDecisionTECH • Created with Datawrapper


Figure 4: This figure demonstrates the average difference in salinity for this week compared with the prior week. It displays a longer term, general trend to see if salinity is increasing, stabilising or decreasing.

2 Water Temperature

2.1 Weekly

Weekly Minimum and Maximum Water Temperature

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Source: FarmDecisionTECH • Created with Datawrapper

Figure 5: These values represent the highest and lowest temperature reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Temperature Average Trend

Average daily temperature at each buoy throughout the Clyde River.

Waterfall Moonlight Rocky Point

Rocky Point

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Rocky Point	13.7 °C	13.4 °C	13.5 °C	13.5 °C	13.6 °C	13.6 °C	13.6 °C
Chinamans Point	13.5 °C	13.2 °C	13.3 °C	13.3 °C	13.4 °C	13.4 °C	13.3 °C
Waterfall Creek	13.6 °C	13.3 °C	13.3 °C	13.3 °C	13.4 °C	13.4 °C	13.3 °C
Angry Man Point	13.2 °C	12.9 °C	13.0 °C	13.0 °C	13.2 °C	13.2 °C	13.2 °C

Waterfall

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Buckenbowra Creek	13.2 °C	13.0 °C	13.1 °C	13.0 °C	13.2 °C	13.1 °C	13.2 °C
Big Island West	13.0 °C	12.8 °C	13.0 °C	13.0 °C	13.1 °C	13.1 °C	13.0 °C
Double Bay	13.0 °C	12.8 °C	12.9 °C	12.8 °C	13.0 °C	12.9 °C	13.1 °C

Moonlight

	Saturday	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday
Moonlight	13.7 °C	13.5 °C	13.5 °C	13.5 °C	13.6 °C	13.7 °C	13.8 °C
Wray Street	13.7 °C	13.5 °C	13.5 °C	13.5 °C	13.7 °C	13.7 °C	13.7 °C
Snapper Point	13.6 °C	13.4 °C	13.5 °C	13.4 °C	13.6 °C	13.6 °C	13.6 °C
Budd Island North	13.6 °C	13.4 °C		13.5 °C	13.6 °C	13.6 °C	13.6 °C

Temperature is displayed in degrees Celsius (°C)


Source: FarmDecisionTECH • Created with Datawrapper

Figure 6: This figure represents the daily average temperature of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week for specific locations within harvest areas.

2.2 Fortnightly

Fortnightly Average Temperature Trend

12-hourly average temperature for each harvest area.




Temperature is displayed in degrees Celsius (°C).
Source: FarmDecisionTECH • Created with Datawrapper

Figure 7: This figure represents the daily average temperature of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week.

Fortnightly Average Temperature Trend

Comparison between this weeks average temperature and the average temperature last week.



Temperature is displayed in degrees Celsius (°C)
Source: FarmDecisionTECH • Created with Datawrapper

Figure 8: This figure demonstrates the average difference in temperature for this week compared with the prior week. It displays a longer term, general trend to see if temperature is increasing, stabilising or decreasing.

2.3 Historical

Historical and Current Average Monthly Water Temperature

Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.




Figure 9: Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Maximum Monthly Water Temperature

Monthly maximum water temperature from historical (1967-73) and recent (2020-22) years. Measured close to the Clyde River entrance.



Temperature is displayed in degrees Celsius (°C).

Source: FarmDecisionTECH • Created with Datawrapper

Figure 10: Monthly maximum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Minimum Monthly Water Temperature

Monthly minimum water temperature from historical (1967-73) and recent (2020-22) years. Measured close to the Clyde River entrance.



Temperature is displayed in degrees Celsius (°C).
Source: FarmDecisionTECH • Created with Datawrapper


Figure 11: Monthly minimum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

3 Precipitation

3.1 Weekly

7-Day Clyde River Precipitation

Daily total precipitation. Weather station is located on Budd Island.




Precipitation is measured in millimeters (mm). 1 mm of rainfall is equal to 1 litre of water per meter squared.
Source: FarmDecisionTECH • Created with Datawrapper

Figure 12: Daily total precipitation at Budd Island for the past week.

3.2 Yearly

Year-to-Date Precipitation

Accumulation of rainfall since the start of the year. Other years are shown in reference to this year.



Rainfall is displayed in millimetres (mm).

Created with Datawrapper


Figure 13: Compares this years total precipitation against previous years.

4 Flow Rate from Tributaries

4.1 Fortnightly

Brooman Tributary Fortnightly Discharge rate

Daily discharge rate totals for Brooman Tributary. This station is roughly 20 km North of the Clyde River.




Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Data: © State of New South Wales through WaterNSW. Quality Codes: 140 = Current rating - may be subject to change, 145 = Telemetry system added point, 255 = Dataset not complete
Source: FarmDecisionTECH • Created with Datawrapper

Figure 14: Fortnightly (daily total) water discharge from Brooman.

Buckenbowra Tributary Fortnightly Discharge rate

Daily discharge rate totals for Buckenbowra Tributary. This station is roughly 8 km West of the Clyde River.



Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Quality Codes: 140 = Best available data, 145 = Data under review, 255 = Dataset not complete


Chart: © State of New South Wales through WaterNSW • Created with Datawrapper

Figure 15: Fortnightly (daily total) water discharge from Buckenbowra.

4.2 Yearly

Cumulative Daily Discharge-Rate Brooman

Cumulative daily water flow for each year from Brooman into the Clyde River. Units represent mega litres (ML). One ML is equal to 1 million litres.



Data: © State of New South Wales through WaterNSW
Chart: FarmDecisionTECH • Created with Datawrapper

Figure 16: Compares this years total water discharge against previous years. Drought years (2018 and 2019) are shown for comparison.