

Clyde River Water Quality Report

Climate Smart Pilots April 26, 2022

www.farmdecisiontech.net.au

Foreword

Funding

This work has been produced by the NSW Primary Industries Climate Change Research Strategy funded by the NSW Climate Change Fund.

NSW Department of Primary Industries Disclaimer

This is a research trial and pilot project, and you should not rely solely on the information or advice provided in these reports.

Feedback and Questions

Please provide feedback and questions to Harvey Bates

Email: harvey.bates@dpi.nsw.gov.au

Ph: 0447 359 557

Contents


1		nity Weekly	
2	2.1	er Temperature Weekly Fortnightly	6
3	3.1	Cipitation Weekly Yearly	
4	4.1	v Rate from Tributaries 1 Fortnightly 1 Yearly 1	
Li	st o	of Figures	
	1 2 3	Weekly Minimum and Maximum Salinity	3
	4 5 6	Average Fortnightly Salinity Difference	5
	7 8 9		7 8
	10 11	Yearly Cumulative Precipitation Budd Island	1 O
	12 13	Fortnightly Discharge Rate Buckenbowra	12 13

1 Salinity

1.1 Weekly

Weekly Minimum and Maximum Salinity

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.




Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH · Created with Datawrapper

Figure 1: These values represent the highest and lowest salinity reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Salinity Average Trend

Average daily salinity at each buoy throughout the Clyde River.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.


Source: FarmDecisionTECH • Created with Datawrapper

Figure 2: This figure represents the daily average salinity of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week for specific locations within harvest areas.

1.2 Fortnightly

Fortnightly Salinity Average Trend

12-hourly average salinity for each harvest area.




Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L. Source: FarmDecisionTECH • Created with Datawrapper

Figure 3: This figure represents the daily average salinity of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week.

Fortnightly Average Salinity Trend

Comparison between this weeks average salinity and the average salinity last week. Left pointing arrow means salinity is decreasing, right pointing means salinity is increasing.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L. Source: FarmDecisionTECH • Created with Datawrapper


Figure 4: This figure demonstrates the average difference in salinity for this week compared with the prior week. It displays a longer term, general trend to see if salinity is increasing, stabilising or decreasing.

2 Water Temperature

2.1 Weekly

Weekly Minimum and Maximum Water Temperature

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Source: FarmDecisionTECH · Created with Datawrapper

Waterfall Moonlight Pocky Point

Figure 5: These values represent the highest and lowest temperature reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Temperature Average Trend

Average daily temperature at each buoy throughout the Clyde River.

Waterfall Mo	onlight Rock	xy Point					
Rocky Point							
	Wednesday	Thursday	Friday	Saturday	Sunday	Monday	Tuesday
Rocky Point	20.3 °C	19.9 °C	19.3 °C	19.2 °C	19.3 °C	19.5 °C	19.8 °C
Chinamans Point	19.9 °C	19.6 °C	19.2 °C	19.0 °C	19.1 °C	19.2 °C	19.5 °C
Waterfall Creek	20.2 °C	19.9 °C	19.3 °C	19.1 °C	19.1 °C	19.1 °C	19.4 °C
Angry Man Point							
Waterfall							
	Wednesday	Thursday	Friday	Saturday	Sunday	Monday	Tuesday
Buckenbowra Creek	19.6 °C	19.4 °C	18.9 °C	18.8 °C	18.9 °C	19.0 °C	19.2 °C
Big Island West	19.7 °C	19.4 °C	18.9 °C	18.9 °C	18.9 °C	18.8 °C	19.0 °C
Double Bay	19.4 °C	19.2 °C	18.8 °C	18.8 °C	18.9 °C	18.8 °C	18.9 °C
Moonlight .							
	Wednesday	Thursday	Friday	Saturday	Sunday	Monday	Tuesday
Moonlight	20.3 °C	19.9 °C	19.2 °C	19.3 °C	19.3 °C	19.5 °C	19.9 °C
Wray Street	20.3 °C	19.8 °C	19.3 °C	19.2 °C	19.3 °C	19.5 °C	19.8 °C
Budd Island North	20.3 °C	19.9 °C	19.3 °C	19.2 °C	19.3 °C	19.4 °C	19.7 °C
Snapper Point							


Temperature is displayed in degrees Celsius (°C)
Source: FarmDecisionTECH • Created with Datawrapper

Figure 6: This figure represents the daily average temperature of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week for specific locations within harvest areas.

2.2 Fortnightly

Fortnightly Average Temperature Trend

12-hourly average temperature for each harvest area.




Temperature is displayed in degrees Celsius (°C).
Source: FarmDecisionTECH • Created with Datawrapper

Figure 7: This figure represents the daily average temperature of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week.

Fortnightly Average Temperature Trend

Comparison between this weeks average temperature and the average temperature last week.



Temperature is displayed in degrees Celsius (°C)
Source: FarmDecisionTECH • Created with Datawrapper

Figure 8: This figure demonstrates the average difference in temperature for this week compared with the prior week. It displays a longer term, general trend to see if temperature is increasing, stabilising or decreasing.

3 Precipitation

3.1 Weekly

7-Day Clyde River Precipitation

Daily total precipitation. Weather station is located on Budd Island.

Apr 21	22	23	24	25	26
0.0 mm	0.0 mm	0.0 mm	0.0 mm	0.0 mm	0.0 mm


 $Precipitation \ is \ measured \ in \ millimeters \ (mm). \ 1 \ mm \ of \ rainfall \ is \ equal \ to \ 1 \ litre \ of \ water \ per \ meter \ squared.$ Source: FarmDecisionTECH \cdot Created with Datawrapper

Figure 9: Daily total precipitation at Budd Island for the past week.

3.2 Yearly

Year-to-Date Precipitation

Accumulation of rainfall since the start of the year. Other years are shown in reference to this year.



Rainfall is displayed in millimetres (mm).

Created with Datawrapper


Figure 10: Compares this years total precipitation against previous years.

4 Flow Rate from Tributaries

4.1 Fortnightly

Brooman Tributary Fortnightly Discharge rate

Daily discharge rate totals for Brooman Tributary. This station is roughly 20 km North of the Clyde River.




Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Data: © State of New South Wales through WaterNSW. Quality Codes: 140 = Current rating - may be subject to change, 145 = Telemetry system added point, 255 = Dataset not complete

Source: FarmDecisionTECH • Created with Datawrapper

Figure 11: Fortnightly (daily total) water discharge from Brooman.

Buckenbowra Tributary Fortnightly Discharge rate

Daily discharge rate totals for Buckenbowra Tributary. This station is roughly 8 km West of the Clyde River.



Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Quality Codes: 140 = Best available data, 145 = Data under review, 255 = Dataset not complete


Chart: © State of New South Wales through WaterNSW \bullet Created with Datawrapper

Figure 12: Fortnightly (daily total) water discharge from Buckenbowra.

4.2 Yearly

Cumulative Daily Discharge-Rate Brooman

Cumulative daily water flow for each year from Brooman into the Clyde River. Units represent mega litres (ML). One ML is equal to 1 million litres.



Data: © State of New South Wales through WaterNSW Chart: FarmDecisionTECH • Created with Datawrapper

Figure 13: Compares this years total water discharge against previous years. Drought years (2018 and 2019) are shown for comparison.