

Clyde River Water Quality Report

Climate Smart Pilots
December 7, 2022

www.farmdecisiontech.net.au

Foreword

Funding

This work has been produced by the NSW Primary Industries Climate Change Research Strategy funded by the NSW Climate Change Fund.

NSW Department of Primary Industries Disclaimer

This is a research trial and pilot project, and you should not rely solely on the information or advice provided in these reports.

Feedback and Questions

Please provide feedback and questions to Harvey Bates

Email: harvey.bates@dpi.nsw.gov.au

Ph: 0447 359 557

Contents

1	Sali	nity Weekly	
		Fortnightly	
2		er Temperature (
		Weekly	
		Fortnightly	
	2.3	Historical	Э
3		cipitation 12	
		Weekly	
	3.2	Yearly	3
4	Flov	v Rate from Tributaries	4
	4.1	Fortnightly	4
	4.2	Yearly	3
L	ist o	of Figures	
	1	Weekly Minimum and Maximum Salinity	3
	2	Average Weekly Salinity Table	
	3	Average Fortnightly Salinity Chart	
	4	Average Fortnightly Salinity Difference	
	5	Weekly Minimum and Maximum Temperature	
	6	Average Weekly Temperature Table	
	7	Average Fortnightly Temperature Chart	
	8	Average Fortnightly Temperature Difference	
	9	Historical Average Water Temperature	
	10	Historical Maximum Water Temperature	
	11	Historical Minimum Water Temperature	
	12	Daily Total Precipitation Budd Island	
	13	Yearly Cumulative Precipitation Budd Island	
	14 15	Fortnightly Discharge From Brooman Tributary	
	16	Fortnightly Discharge Rate Buckenbowra	
	10	ically Cullulative dischalge rate divullati	J

1 Salinity

1.1 Weekly

Weekly Minimum and Maximum Salinity

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 1: These values represent the highest and lowest salinity reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Salinity Average Trend

Waterfall Moonlight Rocky Point

Average daily salinity at each buoy throughout the Clyde River.

Rocky Point								
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday	
Waterfall Creek	27.3 ppt	27.1 ppt	26.9 ppt	27.0 ppt	27.5 ppt	28.3 ppt	28.8 ppt	
Angry Man Point	24.5 ppt	24.7 ppt	24.8 ppt	24.9 ppt	25.4 ppt	26.0 ppt	26.5 ppt	
Chinamans Point	24.5 ppt	24.4 ppt	24.3 ppt	24.4 ppt	25.1 ppt	25.6 ppt	26.0 ppt	
Rocky Point	21.7 ppl	21.9 ppt	21.7 ppt	21.7 ppt	22.3 ppt	22.7 ppt	22.7 pp	
Waterfall								
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday	
Buckenbowra Creek	24.4 ppt	24.4 ppt	24.2 ppt	24.0 ppt	24.6 ppt	24.8 ppt	25.5 ppt	
Big Island West	21.7 ppt	21.8 ppt	21.8 ppt	21.5 ppt	22.6 ppt	23.0 ppt	23.4 ppt	
Double Bay	20.6 ppt	20.3 ppt	20.4 ppt	20.6 ppt	21.1 ppt	21.4 ppt	22.0 ppt	
Moonlight								
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday	
Snapper Point	30.3 ppt	30.0 ppt	30.4 ppt	30.4 ppt	30.8 ppt	31.3 ppt	31.4 ppt	
Moonlight	28.6 ppt	28.4 ppt	28.3 ppt	28.3 ppt	29.1 ppt	29.4 ppt	29.7 ppt	
Wray Street	28.4 ppt	28.4 ppt	28.3 ppt	28.7 ppt	29.0 ppt	29.9 ppt	29.8 ppt	
Budd Island North	27.0 ppt	27.2 ppt	26.8 ppt	26.5 ppt	27.5 ppt	28.1 ppt	28.3 ppt	

Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L. Source: FarmDecisionTECH • Created with Datawrapper


Figure 2: This figure represents the daily average salinity of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week for specific locations within harvest areas.

This data is released by NSW Department of Primary Industries for evaluation purposes only and should not be relied upon for business decisions, carrying out risk assessments or other uses.

1.2 Fortnightly

Fortnightly Salinity Average Trend

12-hourly average salinity for each harvest area.




Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L.

Source: FarmDecisionTECH • Created with Datawrapper

Figure 3: This figure represents the daily average salinity of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of salinity over the past week.

Fortnightly Average Salinity Trend

Comparison between this weeks average salinity and the average salinity last week. Left pointing arrow means salinity is decreasing, right pointing means salinity is increasing.



Units represent parts per thousand (ppt). This is equivalent to both g/kg and g/L. Source: FarmDecisionTECH • Created with Datawrapper


Figure 4: This figure demonstrates the average difference in salinity for this week compared with the prior week. It displays a longer term, general trend to see if salinity is increasing, stabilising or decreasing.

2 Water Temperature

2.1 Weekly

Weekly Minimum and Maximum Water Temperature

These values represent the absolute minimum and maximum recorded values from Buoys within each harvest area.



Source: FarmDecisionTECH · Created with Datawrapper

Waterfall Moonlight Rocky Point

Figure 5: These values represent the highest and lowest temperature reading a buoy has recorded in each of the harvest areas in the past week.

7-Day Temperature Average Trend

Average daily temperature at each buoy throughout the Clyde River.

Rocky Point							
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday
Waterfall Creek	20.8 °C	20.9 °C	21.5 °C	22.4 °C	22.8 °C	22.5 °C	22.5 °C
Angry Man Point	20.7 °C	20.7 °C	21.4 °C	22.2 °C	22.4 °C	22.4 °C	22.5 °C
Chinamans Point	20.7 °C	20.7 °C	21.2 °C	22.1 °C	22.4 °C	22.3 °C	22.4 °C
Rocky Point	20.4 °C	20.4 °C	21.1 °C	21.9 °C	22.2 °C	21.9 °C	22.0 °C
Waterfall							
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday
Buckenbowra Creek	20.8 °C	20.8 °C	21.5 °C	22.4 °C	22.7 °C	22.4 °C	22.8 °C
Double Bay	20.9 °C	21.0 °C	21.6 °C	22.3 °C	22.6 °C	22.6 °C	22.9 °C
Big Island West	20.9 °C	20.9 °C	21.4 °C	22.2 °C	22.5 °C	22.5 °C	22.6 °C
Moonlight							
	Thursday	Friday	Saturday	Sunday	Monday	Tuesday	Wednesday
Snapper Point	20.4 °C	20.5 °C	21.0 °C	21.8 °C	22.0 °C	21.8 °C	22.0 °C
Budd Island North	20.3 °C	20.3 °C	21.0 °C	21.8 °C	21.8 °C	21.6 °C	21.7 °C
Moonlight	20.1 °C	20.2 °C	20.8 °C	21.5 °C	21.6 °C	21.5 °C	21.6 °C
Wray Street	20.0 °C	20.1 °C	20.7 °C	21.4 °C	21.5 °C	21.2 °C	21.4 °C

Figure 6: This figure represents the daily average temperature of each of the buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week for specific locations within harvest areas.

2.2 Fortnightly

Fortnightly Average Temperature Trend

12-hourly average temperature for each harvest area.




Figure 7: This figure represents the daily average temperature of all buoys contained within a harvest area. Its designed to reduce the impact of tides and provide the general trend of temperature over the past week.

Fortnightly Average Temperature Trend

Comparison between this weeks average temperature and the average temperature last week.




Figure 8: This figure demonstrates the average difference in temperature for this week compared with the prior week. It displays a longer term, general trend to see if temperature is increasing, stabilising or decreasing.

2.3 Historical

Historical and Current Average Monthly Water Temperature

Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.




Figure 9: Monthly average water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Maximum Monthly Water Temperature

Monthly maximum water temperature from historical (1967-73) and recent (2020-22) years. Measured close to the Clyde River entrance.




Figure 10: Monthly maximum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

Historical and Current Minimum Monthly Water Temperature

Monthly minimum water temperature from historical (1967-73) and recent (2020-22) years. Measured close to the Clyde River entrance.




Figure 11: Monthly minimum water temperature from historical (1967-73) and recent (2020-22). Measured close to the Clyde River entrance.

3 Precipitation

3.1 Weekly

7-Day Clyde River Precipitation

Daily total precipitation. Weather station is located on Budd Island.




Precipitation is measured in millimeters (mm). 1 mm of rainfall is equal to 1 litre of water per meter squared. Source: FarmDecisionTECH • Created with Datawrapper

Figure 12: Daily total precipitation at Budd Island for the past week.

3.2 Yearly

Year-to-Date Precipitation

Accumulation of rainfall since the start of the year. Other years are shown in reference to this year.



Rainfall is displayed in millimetres (mm).

Created with Datawrapper


Figure 13: Compares this years total precipitation against previous years.

4 Flow Rate from Tributaries

4.1 Fortnightly

Brooman Tributary Fortnightly Discharge rate

Daily discharge rate totals for Brooman Tributary. This station is roughly 20 km North of the Clyde River.




Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Data: © State of New South Wales through WaterNSW. Quality Codes: 140 = Current rating - may be subject to change, 145 = Telemetry system added point, 255 = Dataset not complete

Source: FarmDecisionTECH • Created with Datawrapper

Figure 14: Fortnightly (daily total) water discharge from Brooman.

Buckenbowra Tributary Fortnightly Discharge rate

Daily discharge rate totals for Buckenbowra Tributary. This station is roughly 8 km West of the Clyde River.



Values represent megalitres per day (ML/day). A mega litre is equal to 1 million litres. Quality Codes: 140 = Best available data, 145 = Data under review, 255 = Dataset not complete


Chart: © State of New South Wales through WaterNSW \bullet Created with Datawrapper

Figure 15: Fortnightly (daily total) water discharge from Buckenbowra.

4.2 Yearly

Cumulative Daily Discharge-Rate Brooman

Cumulative daily water flow for each year from Brooman into the Clyde River. Units represent mega litres (ML). One ML is equal to 1 million litres.



Data: © State of New South Wales through WaterNSW Chart: FarmDecisionTECH • Created with Datawrapper

Figure 16: Compares this years total water discharge against previous years. Drought years (2018 and 2019) are shown for comparison.