

Roelof Koekoek

Peter A. Lesky

René F. Swarttouw

Hypergeometric orthogonal polynomials and their q -analogues

– Monograph –

January 11, 2016

Springer

Berlin Heidelberg New York

Hong Kong London

Milan Paris Tokyo

Hypergeometric orthogonal polynomials

In this chapter we deal with all families of hypergeometric orthogonal polynomials appearing in the Askey scheme on page ?? . For each family of orthogonal polynomials we state the most important properties such as a representation as a hypergeometric function, orthogonality relation(s), the three-term recurrence relation, the second-order differential or difference equation, the forward shift (or degree lowering) and backward shift (or degree raising) operator, a Rodrigues-type formula and some generating functions. In each case we use the notation which seems to be most common in the literature. Moreover, in each case we mention the connection between various families by stating the appropriate limit relations. See also [?] for an algebraic approach of this Askey scheme and [?] for a view from asymptotic analysis. For notations the reader is referred to chapter ?? .

9.1 Wilson

Hypergeometric representation

$$\begin{aligned} & \frac{W_n(x^2; a, b, c, d)}{(a+b)_n(a+c)_n(a+d)_n} \\ &= {}_4F_3 \left(\begin{matrix} -n, n+a+b+c+d-1, a+ix, a-ix \\ a+b, a+c, a+d \end{matrix} ; 1 \right). \end{aligned} \quad (9.1.1)$$

Orthogonality relation

If $\operatorname{Re}(a, b, c, d) > 0$ and non-real parameters occur in conjugate pairs, then

$$\begin{aligned} & \frac{1}{2\pi} \int_0^\infty \left| \frac{\Gamma(a+ix)\Gamma(b+ix)\Gamma(c+ix)\Gamma(d+ix)}{\Gamma(2ix)} \right|^2 \\ & \quad \times W_m(x^2; a, b, c, d) W_n(x^2; a, b, c, d) dx \\ & = \frac{\Gamma(n+a+b) \cdots \Gamma(n+c+d)}{\Gamma(2n+a+b+c+d)} (n+a+b+c+d-1)_n n! \delta_{mn}, \end{aligned} \quad (9.1.2)$$

where

$$\begin{aligned} & \Gamma(n+a+b) \cdots \Gamma(n+c+d) \\ & = \Gamma(n+a+b)\Gamma(n+a+c)\Gamma(n+a+d)\Gamma(n+b+c)\Gamma(n+b+d)\Gamma(n+c+d). \end{aligned}$$

If $a < 0$ and $a+b, a+c, a+d$ are positive or a pair of complex conjugates occur with positive real parts, then

$$\begin{aligned} & \frac{1}{2\pi} \int_0^\infty \left| \frac{\Gamma(a+ix)\Gamma(b+ix)\Gamma(c+ix)\Gamma(d+ix)}{\Gamma(2ix)} \right|^2 \\ & \quad \times W_m(x^2; a, b, c, d) W_n(x^2; a, b, c, d) dx \\ & \quad + \frac{\Gamma(a+b)\Gamma(a+c)\Gamma(a+d)\Gamma(b-a)\Gamma(c-a)\Gamma(d-a)}{\Gamma(-2a)} \\ & \quad \times \sum_{\substack{k=0,1,2,\dots \\ a+k < 0}} \frac{(2a)_k (a+1)_k (a+b)_k (a+c)_k (a+d)_k}{(a)_k (a-b+1)_k (a-c+1)_k (a-d+1)_k k!} \\ & \quad \times W_m(-(a+k)^2; a, b, c, d) W_n(-(a+k)^2; a, b, c, d) \\ & = \frac{\Gamma(n+a+b) \cdots \Gamma(n+c+d)}{\Gamma(2n+a+b+c+d)} (n+a+b+c+d-1)_n n! \delta_{mn}. \end{aligned} \quad (9.1.3)$$

Recurrence relation

$$-(a^2 + x^2) \tilde{W}_n(x^2) = A_n \tilde{W}_{n+1}(x^2) - (A_n + C_n) \tilde{W}_n(x^2) + C_n \tilde{W}_{n-1}(x^2), \quad (9.1.4)$$

where

$$\tilde{W}_n(x^2) := \tilde{W}_n(x^2; a, b, c, d) = \frac{W_n(x^2; a, b, c, d)}{(a+b)_n (a+c)_n (a+d)_n}$$

and

$$\begin{cases} A_n = \frac{(n+a+b+c+d-1)(n+a+b)(n+a+c)(n+a+d)}{(2n+a+b+c+d-1)(2n+a+b+c+d)} \\ C_n = \frac{n(n+b+c-1)(n+b+d-1)(n+c+d-1)}{(2n+a+b+c+d-2)(2n+a+b+c+d-1)}. \end{cases}$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + (A_n + C_n - a^2)p_n(x) + A_{n-1}C_n p_{n-1}(x), \quad (9.1.5)$$

where

$$W_n(x^2; a, b, c, d) = (-1)^n (n+a+b+c+d-1)_n p_n(x^2).$$

Difference equation

$$\begin{aligned}
& n(n+a+b+c+d-1)y(x) \\
& = B(x)y(x+i) - [B(x) + D(x)]y(x) + D(x)y(x-i),
\end{aligned} \tag{9.1.6}$$

where

$$y(x) = W_n(x^2; a, b, c, d)$$

and

$$\begin{cases} B(x) = \frac{(a-ix)(b-ix)(c-ix)(d-ix)}{2ix(2ix-1)} \\ D(x) = \frac{(a+ix)(b+ix)(c+ix)(d+ix)}{2ix(2ix+1)}. \end{cases}$$

Forward shift operator

$$\begin{aligned}
& W_n((x + \tfrac{1}{2}i)^2; a, b, c, d) - W_n((x - \tfrac{1}{2}i)^2; a, b, c, d) \\
& = -2inx(n+a+b+c+d-1)W_{n-1}(x^2; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}, d + \tfrac{1}{2})
\end{aligned} \tag{9.1.7}$$

or equivalently

$$\begin{aligned}
& \frac{\delta W_n(x^2; a, b, c, d)}{\delta x^2} \\
& = -n(n+a+b+c+d-1)W_{n-1}(x^2; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}, d + \tfrac{1}{2}).
\end{aligned} \tag{9.1.8}$$

Backward shift operator

$$\begin{aligned}
& (a - \tfrac{1}{2} - ix)(b - \tfrac{1}{2} - ix)(c - \tfrac{1}{2} - ix)(d - \tfrac{1}{2} - ix)W_n((x + \tfrac{1}{2}i)^2; a, b, c, d) \\
& - (a - \tfrac{1}{2} + ix)(b - \tfrac{1}{2} + ix)(c - \tfrac{1}{2} + ix)(d - \tfrac{1}{2} + ix)W_n((x - \tfrac{1}{2}i)^2; a, b, c, d) \\
& = -2ixW_{n+1}(x^2; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2})
\end{aligned} \tag{9.1.9}$$

or equivalently

$$\begin{aligned}
& \frac{\delta [\omega(x; a, b, c, d)W_n(x^2; a, b, c, d)]}{\delta x^2} \\
& = \omega(x; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2})W_{n+1}(x^2; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2}),
\end{aligned} \tag{9.1.10}$$

where

$$\omega(x; a, b, c, d) := \frac{1}{2ix} \left| \frac{\Gamma(a + ix)\Gamma(b + ix)\Gamma(c + ix)\Gamma(d + ix)}{\Gamma(2ix)} \right|^2.$$

Rodrigues-type formula

$$\begin{aligned}
& \omega(x; a, b, c, d)W_n(x^2; a, b, c, d) \\
& = \left(\frac{\delta}{\delta x^2} \right)^n [\omega(x; a + \tfrac{1}{2}n, b + \tfrac{1}{2}n, c + \tfrac{1}{2}n, d + \tfrac{1}{2}n)].
\end{aligned} \tag{9.1.11}$$

Generating functions

$${}_2F_1 \left(\begin{matrix} a + ix, b + ix \\ a + b \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} c - ix, d - ix \\ c + d \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{W_n(x^2; a, b, c, d)t^n}{(a + b)_n(c + d)_nn!}. \tag{9.1.12}$$

$${}_2F_1 \left(\begin{matrix} a + ix, c + ix \\ a + c \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} b - ix, d - ix \\ b + d \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{W_n(x^2; a, b, c, d)t^n}{(a + c)_n(b + d)_nn!}. \tag{9.1.13}$$

$${}_2F_1 \left(\begin{matrix} a + ix, d + ix \\ a + d \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} b - ix, c - ix \\ b + c \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{W_n(x^2; a, b, c, d)t^n}{(a + d)_n(b + c)_nn!}. \tag{9.1.14}$$

$$\begin{aligned}
& (1 - t)^{1-a-b-c-d} \\
& \times {}_4F_3 \left(\begin{matrix} \frac{1}{2}(a + b + c + d - 1), \frac{1}{2}(a + b + c + d), a + ix, a - ix \\ a + b, a + c, a + d \end{matrix}; -\frac{4t}{(1 - t)^2} \right) \\
& = \sum_{n=0}^{\infty} \frac{(a + b + c + d - 1)_n}{(a + b)_n(a + c)_n(a + d)_nn!} W_n(x^2; a, b, c, d)t^n.
\end{aligned} \tag{9.1.15}$$

Limit relations**Wilson \rightarrow Continuous dual Hahn**

The continuous dual Hahn polynomials given by (9.3.1) can be found from the Wilson polynomials by dividing by $(a + d)_n$ and letting $d \rightarrow \infty$:

$$\lim_{d \rightarrow \infty} \frac{W_n(x^2; a, b, c, d)}{(a + d)_n} = S_n(x^2; a, b, c). \tag{9.1.16}$$

9.2 Racah

Hypergeometric representation

$$\begin{aligned} R_n(\lambda(x); \alpha, \beta, \gamma, \delta) \\ = {}_4F_3 \left(\begin{matrix} -n, n + \alpha + \beta + 1, -x, x + \gamma + \delta + 1 \\ \alpha + 1, \beta + \delta + 1, \gamma + 1 \end{matrix} ; 1 \right), \quad n = 0, 1, 2, \dots, N, \end{aligned} \quad (9.2.1)$$

where

$$\lambda(x) = x(x + \gamma + \delta + 1)$$

and

$$\alpha + 1 = -N \quad \text{or} \quad \beta + \delta + 1 = -N \quad \text{or} \quad \gamma + 1 = -N$$

with N a nonnegative integer.

Orthogonality relation

$$\begin{aligned} \sum_{x=0}^N \frac{(\alpha + 1)_x (\beta + \delta + 1)_x (\gamma + 1)_x (\gamma + \delta + 1)_x ((\gamma + \delta + 3)/2)_x}{(-\alpha + \gamma + \delta + 1)_x (-\beta + \gamma + 1)_x ((\gamma + \delta + 1)/2)_x (\delta + 1)_x x!} \\ \times R_m(\lambda(x)) R_n(\lambda(x)) \\ = M \frac{(n + \alpha + \beta + 1)_n (\alpha + \beta - \gamma + 1)_n (\alpha - \delta + 1)_n (\beta + 1)_n n!}{(\alpha + \beta + 2)_{2n} (\alpha + 1)_n (\beta + \delta + 1)_n (\gamma + 1)_n} \delta_{mn}, \end{aligned} \quad (9.2.2)$$

where

$$R_n(\lambda(x)) := R_n(\lambda(x); \alpha, \beta, \gamma, \delta)$$

and

$$M = \begin{cases} \frac{(-\beta)_N (\gamma + \delta + 2)_N}{(-\beta + \gamma + 1)_N (\delta + 1)_N} & \text{if } \alpha + 1 = -N \\ \frac{(-\alpha + \delta)_N (\gamma + \delta + 2)_N}{(-\alpha + \gamma + \delta + 1)_N (\delta + 1)_N} & \text{if } \beta + \delta + 1 = -N \\ \frac{(\alpha + \beta + 2)_N (-\delta)_N}{(\alpha - \delta + 1)_N (\beta + 1)_N} & \text{if } \gamma + 1 = -N. \end{cases}$$

Recurrence relation

$$\lambda(x) R_n(\lambda(x)) = A_n R_{n+1}(\lambda(x)) - (A_n + C_n) R_n(\lambda(x)) + C_n R_{n-1}(\lambda(x)), \quad (9.2.3)$$

where

$$R_n(\lambda(x)) := R_n(\lambda(x); \alpha, \beta, \gamma, \delta)$$

and

$$\begin{cases} A_n = \frac{(n + \alpha + 1)(n + \alpha + \beta + 1)(n + \beta + \delta + 1)(n + \gamma + 1)}{(2n + \alpha + \beta + 1)(2n + \alpha + \beta + 2)} \\ C_n = \frac{n(n + \alpha + \beta - \gamma)(n + \alpha - \delta)(n + \beta)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 1)}, \end{cases}$$

hence

$$A_n = \begin{cases} \frac{(n+\beta-N)(n+\beta+\delta+1)(n+\gamma+1)(n-N)}{(2n+\beta-N)(2n+\beta-N+1)} & \text{if } \alpha+1 = -N \\ \frac{(n+\alpha+1)(n+\alpha+\beta+1)(n+\gamma+1)(n-N)}{(2n+\alpha+\beta+1)(2n+\alpha+\beta+2)} & \text{if } \beta+\delta+1 = -N \\ \frac{(n+\alpha+1)(n+\alpha+\beta+1)(n+\beta+\delta+1)(n-N)}{(2n+\alpha+\beta+1)(2n+\alpha+\beta+2)} & \text{if } \gamma+1 = -N \end{cases}$$

and

$$C_n = \begin{cases} \frac{n(n+\beta)(n+\beta-\gamma-N-1)(n-\delta-N-1)}{(2n+\beta-N-1)(2n+\beta-N)} & \text{if } \alpha+1 = -N \\ \frac{n(n+\alpha+\beta+N+1)(n+\alpha+\beta-\gamma)(n+\beta)}{(2n+\alpha+\beta)(2n+\alpha+\beta+1)} & \text{if } \beta+\delta+1 = -N \\ \frac{n(n+\alpha+\beta+N+1)(n+\alpha-\delta)(n+\beta)}{(2n+\alpha+\beta)(2n+\alpha+\beta+1)} & \text{if } \gamma+1 = -N. \end{cases}$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) - (A_n + C_n)p_n(x) + A_{n-1}C_n p_{n-1}(x), \quad (9.2.4)$$

where

$$R_n(\lambda(x); \alpha, \beta, \gamma, \delta) = \frac{(n+\alpha+\beta+1)_n}{(\alpha+1)_n(\beta+\delta+1)_n(\gamma+1)_n} p_n(\lambda(x)).$$

Difference equation

$$n(n+\alpha+\beta+1)y(x) = B(x)y(x+1) - [B(x) + D(x)]y(x) + D(x)y(x-1), \quad (9.2.5)$$

where

$$y(x) = R_n(\lambda(x); \alpha, \beta, \gamma, \delta)$$

and

$$\begin{cases} B(x) = \frac{(x+\alpha+1)(x+\beta+\delta+1)(x+\gamma+1)(x+\gamma+\delta+1)}{(2x+\gamma+\delta+1)(2x+\gamma+\delta+2)} \\ D(x) = \frac{x(x-\alpha+\gamma+\delta)(x-\beta+\gamma)(x+\delta)}{(2x+\gamma+\delta)(2x+\gamma+\delta+1)}. \end{cases}$$

Forward shift operator

$$\begin{aligned} & R_n(\lambda(x+1); \alpha, \beta, \gamma, \delta) - R_n(\lambda(x); \alpha, \beta, \gamma, \delta) \\ &= \frac{n(n+\alpha+\beta+1)}{(\alpha+1)(\beta+\delta+1)(\gamma+1)} \\ & \quad \times (2x+\gamma+\delta+2)R_{n-1}(\lambda(x); \alpha+1, \beta+1, \gamma+1, \delta) \end{aligned} \quad (9.2.6)$$

or equivalently

$$\begin{aligned} & \frac{\Delta R_n(\lambda(x); \alpha, \beta, \gamma, \delta)}{\Delta \lambda(x)} \\ &= \frac{n(n+\alpha+\beta+1)}{(\alpha+1)(\beta+\delta+1)(\gamma+1)} R_{n-1}(\lambda(x); \alpha+1, \beta+1, \gamma+1, \delta). \end{aligned} \quad (9.2.7)$$

Backward shift operator

$$\begin{aligned}
& (x + \alpha)(x + \beta + \delta)(x + \gamma)(x + \gamma + \delta)R_n(\lambda(x); \alpha, \beta, \gamma, \delta) \\
& - x(x - \beta + \gamma)(x - \alpha + \gamma + \delta)(x + \delta)R_n(\lambda(x - 1); \alpha, \beta, \gamma, \delta) \\
& = \alpha\gamma(\beta + \delta)(2x + \gamma + \delta)R_{n+1}(\lambda(x); \alpha - 1, \beta - 1, \gamma - 1, \delta)
\end{aligned} \tag{9.2.8}$$

or equivalently

$$\begin{aligned}
& \frac{\nabla [\omega(x; \alpha, \beta, \gamma, \delta)R_n(\lambda(x); \alpha, \beta, \gamma, \delta)]}{\nabla \lambda(x)} \\
& = \frac{1}{\gamma + \delta} \omega(x; \alpha - 1, \beta - 1, \gamma - 1, \delta)R_{n+1}(\lambda(x); \alpha - 1, \beta - 1, \gamma - 1, \delta),
\end{aligned} \tag{9.2.9}$$

where

$$\omega(x; \alpha, \beta, \gamma, \delta) = \frac{(\alpha + 1)_x(\beta + \delta + 1)_x(\gamma + 1)_x(\gamma + \delta + 1)_x}{(-\alpha + \gamma + \delta + 1)_x(-\beta + \gamma + 1)_x(\delta + 1)_x x!}.$$

Rodrigues-type formula

$$\begin{aligned}
& \omega(x; \alpha, \beta, \gamma, \delta)R_n(\lambda(x); \alpha, \beta, \gamma, \delta) \\
& = (\gamma + \delta + 1)_n (\nabla_\lambda)^n [\omega(x; \alpha + n, \beta + n, \gamma + n, \delta)],
\end{aligned} \tag{9.2.10}$$

where

$$\nabla_\lambda := \frac{\nabla}{\nabla \lambda(x)}.$$

Generating functions

For $x = 0, 1, 2, \dots, N$ we have

$${}_2F_1 \left(\begin{matrix} -x, -x + \alpha - \gamma - \delta \\ \alpha + 1 \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} x + \beta + \delta + 1, x + \gamma + 1 \\ \beta + 1 \end{matrix}; t \right) = \sum_{n=0}^N \frac{(\beta + \delta + 1)_n (\gamma + 1)_n}{(\beta + 1)_n n!} R_n(\lambda(x); \alpha, \beta, \gamma, \delta) t^n, \tag{9.2.11}$$

$${}_2F_1 \left(\begin{matrix} -x, -x + \beta - \gamma \\ \beta + \delta + 1 \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} x + \alpha + 1, x + \gamma + 1 \\ \alpha - \delta + 1 \end{matrix}; t \right) = \sum_{n=0}^N \frac{(\alpha + 1)_n (\gamma + 1)_n}{(\alpha - \delta + 1)_n n!} R_n(\lambda(x); \alpha, \beta, \gamma, \delta) t^n, \tag{9.2.12}$$

$${}_2F_1 \left(\begin{matrix} -x, -x - \delta \\ \gamma + 1 \end{matrix}; t \right) {}_2F_1 \left(\begin{matrix} x + \alpha + 1, x + \beta + \delta + 1 \\ \alpha + \beta - \gamma + 1 \end{matrix}; t \right) = \sum_{n=0}^N \frac{(\alpha + 1)_n (\beta + \delta + 1)_n}{(\alpha + \beta - \gamma + 1)_n n!} R_n(\lambda(x); \alpha, \beta, \gamma, \delta) t^n, \tag{9.2.13}$$

$$\begin{aligned}
& \left[(1 - t)^{-\alpha - \beta - 1} \right. \\
& \quad \times {}_4F_3 \left(\begin{matrix} \frac{1}{2}(\alpha + \beta + 1), \frac{1}{2}(\alpha + \beta + 2), -x, x + \gamma + \delta + 1 \\ \alpha + 1, \beta + \delta + 1, \gamma + 1 \end{matrix}; -\frac{4t}{(1 - t)^2} \right) \Big]_N \\
& = \sum_{n=0}^N \frac{(\alpha + \beta + 1)_n}{n!} R_n(\lambda(x); \alpha, \beta, \gamma, \delta) t^n.
\end{aligned} \tag{9.2.14}$$

Limit relations**Racah \rightarrow Hahn**

The Hahn polynomials given by (9.5.1) can be obtained from the Racah polynomials by taking $\gamma + 1 = -N$ and letting $\delta \rightarrow \infty$:

$$\lim_{\delta \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, -N - 1, \delta) = Q_n(x; \alpha, \beta, N). \quad (9.2.15)$$

The Hahn polynomials given by (9.5.1) can also be obtained from the Racah polynomials by taking $\delta = -\beta - N - 1$ and letting $\gamma \rightarrow \infty$:

$$\lim_{\gamma \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, \gamma, -\beta - N - 1) = Q_n(x; \alpha, \beta, N). \quad (9.2.16)$$

Another way to do this is to take $\alpha + 1 = -N$ and $\beta \rightarrow \beta + \gamma + N + 1$ and then take the limit $\delta \rightarrow \infty$. In that case we obtain the Hahn polynomials given by (9.5.1) in the following way:

$$\lim_{\delta \rightarrow \infty} R_n(\lambda(x); -N - 1, \beta + \gamma + N + 1, \gamma, \delta) = Q_n(x; \gamma, \beta, N). \quad (9.2.17)$$

Racah \rightarrow Dual Hahn

The dual Hahn polynomials given by (9.6.1) are obtained from the Racah polynomials if we take $\alpha + 1 = -N$ and let $\beta \rightarrow \infty$:

$$\lim_{\beta \rightarrow \infty} R_n(\lambda(x); -N - 1, \beta, \gamma, \delta) = R_n(\lambda(x); \gamma, \delta, N). \quad (9.2.18)$$

The dual Hahn polynomials given by (9.6.1) are also obtained from the Racah polynomials if we take $\beta = -\delta - N - 1$ and let $\alpha \rightarrow \infty$:

$$\lim_{\alpha \rightarrow \infty} R_n(\lambda(x); \alpha, -\delta - N - 1, \gamma, \delta) = R_n(\lambda(x); \gamma, \delta, N). \quad (9.2.19)$$

Finally, the dual Hahn polynomials given by (9.6.1) are also obtained from the Racah polynomials if we take $\gamma + 1 = -N$ and $\delta \rightarrow \alpha + \delta + N + 1$ and take the limit $\beta \rightarrow \infty$:

$$\lim_{\beta \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, -N - 1, \alpha + \delta + N + 1) = R_n(\lambda(x); \alpha, \delta, N). \quad (9.2.20)$$

Remark

If we set $\alpha = a + b - 1$, $\beta = c + d - 1$, $\gamma = a + d - 1$, $\delta = a - d$ and $x \rightarrow -a + ix$ in the definition (9.2.1) of the Racah polynomials we obtain the Wilson polynomials given by (9.1.1):

$$\begin{aligned} R_n(\lambda(-a + ix); a + b - 1, c + d - 1, a + d - 1, a - d) \\ = \tilde{W}_n(x^2; a, b, c, d) = \frac{W_n(x^2; a, b, c, d)}{(a + b)_n(a + c)_n(a + d)_n}. \end{aligned}$$

References

[?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?].

9.3 Continuous dual Hahn**Hypergeometric representation**

$$\frac{S_n(x^2; a, b, c)}{(a + b)_n(a + c)_n} = {}_3F_2 \left(\begin{matrix} -n, a + ix, a - ix \\ a + b, a + c \end{matrix}; 1 \right). \quad (9.3.1)$$

Orthogonality relation

If a, b and c are positive except possibly for a pair of complex conjugates with positive real parts, then

$$\begin{aligned} \frac{1}{2\pi} \int_0^\infty \left| \frac{\Gamma(a + ix)\Gamma(b + ix)\Gamma(c + ix)}{\Gamma(2ix)} \right|^2 S_m(x^2; a, b, c) S_n(x^2; a, b, c) dx \\ = \Gamma(n + a + b)\Gamma(n + a + c)\Gamma(n + b + c)n! \delta_{mn}. \end{aligned} \quad (9.3.2)$$

If $a < 0$ and $a + b, a + c$ are positive or a pair of complex conjugates with positive real parts, then

$$\begin{aligned} \frac{1}{2\pi} \int_0^\infty \left| \frac{\Gamma(a + ix)\Gamma(b + ix)\Gamma(c + ix)}{\Gamma(2ix)} \right|^2 S_m(x^2; a, b, c) S_n(x^2; a, b, c) dx \\ + \frac{\Gamma(a + b)\Gamma(a + c)\Gamma(b - a)\Gamma(c - a)}{\Gamma(-2a)} \\ \times \sum_{\substack{k=0,1,2,\dots \\ a+k < 0}} \frac{(2a)_k(a+1)_k(a+b)_k(a+c)_k}{(a)_k(a-b+1)_k(a-c+1)_k k!} (-1)^k \\ \times S_m(-(a+k)^2; a, b, c) S_n(-(a+k)^2; a, b, c) \\ = \Gamma(n + a + b)\Gamma(n + a + c)\Gamma(n + b + c)n! \delta_{mn}. \end{aligned} \quad (9.3.3)$$

Recurrence relation

$$-(a^2 + x^2) \tilde{S}_n(x^2) = A_n \tilde{S}_{n+1}(x^2) - (A_n + C_n) \tilde{S}_n(x^2) + C_n \tilde{S}_{n-1}(x^2), \quad (9.3.4)$$

where

$$\tilde{S}_n(x^2) := \tilde{S}_n(x^2; a, b, c) = \frac{S_n(x^2; a, b, c)}{(a + b)_n(a + c)_n}$$

and

$$\begin{cases} A_n = (n + a + b)(n + a + c) \\ C_n = n(n + b + c - 1). \end{cases}$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + (A_n + C_n - a^2)p_n(x) + A_{n-1}C_np_{n-1}(x), \quad (9.3.5)$$

where

$$S_n(x^2; a, b, c) = (-1)^n p_n(x^2).$$

Difference equation

$$ny(x) = B(x)y(x+i) - [B(x) + D(x)]y(x) + D(x)y(x-i), \quad (9.3.6)$$

where

$$y(x) = S_n(x^2; a, b, c)$$

and

$$\begin{cases} B(x) = \frac{(a-ix)(b-ix)(c-ix)}{2ix(2ix-1)} \\ D(x) = \frac{(a+ix)(b+ix)(c+ix)}{2ix(2ix+1)}. \end{cases}$$

Forward shift operator

$$\begin{aligned} & S_n((x + \tfrac{1}{2}i)^2; a, b, c) - S_n((x - \tfrac{1}{2}i)^2; a, b, c) \\ &= -2inxS_{n-1}(x^2; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}) \end{aligned} \quad (9.3.7)$$

or equivalently

$$\frac{\delta S_n(x^2; a, b, c)}{\delta x^2} = -nS_{n-1}(x^2; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}). \quad (9.3.8)$$

Backward shift operator

$$\begin{aligned} & (a - \tfrac{1}{2} - ix)(b - \tfrac{1}{2} - ix)(c - \tfrac{1}{2} - ix)S_n((x + \tfrac{1}{2}i)^2; a, b, c) \\ & - (a - \tfrac{1}{2} + ix)(b - \tfrac{1}{2} + ix)(c - \tfrac{1}{2} + ix)S_n((x - \tfrac{1}{2}i)^2; a, b, c) \\ &= -2ixS_{n+1}(x^2; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}) \end{aligned} \quad (9.3.9)$$

or equivalently

$$\begin{aligned} & \frac{\delta [\omega(x; a, b, c)S_n(x^2; a, b, c)]}{\delta x^2} \\ &= \omega(x; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2})S_{n+1}(x^2; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}), \end{aligned} \quad (9.3.10)$$

where

$$\omega(x; a, b, c) = \frac{1}{2ix} \left| \frac{\Gamma(a+ix)\Gamma(b+ix)\Gamma(c+ix)}{\Gamma(2ix)} \right|^2.$$

Rodrigues-type formula

$$\omega(x; a, b, c)S_n(x^2; a, b, c) = \left(\frac{\delta}{\delta x^2} \right)^n [\omega(x; a + \tfrac{1}{2}n, b + \tfrac{1}{2}n, c + \tfrac{1}{2}n)]. \quad (9.3.11)$$

Generating functions

$$(1-t)^{-c+ix} {}_2F_1 \left(\begin{matrix} a+ix, b+ix \\ a+b \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{S_n(x^2; a, b, c)}{(a+b)_n n!} t^n. \quad (9.3.12)$$

$$(1-t)^{-b+ix} {}_2F_1 \left(\begin{matrix} a+ix, c+ix \\ a+c \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{S_n(x^2; a, b, c)}{(a+c)_n n!} t^n. \quad (9.3.13)$$

$$(1-t)^{-a+ix} {}_2F_1 \left(\begin{matrix} b+ix, c+ix \\ b+c \end{matrix}; t \right) = \sum_{n=0}^{\infty} \frac{S_n(x^2; a, b, c)}{(b+c)_n n!} t^n. \quad (9.3.14)$$

$$e^t {}_2F_2 \left(\begin{matrix} a+ix, a-ix \\ a+b, a+c \end{matrix}; -t \right) = \sum_{n=0}^{\infty} \frac{S_n(x^2; a, b, c)}{(a+b)_n (a+c)_n n!} t^n. \quad (9.3.15)$$

$$\begin{aligned} & (1-t)^{-\gamma} {}_3F_2 \left(\begin{matrix} \gamma, a+ix, a-ix \\ a+b, a+c \end{matrix}; \frac{t}{t-1} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n S_n(x^2; a, b, c)}{(a+b)_n (a+c)_n n!} t^n, \quad \gamma \text{ arbitrary.} \end{aligned} \quad (9.3.16)$$

Limit relations**Wilson \rightarrow Continuous dual Hahn**

The continuous dual Hahn polynomials can be found from the Wilson polynomials given by (9.1.1) by dividing by $(a+d)_n$ and letting $d \rightarrow \infty$:

$$\lim_{d \rightarrow \infty} \frac{W_n(x^2; a, b, c, d)}{(a+d)_n} = S_n(x^2; a, b, c).$$

Continuous dual Hahn \rightarrow Meixner-Pollaczek

The Meixner-Pollaczek polynomials given by (9.7.1) can be obtained from the continuous dual Hahn polynomials by the substitutions $x \rightarrow x-t$, $a = \lambda + it$, $b = \lambda - it$ and $c = t \cot \phi$ and the limit $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} \frac{S_n((x-t)^2; \lambda + it, \lambda - it, t \cot \phi)}{t^n n!} = \frac{P_n^{(\lambda)}(x; \phi)}{(\sin \phi)^n}. \quad (9.3.17)$$

Remark

Since we have for $k < n$

$$\frac{(a+b)_n (a+c)_n}{(a+b)_k (a+c)_k} = (a+b+k)_{n-k} (a+c+k)_{n-k},$$

the continuous dual Hahn polynomials defined by (9.3.1) can also be seen as polynomials in the parameters a , b and c .

References

[?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?].

9.4 Continuous Hahn

Hypergeometric representation

$$\begin{aligned} p_n(x; a, b, c, d) \\ = i^n \frac{(a+c)_n (a+d)_n}{n!} {}_3F_2 \left(\begin{matrix} -n, n+a+b+c+d-1, a+ix \\ a+c, a+d \end{matrix}; 1 \right). \end{aligned} \quad (9.4.1)$$

Orthogonality relation

If $\operatorname{Re}(a, b, c, d) > 0$, $c = \bar{a}$ and $d = \bar{b}$, then

$$\begin{aligned} \frac{1}{2\pi} \int_{-\infty}^{\infty} \Gamma(a+ix) \Gamma(b+ix) \Gamma(c-ix) \Gamma(d-ix) p_m(x; a, b, c, d) p_n(x; a, b, c, d) dx \\ = \frac{\Gamma(n+a+c) \Gamma(n+a+d) \Gamma(n+b+c) \Gamma(n+b+d)}{(2n+a+b+c+d-1) \Gamma(n+a+b+c+d-1) n!} \delta_{mn}. \end{aligned} \quad (9.4.2)$$

Recurrence relation

$$(a+ix) \tilde{p}_n(x) = A_n \tilde{p}_{n+1}(x) - (A_n + C_n) \tilde{p}_n(x) + C_n \tilde{p}_{n-1}(x), \quad (9.4.3)$$

where

$$\tilde{p}_n(x) := \tilde{p}_n(x; a, b, c, d) = \frac{n!}{i^n (a+c)_n (a+d)_n} p_n(x; a, b, c, d)$$

and

$$\begin{cases} A_n = -\frac{(n+a+b+c+d-1)(n+a+c)(n+a+d)}{(2n+a+b+c+d-1)(2n+a+b+c+d)} \\ C_n = \frac{n(n+b+c-1)(n+b+d-1)}{(2n+a+b+c+d-2)(2n+a+b+c+d-1)}. \end{cases}$$

Normalized recurrence relation

$$x p_n(x) = p_{n+1}(x) + i(A_n + C_n + a) p_n(x) - A_{n-1} C_n p_{n-1}(x), \quad (9.4.4)$$

where

$$p_n(x; a, b, c, d) = \frac{(n+a+b+c+d-1)_n}{n!} \tilde{p}_n(x).$$

Difference equation

$$\begin{aligned} n(n+a+b+c+d-1)y(x) \\ = B(x)y(x+i) - [B(x) + D(x)]y(x) + D(x)y(x-i), \end{aligned} \quad (9.4.5)$$

where

$$y(x) = p_n(x; a, b, c, d)$$

and

$$\begin{cases} B(x) = (c-ix)(d-ix) \\ D(x) = (a+ix)(b+ix). \end{cases}$$

Forward shift operator

$$\begin{aligned}
& p_n(x + \tfrac{1}{2}i; a, b, c, d) - p_n(x - \tfrac{1}{2}i; a, b, c, d) \\
& = i(n + a + b + c + d - 1)p_{n-1}(x; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}, d + \tfrac{1}{2})
\end{aligned} \tag{9.4.6}$$

or equivalently

$$\frac{\delta p_n(x; a, b, c, d)}{\delta x} = (n + a + b + c + d - 1)p_{n-1}(x; a + \tfrac{1}{2}, b + \tfrac{1}{2}, c + \tfrac{1}{2}, d + \tfrac{1}{2}). \tag{9.4.7}$$

Backward shift operator

$$\begin{aligned}
& (c - \tfrac{1}{2} - ix)(d - \tfrac{1}{2} - ix)p_n(x + \tfrac{1}{2}i; a, b, c, d) \\
& \quad - (a - \tfrac{1}{2} + ix)(b - \tfrac{1}{2} + ix)p_n(x - \tfrac{1}{2}i; a, b, c, d) \\
& = \frac{n+1}{i}p_{n+1}(x; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2})
\end{aligned} \tag{9.4.8}$$

or equivalently

$$\begin{aligned}
& \frac{\delta [\omega(x; a, b, c, d)p_n(x; a, b, c, d)]}{\delta x} \\
& = -(n+1)\omega(x; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2}) \\
& \quad \times p_{n+1}(x; a - \tfrac{1}{2}, b - \tfrac{1}{2}, c - \tfrac{1}{2}, d - \tfrac{1}{2}),
\end{aligned} \tag{9.4.9}$$

where

$$\omega(x; a, b, c, d) = \Gamma(a + ix)\Gamma(b + ix)\Gamma(c - ix)\Gamma(d - ix).$$

Rodrigues-type formula

$$\begin{aligned}
& \omega(x; a, b, c, d)p_n(x; a, b, c, d) \\
& = \frac{(-1)^n}{n!} \left(\frac{\delta}{\delta x} \right)^n [\omega(x; a + \tfrac{1}{2}n, b + \tfrac{1}{2}n, c + \tfrac{1}{2}n, d + \tfrac{1}{2}n)].
\end{aligned} \tag{9.4.10}$$

Generating functions

$${}_1F_1 \left(\begin{matrix} a + ix \\ a + c \end{matrix}; -it \right) {}_1F_1 \left(\begin{matrix} d - ix \\ b + d \end{matrix}; it \right) = \sum_{n=0}^{\infty} \frac{p_n(x; a, b, c, d)}{(a + c)_n(b + d)_n} t^n. \tag{9.4.11}$$

$${}_1F_1 \left(\begin{matrix} a + ix \\ a + d \end{matrix}; -it \right) {}_1F_1 \left(\begin{matrix} c - ix \\ b + c \end{matrix}; it \right) = \sum_{n=0}^{\infty} \frac{p_n(x; a, b, c, d)}{(a + d)_n(b + c)_n} t^n. \tag{9.4.12}$$

$$\begin{aligned}
& (1 - t)^{1-a-b-c-d} {}_3F_2 \left(\begin{matrix} \frac{1}{2}(a + b + c + d - 1), \frac{1}{2}(a + b + c + d), a + ix \\ a + c, a + d \end{matrix}; -\frac{4t}{(1 - t)^2} \right) \\
& = \sum_{n=0}^{\infty} \frac{(a + b + c + d - 1)_n}{(a + c)_n(a + d)_n i^n} p_n(x; a, b, c, d) t^n.
\end{aligned} \tag{9.4.13}$$

Limit relations

Wilson \rightarrow Continuous Hahn

The continuous Hahn polynomials are obtained from the Wilson polynomials given by (9.1.1) by the substitution $a \rightarrow a - it$, $b \rightarrow b - it$, $c \rightarrow c + it$, $d \rightarrow d + it$ and $x \rightarrow x + t$ and the limit $t \rightarrow \infty$ in the following way:

$$\lim_{t \rightarrow \infty} \frac{W_n((x+t)^2; a-it, b-it, c+it, d+it)}{(-2t)^n n!} = p_n(x; a, b, c, d).$$

Continuous Hahn \rightarrow Meixner-Pollaczek

The Meixner-Pollaczek polynomials given by (9.7.1) can be obtained from the continuous Hahn polynomials by setting $x \rightarrow x + t$, $a = \lambda - it$, $c = \lambda + it$ and $b = d = t \tan \phi$ and taking the limit $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} \frac{p_n(x+t; \lambda - it, t \tan \phi, \lambda + it, t \tan \phi)}{t^n} = \frac{P_n^{(\lambda)}(x; \phi)}{(\cos \phi)^n}. \quad (9.4.14)$$

Continuous Hahn \rightarrow Jacobi

The Jacobi polynomials given by (9.8.1) follow from the continuous Hahn polynomials by the substitution $x \rightarrow \frac{1}{2}xt$, $a = \frac{1}{2}(\alpha + 1 - it)$, $b = \frac{1}{2}(\beta + 1 + it)$, $c = \frac{1}{2}(\alpha + 1 + it)$ and $d = \frac{1}{2}(\beta + 1 - it)$, division by t^n and the limit $t \rightarrow \infty$:

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{p_n(\frac{1}{2}xt; \frac{1}{2}(\alpha + 1 - it), \frac{1}{2}(\beta + 1 + it), \frac{1}{2}(\alpha + 1 + it), \frac{1}{2}(\beta + 1 - it))}{t^n} \\ &= P_n^{(\alpha, \beta)}(x). \end{aligned} \quad (9.4.15)$$

Continuous Hahn \rightarrow Pseudo Jacobi

The pseudo Jacobi polynomials given by (9.9.1) follow from the continuous Hahn polynomials by the substitution $x \rightarrow xt$, $a = \frac{1}{2}(-N + iv - 2t)$, $b = \frac{1}{2}(-N - iv + 2t)$, $c = \frac{1}{2}(-N - iv - 2t)$ and $d = \frac{1}{2}(-N + iv + 2t)$, division by t^n and the limit $t \rightarrow \infty$:

$$\begin{aligned} & \lim_{t \rightarrow \infty} \frac{p_n(xt; \frac{1}{2}(-N + iv - 2t), \frac{1}{2}(-N - iv + 2t), \frac{1}{2}(-N - iv - 2t), \frac{1}{2}(-N + iv + 2t))}{t^n} \\ &= \frac{(n - 2N - 1)_n}{n!} P_n(x; v, N). \end{aligned} \quad (9.4.16)$$

Remark

Since we have for $k < n$

$$\frac{(a+b)_n (a+c)_n}{(a+b)_k (a+c)_k} = (a+b+k)_{n-k} (a+c+k)_{n-k},$$

the continuous Hahn polynomials defined by (9.4.1) can also be seen as polynomials in the parameters a , b and c .

References

[?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?].

9.5 Hahn

Hypergeometric representation

$$Q_n(x; \alpha, \beta, N) = {}_3F_2 \left(\begin{matrix} -n, n + \alpha + \beta + 1, -x \\ \alpha + 1, -N \end{matrix}; 1 \right), \quad n = 0, 1, 2, \dots, N. \quad (9.5.1)$$

Orthogonality relation

For $\alpha > -1$ and $\beta > -1$, or for $\alpha < -N$ and $\beta < -N$, we have

$$\begin{aligned} & \sum_{x=0}^N \binom{\alpha+x}{x} \binom{\beta+N-x}{N-x} Q_m(x; \alpha, \beta, N) Q_n(x; \alpha, \beta, N) \\ &= \frac{(-1)^n (n + \alpha + \beta + 1)_{N+1} (\beta + 1)_n n!}{(2n + \alpha + \beta + 1) (\alpha + 1)_n (-N)_n N!} \delta_{mn}. \end{aligned} \quad (9.5.2)$$

Recurrence relation

$$-xQ_n(x) = A_n Q_{n+1}(x) - (A_n + C_n) Q_n(x) + C_n Q_{n-1}(x), \quad (9.5.3)$$

where

$$Q_n(x) := Q_n(x; \alpha, \beta, N)$$

and

$$\begin{cases} A_n = \frac{(n + \alpha + \beta + 1)(n + \alpha + 1)(N - n)}{(2n + \alpha + \beta + 1)(2n + \alpha + \beta + 2)} \\ C_n = \frac{n(n + \alpha + \beta + N + 1)(n + \beta)}{(2n + \alpha + \beta)(2n + \alpha + \beta + 1)}. \end{cases}$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + (A_n + C_n) p_n(x) + A_{n-1} C_n p_{n-1}(x), \quad (9.5.4)$$

where

$$Q_n(x; \alpha, \beta, N) = \frac{(n + \alpha + \beta + 1)_n}{(\alpha + 1)_n (-N)_n} p_n(x).$$

Difference equation

$$n(n + \alpha + \beta + 1)y(x) = B(x)y(x+1) - [B(x) + D(x)]y(x) + D(x)y(x-1), \quad (9.5.5)$$

where

$$y(x) = Q_n(x; \alpha, \beta, N)$$

and

$$\begin{cases} B(x) = (x + \alpha + 1)(x - N) \\ D(x) = x(x - \beta - N - 1). \end{cases}$$

Forward shift operator

$$\begin{aligned} & Q_n(x+1; \alpha, \beta, N) - Q_n(x; \alpha, \beta, N) \\ &= -\frac{n(n + \alpha + \beta + 1)}{(\alpha + 1)N} Q_{n-1}(x; \alpha + 1, \beta + 1, N - 1) \end{aligned} \quad (9.5.6)$$

or equivalently

$$\Delta Q_n(x; \alpha, \beta, N) = -\frac{n(n + \alpha + \beta + 1)}{(\alpha + 1)N} Q_{n-1}(x; \alpha + 1, \beta + 1, N - 1). \quad (9.5.7)$$

Backward shift operator

$$\begin{aligned}
& (x + \alpha)(N + 1 - x)Q_n(x; \alpha, \beta, N) - x(\beta + N + 1 - x)Q_n(x - 1; \alpha, \beta, N) \\
& = \alpha(N + 1)Q_{n+1}(x; \alpha - 1, \beta - 1, N + 1)
\end{aligned} \tag{9.5.8}$$

or equivalently

$$\begin{aligned}
& \nabla [\omega(x; \alpha, \beta, N)Q_n(x; \alpha, \beta, N)] \\
& = \frac{N + 1}{\beta} \omega(x; \alpha - 1, \beta - 1, N + 1)Q_{n+1}(x; \alpha - 1, \beta - 1, N + 1),
\end{aligned} \tag{9.5.9}$$

where

$$\omega(x; \alpha, \beta, N) = \binom{\alpha + x}{x} \binom{\beta + N - x}{N - x}.$$

Rodrigues-type formula

$$\begin{aligned}
& \omega(x; \alpha, \beta, N)Q_n(x; \alpha, \beta, N) \\
& = \frac{(-1)^n(\beta + 1)_n}{(-N)_n} \nabla^n [\omega(x; \alpha + n, \beta + n, N - n)].
\end{aligned} \tag{9.5.10}$$

Generating functions

For $x = 0, 1, 2, \dots, N$ we have

$${}_1F_1 \left(\begin{matrix} -x \\ \alpha + 1 \end{matrix}; -t \right) {}_1F_1 \left(\begin{matrix} x - N \\ \beta + 1 \end{matrix}; t \right) = \sum_{n=0}^N \frac{(-N)_n}{(\beta + 1)_n n!} Q_n(x; \alpha, \beta, N) t^n. \tag{9.5.11}$$

$$\begin{aligned}
& {}_2F_0 \left(\begin{matrix} -x, -x + \beta + N + 1 \\ - \end{matrix}; -t \right) {}_2F_0 \left(\begin{matrix} x - N, x + \alpha + 1 \\ - \end{matrix}; t \right) \\
& = \sum_{n=0}^N \frac{(-N)_n (\alpha + 1)_n}{n!} Q_n(x; \alpha, \beta, N) t^n.
\end{aligned} \tag{9.5.12}$$

$$\begin{aligned}
& \left[(1 - t)^{-\alpha - \beta - 1} {}_3F_2 \left(\begin{matrix} \frac{1}{2}(\alpha + \beta + 1), \frac{1}{2}(\alpha + \beta + 2), -x \\ \alpha + 1, -N \end{matrix}; -\frac{4t}{(1 - t)^2} \right) \right]_N \\
& = \sum_{n=0}^N \frac{(\alpha + \beta + 1)_n}{n!} Q_n(x; \alpha, \beta, N) t^n.
\end{aligned} \tag{9.5.13}$$

Limit relations**Racah \rightarrow Hahn**

If we take $\gamma + 1 = -N$ and let $\delta \rightarrow \infty$ in the definition (9.2.1) of the Racah polynomials, we obtain the Hahn polynomials. Hence

$$\lim_{\delta \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, -N - 1, \delta) = Q_n(x; \alpha, \beta, N).$$

And if we take $\delta = -\beta - N - 1$ and let $\gamma \rightarrow \infty$ in the definition (9.2.1) of the Racah polynomials, we also obtain the Hahn polynomials:

$$\lim_{\gamma \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, \gamma, -\beta - N - 1) = Q_n(x; \alpha, \beta, N).$$

Another way to do this is to take $\alpha + 1 = -N$ and $\beta \rightarrow \beta + \gamma + N + 1$ in the definition (9.2.1) of the Racah polynomials and then take the limit $\delta \rightarrow \infty$. In that case we obtain the Hahn polynomials in the following way:

$$\lim_{\delta \rightarrow \infty} R_n(\lambda(x); -N-1, \beta + \gamma + N + 1, \gamma, \delta) = Q_n(x; \gamma, \beta, N).$$

Hahn \rightarrow Jacobi

To find the Jacobi polynomials given by (9.8.1) from the Hahn polynomials we take $x \rightarrow Nx$ and let $N \rightarrow \infty$. In fact we have

$$\lim_{N \rightarrow \infty} Q_n(Nx; \alpha, \beta, N) = \frac{P_n^{(\alpha, \beta)}(1 - 2x)}{P_n^{(\alpha, \beta)}(1)}. \quad (9.5.14)$$

Hahn \rightarrow Meixner

The Meixner polynomials given by (9.10.1) can be obtained from the Hahn polynomials by taking $\alpha = b - 1$, $\beta = N(1 - c)c^{-1}$ and letting $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} Q_n(x; b-1, N(1-c)c^{-1}, N) = M_n(x; b, c). \quad (9.5.15)$$

Hahn \rightarrow Krawtchouk

The Krawtchouk polynomials given by (9.11.1) are obtained from the Hahn polynomials if we take $\alpha = pt$ and $\beta = (1-p)t$ and let $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} Q_n(x; pt, (1-p)t, N) = K_n(x; p, N). \quad (9.5.16)$$

Remark

If we interchange the role of x and n in (9.5.1) we obtain the dual Hahn polynomials given by (9.6.1).

Since

$$Q_n(x; \alpha, \beta, N) = R_x(\lambda(n); \alpha, \beta, N)$$

we obtain the dual orthogonality relation for the Hahn polynomials from the orthogonality relation (9.6.2) of the dual Hahn polynomials:

$$\begin{aligned} & \sum_{n=0}^N \frac{(2n+\alpha+\beta+1)(\alpha+1)_n(-N)_n N!}{(-1)^n(n+\alpha+\beta+1)_{N+1}(\beta+1)_n n!} Q_n(x; \alpha, \beta, N) Q_n(y; \alpha, \beta, N) \\ &= \frac{\delta_{xy}}{\binom{\alpha+x}{x} \binom{\beta+N-x}{N-x}}, \quad x, y \in \{0, 1, 2, \dots, N\}. \end{aligned}$$

References

[illegible]

9.6 Dual Hahn

Hypergeometric representation

$$R_n(\lambda(x); \gamma, \delta, N) = {}_3F_2 \left(\begin{matrix} -n, -x, x + \gamma + \delta + 1 \\ \gamma + 1, -N \end{matrix}; 1 \right), \quad n = 0, 1, 2, \dots, N, \quad (9.6.1)$$

where

$$\lambda(x) = x(x + \gamma + \delta + 1).$$

Orthogonality relation

For $\gamma > -1$ and $\delta > -1$, or for $\gamma < -N$ and $\delta < -N$, we have

$$\begin{aligned} \sum_{x=0}^N \frac{(2x + \gamma + \delta + 1)(\gamma + 1)_x (-N)_x N!}{(-1)^x (x + \gamma + \delta + 1)_{N+1} (\delta + 1)_x x!} R_m(\lambda(x); \gamma, \delta, N) R_n(\lambda(x); \gamma, \delta, N) \\ = \frac{\delta_{mn}}{\binom{\gamma + n}{n} \binom{\delta + N - n}{N - n}}. \end{aligned} \quad (9.6.2)$$

Recurrence relation

$$\lambda(x) R_n(\lambda(x)) = A_n R_{n+1}(\lambda(x)) - (A_n + C_n) R_n(\lambda(x)) + C_n R_{n-1}(\lambda(x)), \quad (9.6.3)$$

where

$$R_n(\lambda(x)) := R_n(\lambda(x); \gamma, \delta, N)$$

and

$$\begin{cases} A_n = (n + \gamma + 1)(n - N) \\ C_n = n(n - \delta - N - 1). \end{cases}$$

Normalized recurrence relation

$$x p_n(x) = p_{n+1}(x) - (A_n + C_n) p_n(x) + A_{n-1} C_n p_{n-1}(x), \quad (9.6.4)$$

where

$$R_n(\lambda(x); \gamma, \delta, N) = \frac{1}{(\gamma + 1)_n (-N)_n} p_n(\lambda(x)).$$

Difference equation

$$-ny(x) = B(x)y(x+1) - [B(x) + D(x)]y(x) + D(x)y(x-1), \quad (9.6.5)$$

where

$$y(x) = R_n(\lambda(x); \gamma, \delta, N)$$

and

$$\begin{cases} B(x) = \frac{(x + \gamma + 1)(x + \gamma + \delta + 1)(N - x)}{(2x + \gamma + \delta + 1)(2x + \gamma + \delta + 2)} \\ D(x) = \frac{x(x + \gamma + \delta + N + 1)(x + \delta)}{(2x + \gamma + \delta)(2x + \gamma + \delta + 1)}. \end{cases}$$

Forward shift operator

$$\begin{aligned}
& R_n(\lambda(x+1); \gamma, \delta, N) - R_n(\lambda(x); \gamma, \delta, N) \\
&= -\frac{n(2x + \gamma + \delta + 2)}{(\gamma + 1)N} R_{n-1}(\lambda(x); \gamma + 1, \delta, N - 1)
\end{aligned} \tag{9.6.6}$$

or equivalently

$$\frac{\Delta R_n(\lambda(x); \gamma, \delta, N)}{\Delta \lambda(x)} = -\frac{n}{(\gamma + 1)N} R_{n-1}(\lambda(x); \gamma + 1, \delta, N - 1). \tag{9.6.7}$$

Backward shift operator

$$\begin{aligned}
& (x + \gamma)(x + \gamma + \delta)(N + 1 - x)R_n(\lambda(x); \gamma, \delta, N) \\
& \quad - x(x + \gamma + \delta + N + 1)(x + \delta)R_n(\lambda(x - 1); \gamma, \delta, N) \\
&= \gamma(N + 1)(2x + \gamma + \delta)R_{n+1}(\lambda(x); \gamma - 1, \delta, N + 1)
\end{aligned} \tag{9.6.8}$$

or equivalently

$$\begin{aligned}
& \frac{\nabla [\omega(x; \gamma, \delta, N)R_n(\lambda(x); \gamma, \delta, N)]}{\nabla \lambda(x)} \\
&= \frac{1}{\gamma + \delta} \omega(x; \gamma - 1, \delta, N + 1)R_{n+1}(\lambda(x); \gamma - 1, \delta, N + 1),
\end{aligned} \tag{9.6.9}$$

where

$$\omega(x; \gamma, \delta, N) = \frac{(-1)^x (\gamma + 1)_x (\gamma + \delta + 1)_x (-N)_x}{(\gamma + \delta + N + 2)_x (\delta + 1)_x x!}.$$

Rodrigues-type formula

$$\omega(x; \gamma, \delta, N) R_n(\lambda(x); \gamma, \delta, N) = (\gamma + \delta + 1)_n (\nabla_\lambda)^n [\omega(x; \gamma + n, \delta, N - n)], \quad (9.6.10)$$

where

$$\nabla_\lambda := \frac{\nabla}{\nabla \lambda(x)}.$$

Generating functions

For $x = 0, 1, 2, \dots, N$ we have

$$(1-t)^{N-x} {}_2F_1 \left(\begin{matrix} -x, -x-\delta \\ \gamma+1 \end{matrix}; t \right) = \sum_{n=0}^N \frac{(-N)_n}{n!} R_n(\lambda(x); \gamma, \delta, N) t^n. \quad (9.6.11)$$

$$\begin{aligned} (1-t)^x {}_2F_1 \left(\begin{matrix} x-N, x+\gamma+1 \\ -\delta-N \end{matrix}; t \right) \\ = \sum_{n=0}^N \frac{(\gamma+1)_n (-N)_n}{(-\delta-N)_n n!} R_n(\lambda(x); \gamma, \delta, N) t^n. \end{aligned} \quad (9.6.12)$$

$$\left[e^t {}_2F_2 \left(\begin{matrix} -x, x+\gamma+\delta+1 \\ \gamma+1, -N \end{matrix}; -t \right) \right]_N = \sum_{n=0}^N \frac{R_n(\lambda(x); \gamma, \delta, N)}{n!} t^n. \quad (9.6.13)$$

$$\begin{aligned} \left[(1-t)^{-\varepsilon} {}_3F_2 \left(\begin{matrix} \varepsilon, -x, x+\gamma+\delta+1 \\ \gamma+1, -N \end{matrix}; \frac{t}{t-1} \right) \right]_N \\ = \sum_{n=0}^N \frac{(\varepsilon)_n}{n!} R_n(\lambda(x); \gamma, \delta, N) t^n, \quad \varepsilon \text{ arbitrary.} \end{aligned} \quad (9.6.14)$$

Limit relations**Racah \rightarrow Dual Hahn**

If we take $\alpha + 1 = -N$ and let $\beta \rightarrow \infty$ in the definition (9.2.1) of the Racah polynomials, then we obtain the dual Hahn polynomials:

$$\lim_{\beta \rightarrow \infty} R_n(\lambda(x); -N-1, \beta, \gamma, \delta) = R_n(\lambda(x); \gamma, \delta, N).$$

And if we take $\beta = -\delta - N - 1$ and let $\alpha \rightarrow \infty$ in the definition (9.2.1) of the Racah polynomials, then we also obtain the dual Hahn polynomials:

$$\lim_{\alpha \rightarrow \infty} R_n(\lambda(x); \alpha, -\delta - N - 1, \gamma, \delta) = R_n(\lambda(x); \gamma, \delta, N).$$

Finally, if we take $\gamma + 1 = -N$ and $\delta \rightarrow \alpha + \delta + N + 1$ in the definition (9.2.1) of the Racah polynomials and take the limit $\beta \rightarrow \infty$ we find the dual Hahn polynomials in the following way:

$$\lim_{\beta \rightarrow \infty} R_n(\lambda(x); \alpha, \beta, -N-1, \alpha + \delta + N + 1) = R_n(\lambda(x); \alpha, \delta, N).$$

Dual Hahn \rightarrow Meixner

The Meixner polynomials given by (9.10.1) are obtained from the dual Hahn polynomials if we take $\gamma = \beta - 1$ and $\delta = N(1-c)c^{-1}$ and let $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} R_n(\lambda(x); \beta - 1, N(1-c)c^{-1}, N) = M_n(x; \beta, c). \quad (9.6.15)$$

Dual Hahn \rightarrow Krawtchouk

The Krawtchouk polynomials given by (9.11.1) can be obtained from the dual Hahn polynomials by setting $\gamma = pt$, $\delta = (1-p)t$ and letting $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} R_n(\lambda(x); pt, (1-p)t, N) = K_n(x; p, N). \quad (9.6.16)$$

Remark

If we interchange the role of x and n in the definition (9.6.1) of the dual Hahn polynomials we obtain the Hahn polynomials given by (9.5.1).

Since

$$R_n(\lambda(x); \gamma, \delta, N) = Q_x(n; \gamma, \delta, N)$$

we obtain the dual orthogonality relation for the dual Hahn polynomials from the orthogonality relation (9.5.2) for the Hahn polynomials:

$$\begin{aligned} & \sum_{n=0}^N \binom{\gamma+n}{n} \binom{\delta+N-n}{N-n} R_n(\lambda(x); \gamma, \delta, N) R_n(\lambda(y); \gamma, \delta, N) \\ &= \frac{(-1)^x (x+\gamma+\delta+1)_{N+1} (\delta+1)_x x!}{(2x+\gamma+\delta+1)(\gamma+1)_x (-N)_x N!} \delta_{xy}, \quad x, y \in \{0, 1, 2, \dots, N\}. \end{aligned}$$

References

[?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?].

9.7 Meixner-Pollaczek**Hypergeometric representation**

$$P_n^{(\lambda)}(x; \phi) = \frac{(2\lambda)_n}{n!} e^{in\phi} {}_2F_1 \left(\begin{matrix} -n, \lambda + ix \\ 2\lambda \end{matrix}; 1 - e^{-2i\phi} \right). \quad (9.7.1)$$

Orthogonality relation

$$\frac{1}{2\pi} \int_{-\infty}^{\infty} e^{(2\phi-\pi)x} |\Gamma(\lambda + ix)|^2 P_m^{(\lambda)}(x; \phi) P_n^{(\lambda)}(x; \phi) dx = \frac{\Gamma(n+2\lambda)}{(2\sin\phi)^{2\lambda} n!} \delta_{mn}, \quad (9.7.2)$$

Recurrence relation

$$\begin{aligned} & (n+1)P_{n+1}^{(\lambda)}(x; \phi) - 2[x\sin\phi + (n+\lambda)\cos\phi]P_n^{(\lambda)}(x; \phi) \\ & + (n+2\lambda-1)P_{n-1}^{(\lambda)}(x; \phi) = 0. \end{aligned} \quad (9.7.3)$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) - \left(\frac{n+\lambda}{\tan\phi} \right) p_n(x) + \frac{n(n+2\lambda-1)}{4\sin^2\phi} p_{n-1}(x), \quad (9.7.4)$$

where

$$P_n^{(\lambda)}(x; \phi) = \frac{(2\sin\phi)^n}{n!} p_n(x).$$

Difference equation

$$\begin{aligned} e^{i\phi}(\lambda - ix)y(x+i) + 2i[x \cos \phi - (n+\lambda) \sin \phi]y(x) \\ - e^{-i\phi}(\lambda + ix)y(x-i) = 0, \quad y(x) = P_n^{(\lambda)}(x; \phi). \end{aligned} \quad (9.7.5)$$

Forward shift operator

$$P_n^{(\lambda)}(x + \tfrac{1}{2}i; \phi) - P_n^{(\lambda)}(x - \tfrac{1}{2}i; \phi) = (e^{i\phi} - e^{-i\phi})P_{n-1}^{(\lambda+\frac{1}{2})}(x; \phi) \quad (9.7.6)$$

or equivalently

$$\frac{\delta P_n^{(\lambda)}(x; \phi)}{\delta x} = 2 \sin \phi P_{n-1}^{(\lambda+\frac{1}{2})}(x; \phi). \quad (9.7.7)$$

Backward shift operator

$$\begin{aligned} e^{i\phi}(\lambda - \tfrac{1}{2} - ix)P_n^{(\lambda)}(x + \tfrac{1}{2}i; \phi) + e^{-i\phi}(\lambda - \tfrac{1}{2} + ix)P_n^{(\lambda)}(x - \tfrac{1}{2}i; \phi) \\ = (n+1)P_{n+1}^{(\lambda-\frac{1}{2})}(x; \phi) \end{aligned} \quad (9.7.8)$$

or equivalently

$$\frac{\delta \left[\omega(x; \lambda, \phi) P_n^{(\lambda)}(x; \phi) \right]}{\delta x} = -(n+1) \omega(x; \lambda - \tfrac{1}{2}, \phi) P_{n+1}^{(\lambda-\frac{1}{2})}(x; \phi), \quad (9.7.9)$$

where

$$\omega(x; \lambda, \phi) = \Gamma(\lambda + ix) \Gamma(\lambda - ix) e^{(2\phi - \pi)x}.$$

Rodrigues-type formula

$$\omega(x; \lambda, \phi) P_n^{(\lambda)}(x; \phi) = \frac{(-1)^n}{n!} \left(\frac{\delta}{\delta x} \right)^n \left[\omega(x; \lambda + \tfrac{1}{2}n, \phi) \right]. \quad (9.7.10)$$

Generating functions

$$(1 - e^{i\phi}t)^{-\lambda+ix}(1 - e^{-i\phi}t)^{-\lambda-ix} = \sum_{n=0}^{\infty} P_n^{(\lambda)}(x; \phi) t^n. \quad (9.7.11)$$

$$e^t {}_1F_1 \left(\begin{matrix} \lambda + ix \\ 2\lambda \end{matrix}; (e^{-2i\phi} - 1)t \right) = \sum_{n=0}^{\infty} \frac{P_n^{(\lambda)}(x; \phi)}{(2\lambda)_n e^{in\phi}} t^n. \quad (9.7.12)$$

$$\begin{aligned} & (1-t)^{-\gamma} {}_2F_1 \left(\begin{matrix} \gamma, \lambda + ix \\ 2\lambda \end{matrix}; \frac{(1 - e^{-2i\phi})t}{t-1} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(2\lambda)_n} \frac{P_n^{(\lambda)}(x; \phi)}{e^{in\phi}} t^n, \quad \gamma \text{ arbitrary.} \end{aligned} \quad (9.7.13)$$

Limit relations**Continuous dual Hahn \rightarrow Meixner-Pollaczek**

The Meixner-Pollaczek polynomials can be obtained from the continuous dual Hahn polynomials given by (9.3.1) by the substitutions $x \rightarrow x - t$, $a = \lambda + it$, $b = \lambda - it$ and $c = t \cot \phi$ and the limit $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} \frac{S_n((x-t)^2; \lambda + it, \lambda - it, t \cot \phi)}{t^n n!} = \frac{P_n^{(\lambda)}(x; \phi)}{(\sin \phi)^n}.$$

Continuous Hahn \rightarrow Meixner-Pollaczek

By setting $x \rightarrow x + t$, $a = \lambda - it$, $c = \lambda + it$ and $b = d = t \tan \phi$ in the definition (9.4.1) of the continuous Hahn polynomials and taking the limit $t \rightarrow \infty$ we obtain the Meixner-Pollaczek polynomials:

$$\lim_{t \rightarrow \infty} \frac{p_n(x+t; \lambda - it, t \tan \phi, \lambda + it, t \tan \phi)}{t^n n!} = \frac{P_n^{(\lambda)}(x; \phi)}{(\cos \phi)^n}.$$

$$\begin{aligned} & \int_1^\infty (x+1)^\alpha (x-1)^\beta P_m^{(\alpha,\beta)}(-x) P_n^{(\alpha,\beta)}(-x) dx \\ &= -\frac{2^{\alpha+\beta+1}}{2n+\alpha+\beta+1} \frac{\Gamma(-n-\alpha-\beta)\Gamma(n+\alpha+\beta+1)}{\Gamma(-n-\alpha)n!} \delta_{mn}. \end{aligned} \quad (9.8.3)$$

Recurrence relation

$$\begin{aligned}
xP_n^{(\alpha,\beta)}(x) &= \frac{2(n+1)(n+\alpha+\beta+1)}{(2n+\alpha+\beta+1)(2n+\alpha+\beta+2)}P_{n+1}^{(\alpha,\beta)}(x) \\
&\quad + \frac{\beta^2 - \alpha^2}{(2n+\alpha+\beta)(2n+\alpha+\beta+2)}P_n^{(\alpha,\beta)}(x) \\
&\quad + \frac{2(n+\alpha)(n+\beta)}{(2n+\alpha+\beta)(2n+\alpha+\beta+1)}P_{n-1}^{(\alpha,\beta)}(x).
\end{aligned} \tag{9.8.4}$$

Normalized recurrence relation

$$\begin{aligned}
xp_n(x) &= p_{n+1}(x) + \frac{\beta^2 - \alpha^2}{(2n+\alpha+\beta)(2n+\alpha+\beta+2)}p_n(x) \\
&\quad + \frac{4n(n+\alpha)(n+\beta)(n+\alpha+\beta)}{(2n+\alpha+\beta-1)(2n+\alpha+\beta)^2(2n+\alpha+\beta+1)}p_{n-1}(x)
\end{aligned} \tag{9.8.5}$$

where

$$P_n^{(\alpha,\beta)}(x) = \frac{(n+\alpha+\beta+1)_n}{2^n n!} p_n(x).$$

Differential equation

$$\begin{aligned}
(1-x^2)y''(x) + [\beta - \alpha - (\alpha + \beta + 2)x]y'(x) \\
+ n(n+\alpha+\beta+1)y(x) = 0, \quad y(x) = P_n^{(\alpha,\beta)}(x).
\end{aligned} \tag{9.8.6}$$

Forward shift operator

$$\frac{d}{dx}P_n^{(\alpha,\beta)}(x) = \frac{n+\alpha+\beta+1}{2}P_{n-1}^{(\alpha+1,\beta+1)}(x). \tag{9.8.7}$$

Backward shift operator

$$\begin{aligned}
(1-x^2)\frac{d}{dx}P_n^{(\alpha,\beta)}(x) + [(\beta - \alpha) - (\alpha + \beta)x]P_n^{(\alpha,\beta)}(x) \\
= -2(n+1)P_{n+1}^{(\alpha-1,\beta-1)}(x)
\end{aligned} \tag{9.8.8}$$

or equivalently

$$\begin{aligned}
\frac{d}{dx} \left[(1-x)^\alpha (1+x)^\beta P_n^{(\alpha,\beta)}(x) \right] \\
= -2(n+1)(1-x)^{\alpha-1}(1+x)^{\beta-1}P_{n+1}^{(\alpha-1,\beta-1)}(x).
\end{aligned} \tag{9.8.9}$$

Rodrigues-type formula

$$(1-x)^\alpha (1+x)^\beta P_n^{(\alpha,\beta)}(x) = \frac{(-1)^n}{2^n n!} \left(\frac{d}{dx} \right)^n \left[(1-x)^{n+\alpha} (1+x)^{n+\beta} \right]. \tag{9.8.10}$$

Generating functions

$$\frac{2^{\alpha+\beta}}{R(1+R-t)^{\alpha}(1+R+t)^{\beta}} = \sum_{n=0}^{\infty} P_n^{(\alpha,\beta)}(x)t^n, \quad R = \sqrt{1-2xt+t^2}. \quad (9.8.11)$$

$$\begin{aligned} & {}_0F_1\left(\begin{matrix} - \\ \alpha+1 \end{matrix}; \frac{(x-1)t}{2}\right) {}_0F_1\left(\begin{matrix} - \\ \beta+1 \end{matrix}; \frac{(x+1)t}{2}\right) \\ &= \sum_{n=0}^{\infty} \frac{P_n^{(\alpha,\beta)}(x)}{(\alpha+1)_n(\beta+1)_n} t^n. \end{aligned} \quad (9.8.12)$$

$$\begin{aligned} & (1-t)^{-\alpha-\beta-1} {}_2F_1\left(\begin{matrix} \frac{1}{2}(\alpha+\beta+1), \frac{1}{2}(\alpha+\beta+2) \\ \alpha+1 \end{matrix}; \frac{2(x-1)t}{(1-t)^2}\right) \\ &= \sum_{n=0}^{\infty} \frac{(\alpha+\beta+1)_n}{(\alpha+1)_n} P_n^{(\alpha,\beta)}(x)t^n. \end{aligned} \quad (9.8.13)$$

$$\begin{aligned} & (1+t)^{-\alpha-\beta-1} {}_2F_1\left(\begin{matrix} \frac{1}{2}(\alpha+\beta+1), \frac{1}{2}(\alpha+\beta+2) \\ \beta+1 \end{matrix}; \frac{2(x+1)t}{(1+t)^2}\right) \\ &= \sum_{n=0}^{\infty} \frac{(\alpha+\beta+1)_n}{(\beta+1)_n} P_n^{(\alpha,\beta)}(x)t^n. \end{aligned} \quad (9.8.14)$$

$$\begin{aligned} & {}_2F_1\left(\begin{matrix} \gamma, \alpha+\beta+1-\gamma \\ \alpha+1 \end{matrix}; \frac{1-R-t}{2}\right) {}_2F_1\left(\begin{matrix} \gamma, \alpha+\beta+1-\gamma \\ \beta+1 \end{matrix}; \frac{1-R+t}{2}\right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n(\alpha+\beta+1-\gamma)_n}{(\alpha+1)_n(\beta+1)_n} P_n^{(\alpha,\beta)}(x)t^n, \quad R = \sqrt{1-2xt+t^2} \end{aligned} \quad (9.8.15)$$

with γ arbitrary.

Limit relations**Wilson \rightarrow Jacobi**

The Jacobi polynomials can be found from the Wilson polynomials given by (9.1.1) by substituting $a = b = \frac{1}{2}(\alpha+1)$, $c = \frac{1}{2}(\beta+1) + it$, $d = \frac{1}{2}(\beta+1) - it$ and $x \rightarrow t\sqrt{\frac{1}{2}(1-x)}$ in the definition (9.1.1) of the Wilson polynomials and taking the limit $t \rightarrow \infty$. In fact we have

$$\lim_{t \rightarrow \infty} \frac{W_n(\frac{1}{2}(1-x)t^2; \frac{1}{2}(\alpha+1), \frac{1}{2}(\alpha+1), \frac{1}{2}(\beta+1) + it, \frac{1}{2}(\beta+1) - it)}{t^{2n}n!} = P_n^{(\alpha,\beta)}(x).$$

Continuous Hahn \rightarrow Jacobi

The Jacobi polynomials follow from the continuous Hahn polynomials given by (9.4.1) by using the substitution $x \rightarrow \frac{1}{2}xt$, $a = \frac{1}{2}(\alpha+1-it)$, $b = \frac{1}{2}(\beta+1+it)$, $c = \frac{1}{2}(\alpha+1+it)$ and $d = \frac{1}{2}(\beta+1-it)$ in (9.4.1), division by t^n and the limit $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} \frac{p_n(\frac{1}{2}xt; \frac{1}{2}(\alpha+1-it), \frac{1}{2}(\beta+1+it), \frac{1}{2}(\alpha+1+it), \frac{1}{2}(\beta+1-it))}{t^n} = P_n^{(\alpha,\beta)}(x).$$

Hahn \rightarrow Jacobi

To find the Jacobi polynomials from the Hahn polynomials given by (9.5.1) we take $x \rightarrow Nx$ in (9.5.1) and let $N \rightarrow \infty$. In fact we have

$$\lim_{N \rightarrow \infty} Q_n(Nx; \alpha, \beta, N) = \frac{P_n^{(\alpha, \beta)}(1 - 2x)}{P_n^{(\alpha, \beta)}(1)}.$$

Jacobi \rightarrow Laguerre

The Laguerre polynomials given by (9.12.1) can be obtained from the Jacobi polynomials by setting $x \rightarrow 1 - 2\beta^{-1}x$ and then the limit $\beta \rightarrow \infty$:

$$\lim_{\beta \rightarrow \infty} P_n^{(\alpha, \beta)}(1 - 2\beta^{-1}x) = L_n^{(\alpha)}(x). \quad (9.8.16)$$

Jacobi \rightarrow Bessel

The Bessel polynomials given by (9.13.1) are obtained from the Jacobi polynomials if we take $\beta = a - \alpha$ and let $\alpha \rightarrow -\infty$:

$$\lim_{\alpha \rightarrow -\infty} \frac{P_n^{(\alpha, a-\alpha)}(1 + \alpha x)}{P_n^{(\alpha, a-\alpha)}(1)} = y_n(x; a). \quad (9.8.17)$$

Jacobi \rightarrow Hermite

The Hermite polynomials given by (9.15.1) follow from the Jacobi polynomials by taking $\beta = \alpha$ and letting $\alpha \rightarrow \infty$ in the following way:

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\frac{1}{2}n} P_n^{(\alpha, \alpha)}(\alpha^{-\frac{1}{2}}x) = \frac{H_n(x)}{2^n n!}. \quad (9.8.18)$$

Remarks

The definition (9.8.1) of the Jacobi polynomials can also be written as:

$$P_n^{(\alpha, \beta)}(x) = \frac{1}{n!} \sum_{k=0}^n \frac{(-n)_k}{k!} (n + \alpha + \beta + 1)_k (\alpha + k + 1)_{n-k} \left(\frac{1-x}{2} \right)^k.$$

In this way the Jacobi polynomials can also be seen as polynomials in the parameters α and β . Therefore they can be defined for all α and β . Then we have the following connection with the Meixner polynomials given by (9.10.1):

$$\frac{(\beta)_n}{n!} M_n(x; \beta, c) = P_n^{(\beta-1, -n-\beta-x)}((2-c)c^{-1}).$$

The Jacobi polynomials are related to the pseudo Jacobi polynomials defined by (9.9.1) in the following way:

$$P_n(x; \nu, N) = \frac{(-2i)^n n!}{(n - 2N - 1)_n} P_n^{(-N-1+i\nu, -N-1-i\nu)}(ix).$$

The Jacobi polynomials are also related to the Gegenbauer (or ultraspherical) polynomials given by (9.8.19) by the quadratic transformations:

$$C_{2n}^{(\lambda)}(x) = \frac{(\lambda)_n}{(\frac{1}{2})_n} P_n^{(\lambda-\frac{1}{2}, -\frac{1}{2})}(2x^2 - 1)$$

and

$$C_{2n+1}^{(\lambda)}(x) = \frac{(\lambda)_{n+1}}{(\frac{1}{2})_{n+1}} x P_n^{(\lambda-\frac{1}{2}, \frac{1}{2})}(2x^2 - 1).$$

$$\frac{d}{dx}C_n^{(\lambda)}(x) = 2\lambda C_{n-1}^{(\lambda+1)}(x). \quad (9.8.24)$$

Backward shift operator

$$(1-x^2)\frac{d}{dx}C_n^{(\lambda)}(x) + (1-2\lambda)x C_n^{(\lambda)}(x) = -\frac{(n+1)(2\lambda+n-1)}{2(\lambda-1)}C_{n+1}^{(\lambda-1)}(x) \quad (9.8.25)$$

or equivalently

$$\begin{aligned} & \frac{d}{dx} \left[(1-x^2)^{\lambda-\frac{1}{2}} C_n^{(\lambda)}(x) \right] \\ &= -\frac{(n+1)(2\lambda+n-1)}{2(\lambda-1)} (1-x^2)^{\lambda-\frac{3}{2}} C_{n+1}^{(\lambda-1)}(x). \end{aligned} \quad (9.8.26)$$

Rodrigues-type formula

$$(1-x^2)^{\lambda-\frac{1}{2}} C_n^{(\lambda)}(x) = \frac{(2\lambda)_n (-1)^n}{(\lambda+\frac{1}{2})_n 2^n n!} \left(\frac{d}{dx} \right)^n \left[(1-x^2)^{\lambda+n-\frac{1}{2}} \right]. \quad (9.8.27)$$

Generating functions

$$(1-2xt+t^2)^{-\lambda} = \sum_{n=0}^{\infty} C_n^{(\lambda)}(x) t^n. \quad (9.8.28)$$

$$R^{-1} \left(\frac{1+R-xt}{2} \right)^{\frac{1}{2}-\lambda} = \sum_{n=0}^{\infty} \frac{(\lambda+\frac{1}{2})_n}{(2\lambda)_n} C_n^{(\lambda)}(x) t^n, \quad R = \sqrt{1-2xt+t^2}. \quad (9.8.29)$$

$${}_0F_1 \left(\begin{matrix} - \\ \lambda+\frac{1}{2} \end{matrix}; \frac{(x-1)t}{2} \right) {}_0F_1 \left(\begin{matrix} - \\ \lambda+\frac{1}{2} \end{matrix}; \frac{(x+1)t}{2} \right) = \sum_{n=0}^{\infty} \frac{C_n^{(\lambda)}(x)}{(2\lambda)_n (\lambda+\frac{1}{2})_n} t^n. \quad (9.8.30)$$

$$e^{xt} {}_0F_1 \left(\begin{matrix} - \\ \lambda+\frac{1}{2} \end{matrix}; \frac{(x^2-1)t^2}{4} \right) = \sum_{n=0}^{\infty} \frac{C_n^{(\lambda)}(x)}{(2\lambda)_n} t^n. \quad (9.8.31)$$

$$\begin{aligned} & {}_2F_1 \left(\begin{matrix} \gamma, 2\lambda-\gamma \\ \lambda+\frac{1}{2} \end{matrix}; \frac{1-R-t}{2} \right) {}_2F_1 \left(\begin{matrix} \gamma, 2\lambda-\gamma \\ \lambda+\frac{1}{2} \end{matrix}; \frac{1-R+t}{2} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n (2\lambda-\gamma)_n}{(2\lambda)_n (\lambda+\frac{1}{2})_n} C_n^{(\lambda)}(x) t^n, \quad R = \sqrt{1-2xt+t^2}, \quad \gamma \text{ arbitrary}. \end{aligned} \quad (9.8.32)$$

$$\begin{aligned} & (1-xt)^{-\gamma} {}_2F_1 \left(\begin{matrix} \frac{1}{2}\gamma, \frac{1}{2}\gamma+\frac{1}{2} \\ \lambda+\frac{1}{2} \end{matrix}; \frac{(x^2-1)t^2}{(1-xt)^2} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(2\lambda)_n} C_n^{(\lambda)}(x) t^n, \quad \gamma \text{ arbitrary}. \end{aligned} \quad (9.8.33)$$

Limit relation**Gegenbauer / Ultraspherical \rightarrow Hermite**

The Hermite polynomials given by (9.15.1) follow from the Gegenbauer (or ultraspherical) polynomials by taking $\lambda = \alpha + \frac{1}{2}$ and letting $\alpha \rightarrow \infty$ in the following way:

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\frac{1}{2}n} C_n^{(\alpha+\frac{1}{2})}(\alpha^{-\frac{1}{2}}x) = \frac{H_n(x)}{n!}. \quad (9.8.34)$$

$$2xU_n(x) = U_{n+1}(x) + U_{n-1}(x), \quad U_0(x) = 1 \quad \text{and} \quad U_1(x) = 2x. \quad (9.8.40)$$

Normalized recurrence relations

$$xp_n(x) = p_{n+1}(x) + \frac{1}{4}p_{n-1}(x), \quad (9.8.41)$$

where

$$T_1(x) = p_1(x) = x \quad \text{and} \quad T_n(x) = 2^n p_n(x), \quad n \neq 1.$$

$$xp_n(x) = p_{n+1}(x) + \frac{1}{4}p_{n-1}(x), \quad (9.8.42)$$

where

$$U_n(x) = 2^n p_n(x).$$

Differential equations

$$(1-x^2)y''(x) - xy'(x) + n^2y(x) = 0, \quad y(x) = T_n(x). \quad (9.8.43)$$

$$(1-x^2)y''(x) - 3xy'(x) + n(n+2)y(x) = 0, \quad y(x) = U_n(x). \quad (9.8.44)$$

Forward shift operator

$$\frac{d}{dx}T_n(x) = nU_{n-1}(x). \quad (9.8.45)$$

Backward shift operator

$$(1-x^2)\frac{d}{dx}U_n(x) - xU_n(x) = -(n+1)T_{n+1}(x) \quad (9.8.46)$$

or equivalently

$$\frac{d}{dx} \left[(1-x^2)^{\frac{1}{2}} U_n(x) \right] = -(n+1) (1-x^2)^{-\frac{1}{2}} T_{n+1}(x). \quad (9.8.47)$$

Rodrigues-type formulas

$$(1-x^2)^{-\frac{1}{2}} T_n(x) = \frac{(-1)^n}{(\frac{1}{2})_n 2^n} \left(\frac{d}{dx} \right)^n \left[(1-x^2)^{n-\frac{1}{2}} \right]. \quad (9.8.48)$$

$$(1-x^2)^{\frac{1}{2}} U_n(x) = \frac{(n+1)(-1)^n}{(\frac{3}{2})_n 2^n} \left(\frac{d}{dx} \right)^n \left[(1-x^2)^{n+\frac{1}{2}} \right]. \quad (9.8.49)$$

Generating functions

$$\frac{1-xt}{1-2xt+t^2} = \sum_{n=0}^{\infty} T_n(x)t^n. \quad (9.8.50)$$

$$R^{-1} \sqrt{\frac{1}{2}(1+R-xt)} = \sum_{n=0}^{\infty} \frac{(\frac{1}{2})_n}{n!} T_n(x)t^n, \quad R = \sqrt{1-2xt+t^2}. \quad (9.8.51)$$

$${}_0F_1 \left(\frac{-}{\frac{1}{2}}; \frac{(x-1)t}{2} \right) {}_0F_1 \left(\frac{-}{\frac{1}{2}}; \frac{(x+1)t}{2} \right) = \sum_{n=0}^{\infty} \frac{T_n(x)}{(\frac{1}{2})_n n!} t^n. \quad (9.8.52)$$

$$e^{xt} {}_0F_1 \left(\frac{-}{\frac{1}{2}}; \frac{(x^2-1)t^2}{4} \right) = \sum_{n=0}^{\infty} \frac{T_n(x)}{n!} t^n. \quad (9.8.53)$$

$$\begin{aligned}
& {}_2F_1\left(\gamma, -\gamma; \frac{1-R-t}{2}; \frac{1-R+t}{2}\right) {}_2F_1\left(\gamma, -\gamma; \frac{1-R+t}{2}; \frac{1-R-t}{2}\right) \\
&= \sum_{n=0}^{\infty} \frac{(\gamma)_n (-\gamma)_n}{\left(\frac{1}{2}\right)_n n!} T_n(x) t^n, \quad R = \sqrt{1-2xt+t^2}, \quad \gamma \text{ arbitrary.}
\end{aligned} \tag{9.8.54}$$

$$\begin{aligned}
& (1-xt)^{-\gamma} {}_2F_1\left(\frac{1}{2}\gamma, \frac{1}{2}\gamma + \frac{1}{2}; \frac{(x^2-1)t^2}{(1-xt)^2}; \frac{1}{2}\right) \\
&= \sum_{n=0}^{\infty} \frac{(\gamma)_n}{n!} T_n(x) t^n, \quad \gamma \text{ arbitrary.}
\end{aligned} \tag{9.8.55}$$

$$\frac{1}{1-2xt+t^2} = \sum_{n=0}^{\infty} U_n(x) t^n. \tag{9.8.56}$$

$$\frac{1}{R\sqrt{\frac{1}{2}(1+R-xt)}} = \sum_{n=0}^{\infty} \frac{\left(\frac{3}{2}\right)_n}{(n+1)!} U_n(x) t^n, \quad R = \sqrt{1-2xt+t^2}. \tag{9.8.57}$$

$${}_0F_1\left(-; \frac{(x-1)t}{2}\right) {}_0F_1\left(-; \frac{(x+1)t}{2}\right) = \sum_{n=0}^{\infty} \frac{U_n(x)}{\left(\frac{3}{2}\right)_n (n+1)!} t^n. \tag{9.8.58}$$

$$e^{xt} {}_0F_1\left(-; \frac{(x^2-1)t^2}{4}\right) = \sum_{n=0}^{\infty} \frac{U_n(x)}{(n+1)!} t^n. \tag{9.8.59}$$

$$\begin{aligned}
& {}_2F_1\left(\gamma, 2-\gamma; \frac{1-R-t}{2}; \frac{1-R+t}{2}\right) {}_2F_1\left(\gamma, 2-\gamma; \frac{1-R+t}{2}; \frac{1-R-t}{2}\right) \\
&= \sum_{n=0}^{\infty} \frac{(\gamma)_n (2-\gamma)_n}{\left(\frac{3}{2}\right)_n (n+1)!} U_n(x) t^n, \quad R = \sqrt{1-2xt+t^2}, \quad \gamma \text{ arbitrary.}
\end{aligned} \tag{9.8.60}$$

$$\begin{aligned}
& (1-xt)^{-\gamma} {}_2F_1\left(\frac{1}{2}\gamma, \frac{1}{2}\gamma + \frac{1}{2}; \frac{(x^2-1)t^2}{(1-xt)^2}; \frac{3}{2}\right) \\
&= \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(n+1)!} U_n(x) t^n, \quad \gamma \text{ arbitrary.}
\end{aligned} \tag{9.8.61}$$

Remarks

The Chebyshev polynomials can also be written as:

$$T_n(x) = \cos(n\theta), \quad x = \cos \theta$$

and

$$U_n(x) = \frac{\sin(n+1)\theta}{\sin \theta}, \quad x = \cos \theta.$$

Further we have

$$U_n(x) = C_n^{(1)}(x)$$

where $C_n^{(\lambda)}(x)$ denotes the Gegenbauer (or ultraspherical) polynomial given by (9.8.19) in the preceding subsection.

References

[?], [?].

9.8.3 Legendre / Spherical**Hypergeometric representation**

The Legendre (or spherical) polynomials are Jacobi polynomials with $\alpha = \beta = 0$:

$$P_n(x) = P_n^{(0,0)}(x) = {}_2F_1 \left(\begin{matrix} -n, n+1 \\ 1 \end{matrix}; \frac{1-x}{2} \right). \quad (9.8.62)$$

Orthogonality relation

$$\int_{-1}^1 P_m(x) P_n(x) dx = \frac{2}{2n+1} \delta_{mn}. \quad (9.8.63)$$

Recurrence relation

$$(2n+1)xP_n(x) = (n+1)P_{n+1}(x) + nP_{n-1}(x). \quad (9.8.64)$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + \frac{n^2}{(2n-1)(2n+1)} p_{n-1}(x), \quad (9.8.65)$$

where

$$P_n(x) = \binom{2n}{n} \frac{1}{2^n} p_n(x).$$

Differential equation

$$(1-x^2)y''(x) - 2xy'(x) + n(n+1)y(x) = 0, \quad y(x) = P_n(x). \quad (9.8.66)$$

Rodrigues-type formula

$$P_n(x) = \frac{(-1)^n}{2^n n!} \left(\frac{d}{dx} \right)^n [(1-x^2)^n]. \quad (9.8.67)$$

Generating functions

$$\frac{1}{\sqrt{1-2xt+t^2}} = \sum_{n=0}^{\infty} P_n(x) t^n. \quad (9.8.68)$$

$${}_0F_1 \left(\begin{matrix} - \\ 1 \end{matrix}; \frac{(x-1)t}{2} \right) {}_0F_1 \left(\begin{matrix} - \\ 1 \end{matrix}; \frac{(x+1)t}{2} \right) = \sum_{n=0}^{\infty} \frac{P_n(x)}{(n!)^2} t^n. \quad (9.8.69)$$

$$e^{xt} {}_0F_1 \left(\begin{matrix} - \\ 1 \end{matrix}; \frac{(x^2-1)t^2}{4} \right) = \sum_{n=0}^{\infty} \frac{P_n(x)}{n!} t^n. \quad (9.8.70)$$

$$\begin{aligned} & {}_2F_1 \left(\begin{matrix} \gamma, 1-\gamma \\ 1 \end{matrix}; \frac{1-R-t}{2} \right) {}_2F_1 \left(\begin{matrix} \gamma, 1-\gamma \\ 1 \end{matrix}; \frac{1-R+t}{2} \right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n (1-\gamma)_n}{(n!)^2} P_n(x) t^n, \quad R = \sqrt{1-2xt+t^2}, \quad \gamma \text{ arbitrary.} \end{aligned} \quad (9.8.71)$$

$$\begin{aligned} & (1-xt)^{-\gamma} {}_2F_1\left(\begin{matrix} \frac{1}{2}\gamma, \frac{1}{2}\gamma + \frac{1}{2} \\ 1 \end{matrix}; \frac{(x^2-1)t^2}{(1-xt)^2}\right) \\ &= \sum_{n=0}^{\infty} \frac{(\gamma)_n}{n!} P_n(x) t^n, \quad \gamma \text{ arbitrary.} \end{aligned} \tag{9.8.72}$$

References

[illegible]

9.9 Pseudo Jacobi

Hypergeometric representation

$$\begin{aligned} P_n(x; \nu, N) &= \frac{(-2i)^n (-N + i\nu)_n}{(n - 2N - 1)_n} {}_2F_1 \left(\begin{matrix} -n, n - 2N - 1 \\ -N + i\nu \end{matrix}; \frac{1 - ix}{2} \right) \\ &= (x + i)^n {}_2F_1 \left(\begin{matrix} -n, N + 1 - n - i\nu \\ 2N + 2 - 2n \end{matrix}; \frac{2}{1 - ix} \right), \quad n = 0, 1, 2, \dots, N. \end{aligned} \quad (9.9.1)$$

Orthogonality relation

$$\begin{aligned} &\frac{1}{2\pi} \int_{-\infty}^{\infty} (1 + x^2)^{-N-1} e^{2\nu \arctan x} P_m(x; \nu, N) P_n(x; \nu, N) dx \\ &= \frac{\Gamma(2N + 1 - 2n) \Gamma(2N + 2 - 2n) 2^{2n-2N-1} n!}{\Gamma(2N + 2 - n) |\Gamma(N + 1 - n + i\nu)|^2} \delta_{mn}. \end{aligned} \quad (9.9.2)$$

Recurrence relation

$$\begin{aligned} xP_n(x; \nu, N) &= P_{n+1}(x; \nu, N) + \frac{(N + 1)\nu}{(n - N - 1)(n - N)} P_n(x; \nu, N) \\ &\quad - \frac{n(n - 2N - 2)}{(2n - 2N - 3)(n - N - 1)^2(2n - 2N - 1)} \\ &\quad \times (n - N - 1 - i\nu)(n - N - 1 + i\nu) P_{n-1}(x; \nu, N). \end{aligned} \quad (9.9.3)$$

Normalized recurrence relation

$$\begin{aligned} xp_n(x) &= p_{n+1}(x) + \frac{(N + 1)\nu}{(n - N - 1)(n - N)} p_n(x) \\ &\quad - \frac{n(n - 2N - 2)(n - N - 1 - i\nu)(n - N - 1 + i\nu)}{(2n - 2N - 3)(n - N - 1)^2(2n - 2N - 1)} p_{n-1}(x), \end{aligned} \quad (9.9.4)$$

where

$$P_n(x; \nu, N) = p_n(x).$$

Differential equation

$$(1 + x^2)y''(x) + 2(\nu - Nx)y'(x) - n(n - 2N - 1)y(x) = 0, \quad (9.9.5)$$

where

$$y(x) = P_n(x; \nu, N).$$

Forward shift operator

$$\frac{d}{dx} P_n(x; \nu, N) = nP_{n-1}(x; \nu, N - 1). \quad (9.9.6)$$

Backward shift operator

$$\begin{aligned}
& (1+x^2) \frac{d}{dx} P_n(x; \nu, N) + 2[\nu - (N+1)x] P_n(x; \nu, N) \\
& = (n - 2N - 2) P_{n+1}(x; \nu, N+1)
\end{aligned} \tag{9.9.7}$$

or equivalently

$$\begin{aligned}
& \frac{d}{dx} [(1+x^2)^{-N-1} e^{2\nu \arctan x} P_n(x; \nu, N)] \\
& = (n - 2N - 2) (1+x^2)^{-N-2} e^{2\nu \arctan x} P_{n+1}(x; \nu, N+1).
\end{aligned} \tag{9.9.8}$$

Rodrigues-type formula

$$P_n(x; \nu, N) = \frac{(1+x^2)^{N+1} e^{-2\nu \arctan x}}{(n-2N-1)_n} \left(\frac{d}{dx} \right)^n [(1+x^2)^{n-N-1} e^{2\nu \arctan x}]. \tag{9.9.9}$$

Generating function

$$\begin{aligned}
& \left[{}_0F_1 \left(\begin{matrix} - \\ -N+iv \end{matrix}; (x+i)t \right) {}_0F_1 \left(\begin{matrix} - \\ -N-iv \end{matrix}; (x-i)t \right) \right]_N \\
& = \sum_{n=0}^N \frac{(n-2N-1)_n}{(-N+iv)_n (-N-iv)_n n!} P_n(x; \nu, N) t^n.
\end{aligned} \tag{9.9.10}$$

Limit relation**Continuous Hahn \rightarrow Pseudo Jacobi**

The pseudo Jacobi polynomials follow from the continuous Hahn polynomials given by (9.4.1) by the substitutions $x \rightarrow xt$, $a = \frac{1}{2}(-N+iv-2t)$, $b = \frac{1}{2}(-N-iv+2t)$, $c = \frac{1}{2}(-N-iv-2t)$ and $d = \frac{1}{2}(-N+iv+2t)$, division by t^n and the limit $t \rightarrow \infty$:

$$\begin{aligned}
& \lim_{t \rightarrow \infty} \frac{p_n(xt; \frac{1}{2}(-N+iv-2t), \frac{1}{2}(-N-iv+2t), \frac{1}{2}(-N+iv-2t), \frac{1}{2}(-N-iv+2t))}{t^n} \\
& = \frac{(n-2N-1)_n}{n!} P_n(x; \nu, N).
\end{aligned}$$

Remarks

Since we have for $k < n$

$$\frac{(-N+iv)_n}{(-N+iv)_k} = (-N+iv+k)_{n-k},$$

the pseudo Jacobi polynomials given by (9.9.1) can also be seen as polynomials in the parameter ν .

The weight function for the pseudo Jacobi polynomials can be written as

$$(1+x^2)^{-N-1} e^{2\nu \arctan x} = (1+ix)^{-N-1-iv} (1-ix)^{-N-1+iv}.$$

The pseudo Jacobi polynomials are related to the Jacobi polynomials defined by (9.8.1) in the following way:

$$P_n(x; \nu, N) = \frac{(-2i)^n n!}{(n-2N-1)_n} P_n^{(-N-1+iv, -N-1-iv)}(ix).$$

If we set $x \rightarrow \nu x$ in the definition (9.9.1) of the pseudo Jacobi polynomials and take the limit $\nu \rightarrow \infty$ we obtain a special case of the Bessel polynomials given by (9.13.1) in the following way:

$$\lim_{\nu \rightarrow \infty} \frac{P_n(\nu x; \nu, N)}{\nu^n} = \frac{2^n}{(n-2N-1)_n} y_n(x; -2N-2).$$

References

[?], [?], [?].

9.10 Meixner**Hypergeometric representation**

$$M_n(x; \beta, c) = {}_2F_1 \left(\begin{matrix} -n, -x \\ \beta \end{matrix}; 1 - \frac{1}{c} \right). \quad (9.10.1)$$

Orthogonality relation

$$\sum_{x=0}^{\infty} \frac{(\beta)_x}{x!} c^x M_m(x; \beta, c) M_n(x; \beta, c) = \frac{c^{-n} n!}{(\beta)_n (1-c)^\beta} \delta_{mn} \quad (9.10.2)$$

Recurrence relation

$$(c-1)xM_n(x; \beta, c) = c(n+\beta)M_{n+1}(x; \beta, c) - [n + (n+\beta)c]M_n(x; \beta, c) + nM_{n-1}(x; \beta, c). \quad (9.10.3)$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + \frac{n + (n+\beta)c}{1-c} p_n(x) + \frac{n(n+\beta-1)c}{(1-c)^2} p_{n-1}(x), \quad (9.10.4)$$

where

$$M_n(x; \beta, c) = \frac{1}{(\beta)_n} \left(\frac{c-1}{c} \right)^n p_n(x).$$

Difference equation

$$n(c-1)y(x) = c(x+\beta)y(x+1) - [x + (x+\beta)c]y(x) + xy(x-1), \quad (9.10.5)$$

where

$$y(x) = M_n(x; \beta, c).$$

Forward shift operator

$$M_n(x+1; \beta, c) - M_n(x; \beta, c) = \frac{n}{\beta} \left(\frac{c-1}{c} \right) M_{n-1}(x; \beta+1, c) \quad (9.10.6)$$

or equivalently

$$\Delta M_n(x; \beta, c) = \frac{n}{\beta} \left(\frac{c-1}{c} \right) M_{n-1}(x; \beta+1, c). \quad (9.10.7)$$

Backward shift operator

$$c(\beta+x-1)M_n(x; \beta, c) - xM_n(x-1; \beta, c) = c(\beta-1)M_{n+1}(x; \beta-1, c) \quad (9.10.8)$$

or equivalently

$$\nabla \left[\frac{(\beta)_x c^x}{x!} M_n(x; \beta, c) \right] = \frac{(\beta-1)_x c^x}{x!} M_{n+1}(x; \beta-1, c). \quad (9.10.9)$$

Rodrigues-type formula

$$\frac{(\beta)_x c^x}{x!} M_n(x; \beta, c) = \nabla^n \left[\frac{(\beta + n)_x c^x}{x!} \right]. \quad (9.10.10)$$

Generating functions

$$\left(1 - \frac{t}{c}\right)^x (1-t)^{-x-\beta} = \sum_{n=0}^{\infty} \frac{(\beta)_n}{n!} M_n(x; \beta, c) t^n. \quad (9.10.11)$$

$$e^t {}_1F_1 \left(\begin{matrix} -x \\ \beta \end{matrix}; \left(\frac{1-c}{c} \right) t \right) = \sum_{n=0}^{\infty} \frac{M_n(x; \beta, c)}{n!} t^n. \quad (9.10.12)$$

$$(1-t)^{-\gamma} {}_2F_1 \left(\begin{matrix} \gamma, -x \\ \beta \end{matrix}; \frac{(1-c)t}{c(1-t)} \right) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{n!} M_n(x; \beta, c) t^n, \quad \gamma \text{ arbitrary.} \quad (9.10.13)$$

Limit relations**Hahn \rightarrow Meixner**

If we take $\alpha = b - 1$, $\beta = N(1 - c)c^{-1}$ in the definition (9.5.1) of the Hahn polynomials and let $N \rightarrow \infty$ we find the Meixner polynomials:

$$\lim_{N \rightarrow \infty} Q_n(x; b - 1, N(1 - c)c^{-1}, N) = M_n(x; b, c).$$

Dual Hahn \rightarrow Meixner

To obtain the Meixner polynomials from the dual Hahn polynomials we have to take $\gamma = \beta - 1$ and $\delta = N(1 - c)c^{-1}$ in the definition (9.6.1) of the dual Hahn polynomials and let $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} R_n(\lambda(x); \beta - 1, N(1 - c)c^{-1}, N) = M_n(x; \beta, c).$$

Meixner \rightarrow Laguerre

The Laguerre polynomials given by (9.12.1) are obtained from the Meixner polynomials if we take $\beta = \alpha + 1$ and $x \rightarrow (1 - c)^{-1}x$ and let $c \rightarrow 1$:

$$\lim_{c \rightarrow 1} M_n((1 - c)^{-1}x; \alpha + 1, c) = \frac{L_n^{(\alpha)}(x)}{L_n^{(\alpha)}(0)}. \quad (9.10.14)$$

Meixner \rightarrow Charlier

The Charlier polynomials given by (9.14.1) are obtained from the Meixner polynomials if we take $c = (a + \beta)^{-1}a$ and let $\beta \rightarrow \infty$:

$$\lim_{\beta \rightarrow \infty} M_n(x; \beta, (a + \beta)^{-1}a) = C_n(x; a). \quad (9.10.15)$$

Remarks

The Meixner polynomials are related to the Jacobi polynomials given by (9.8.1) in the following way:

$$\frac{(\beta)_n}{n!} M_n(x; \beta, c) = P_n^{\beta-1, -n-\beta-x}((2 - c)c^{-1}).$$

The Meixner polynomials are also related to the Krawtchouk polynomials given by (9.11.1) in the following way:

$$K_n(x; p, N) = M_n(x; -N, (p - 1)^{-1}p).$$

References

[illegible]

9.11 Krawtchouk

Hypergeometric representation

$$K_n(x; p, N) = {}_2F_1\left(\begin{matrix} -n, -x \\ -N \end{matrix}; \frac{1}{p}\right), \quad n = 0, 1, 2, \dots, N. \quad (9.11.1)$$

Orthogonality relation

$$\begin{aligned} & \sum_{x=0}^N \binom{N}{x} p^x (1-p)^{N-x} K_m(x; p, N) K_n(x; p, N) \\ &= \frac{(-1)^n n!}{(-N)_n} \left(\frac{1-p}{p} \right)^n \delta_{mn}, \quad 0 < p < 1. \end{aligned} \quad (9.11.2)$$

Recurrence relation

$$\begin{aligned}
 -xK_n(x; p, N) &= p(N-n)K_{n+1}(x; p, N) \\
 &\quad - [p(N-n) + n(1-p)]K_n(x; p, N) \\
 &\quad + n(1-p)K_{n-1}(x; p, N). \tag{9.11.3}
 \end{aligned}$$

Normalized recurrence relation

$$\begin{aligned} xp_n(x) &= p_{n+1}(x) + [p(N-n) + n(1-p)]p_n(x) \\ &\quad + np(1-p)(N+1-n)p_{n-1}(x), \end{aligned} \quad (9.11.4)$$

where

$$K_n(x; p, N) = \frac{1}{(-N)_n p^n} p_n(x).$$

Difference equation

$$\begin{aligned}
 -ny(x) &= p(N-x)y(x+1) \\
 &\quad - [p(N-x) + x(1-p)]y(x) + x(1-p)y(x-1), \quad (9.11.5)
 \end{aligned}$$

where

$$y(x) = K_n(x; p, N).$$

Forward shift operator

$$K_n(x+1; p, N) - K_n(x; p, N) = -\frac{n}{Np} K_{n-1}(x; p, N-1) \quad (9.11.6)$$

or equivalently

$$\Delta K_n(x; p, N) = -\frac{n}{Np} K_{n-1}(x; p, N-1). \quad (9.11.7)$$

Backward shift operator

$$\begin{aligned}
& (N+1-x)K_n(x; p, N) - x \left(\frac{1-p}{p} \right) K_n(x-1; p, N) \\
& = (N+1)K_{n+1}(x; p, N+1)
\end{aligned} \tag{9.11.8}$$

or equivalently

$$\nabla \left[\binom{N}{x} \left(\frac{p}{1-p} \right)^x K_n(x; p, N) \right] = \binom{N+1}{x} \left(\frac{p}{1-p} \right)^x K_{n+1}(x; p, N+1). \tag{9.11.9}$$

Rodrigues-type formula

$$\binom{N}{x} \left(\frac{p}{1-p} \right)^x K_n(x; p, N) = \nabla^n \left[\binom{N-n}{x} \left(\frac{p}{1-p} \right)^x \right]. \tag{9.11.10}$$

Generating functions

For $x = 0, 1, 2, \dots, N$ we have

$$\left(1 - \frac{(1-p)}{p} t \right)^x (1+t)^{N-x} = \sum_{n=0}^N \binom{N}{n} K_n(x; p, N) t^n. \tag{9.11.11}$$

$$\left[e^t {}_1F_1 \left(\begin{matrix} -x \\ -N \end{matrix}; -\frac{t}{p} \right) \right]_N = \sum_{n=0}^N \frac{K_n(x; p, N)}{n!} t^n. \tag{9.11.12}$$

$$\begin{aligned}
& \left[(1-t)^{-\gamma} {}_2F_1 \left(\begin{matrix} \gamma, -x \\ -N \end{matrix}; \frac{t}{p(t-1)} \right) \right]_N \\
& = \sum_{n=0}^N \frac{(\gamma)_n}{n!} K_n(x; p, N) t^n, \quad \gamma \text{ arbitrary.}
\end{aligned} \tag{9.11.13}$$

Limit relations**Hahn \rightarrow Krawtchouk**

If we take $\alpha = pt$ and $\beta = (1-p)t$ in the definition (9.5.1) of the Hahn polynomials and let $t \rightarrow \infty$ we obtain the Krawtchouk polynomials:

$$\lim_{t \rightarrow \infty} Q_n(x; pt, (1-p)t, N) = K_n(x; p, N).$$

Dual Hahn \rightarrow Krawtchouk

The Krawtchouk polynomials follow from the dual Hahn polynomials given by (9.6.1) if we set $\gamma = pt$, $\delta = (1-p)t$ and let $t \rightarrow \infty$:

$$\lim_{t \rightarrow \infty} R_n(\lambda(x); pt, (1-p)t, N) = K_n(x; p, N).$$

Krawtchouk \rightarrow Charlier

The Charlier polynomials given by (9.14.1) can be found from the Krawtchouk polynomials by taking $p = N^{-1}a$ and letting $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} K_n(x; N^{-1}a, N) = C_n(x; a). \tag{9.11.14}$$

Krawtchouk \rightarrow Hermite

The Hermite polynomials given by (9.15.1) follow from the Krawtchouk polynomials by setting $x \rightarrow pN + x\sqrt{2p(1-p)N}$ and then letting $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} \sqrt{\binom{N}{n}} K_n(pN + x\sqrt{2p(1-p)N}; p, N) = \frac{(-1)^n H_n(x)}{\sqrt{2^n n! \left(\frac{p}{1-p}\right)^n}}. \quad (9.11.15)$$

Remarks

The Krawtchouk polynomials are self-dual, which means that

$$K_n(x; p, N) = K_x(n; p, N), \quad n, x \in \{0, 1, 2, \dots, N\}.$$

By using this relation we easily obtain the so-called dual orthogonality relation from the orthogonality relation (9.11.2):

$$\sum_{n=0}^N \binom{N}{n} p^n (1-p)^{N-n} K_n(x; p, N) K_n(y; p, N) = \frac{\left(\frac{1-p}{p}\right)^x}{\binom{N}{x}} \delta_{xy},$$

where $0 < p < 1$ and $x, y \in \{0, 1, 2, \dots, N\}$.

The Krawtchouk polynomials are related to the Meixner polynomials given by (9.10.1) in the following way:

$$K_n(x; p, N) = M_n(x; -N, (p-1)^{-1}p).$$

References

[illegible]

9.12 Laguerre

Hypergeometric representation

$$L_n^{(\alpha)}(x) = \frac{(\alpha+1)_n}{n!} {}_1F_1\left(\begin{matrix} -n \\ \alpha+1 \end{matrix}; x\right). \quad (9.12.1)$$

Orthogonality relation

$$\int_0^\infty e^{-x} x^\alpha L_m^{(\alpha)}(x) L_n^{(\alpha)}(x) dx = \frac{\Gamma(n+\alpha+1)}{n!} \delta_{mn}, \quad \alpha > -1. \quad (9.12.2)$$

Recurrence relation

$$(n+1)L_{n+1}^{(\alpha)}(x) - (2n+\alpha+1-x)L_n^{(\alpha)}(x) + (n+\alpha)L_{n-1}^{(\alpha)}(x) = 0. \quad (9.12.3)$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + (2n + \alpha + 1)p_n(x) + n(n + \alpha)p_{n-1}(x), \quad (9.12.4)$$

where

$$L_n^{(\alpha)}(x) = \frac{(-1)^n}{n!} p_n(x).$$

Differential equation

$$xy''(x) + (\alpha + 1 - x)y'(x) + ny(x) = 0, \quad y(x) = L_n^{(\alpha)}(x). \quad (9.12.5)$$

Forward shift operator

$$\frac{d}{dx}L_n^{(\alpha)}(x) = -L_{n-1}^{(\alpha+1)}(x). \quad (9.12.6)$$

Backward shift operator

$$x \frac{d}{dx}L_n^{(\alpha)}(x) + (\alpha - x)L_n^{(\alpha)}(x) = (n+1)L_{n+1}^{(\alpha-1)}(x) \quad (9.12.7)$$

or equivalently

$$\frac{d}{dx} \left[e^{-x} x^\alpha L_n^{(\alpha)}(x) \right] = (n+1) e^{-x} x^{\alpha-1} L_{n+1}^{(\alpha-1)}(x). \quad (9.12.8)$$

Rodrigues-type formula

$$e^{-x} x^\alpha L_n^{(\alpha)}(x) = \frac{1}{n!} \left(\frac{d}{dx} \right)^n [e^{-x} x^{n+\alpha}]. \quad (9.12.9)$$

Generating functions

$$(1-t)^{-\alpha-1} \exp\left(\frac{xt}{t-1}\right) = \sum_{n=0}^{\infty} L_n^{(\alpha)}(x) t^n. \quad (9.12.10)$$

$$e^t {}_0F_1\left(\begin{matrix} - \\ \alpha+1 \end{matrix}; -xt\right) = \sum_{n=0}^{\infty} \frac{L_n^{(\alpha)}(x)}{(\alpha+1)_n} t^n. \quad (9.12.11)$$

$$(1-t)^{-\gamma} {}_1F_1\left(\begin{matrix} \gamma \\ \alpha+1 \end{matrix}; \frac{xt}{t-1}\right) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(\alpha+1)_n} L_n^{(\alpha)}(x) t^n, \quad \gamma \text{ arbitrary.} \quad (9.12.12)$$

Limit relations
Meixner-Pollaczek \rightarrow Laguerre

The Laguerre polynomials can be obtained from the Meixner-Pollaczek polynomials given by (9.7.1) by the substitution $\lambda = \frac{1}{2}(\alpha+1)$, $x \rightarrow -\frac{1}{2}\phi^{-1}x$ and the limit $\phi \rightarrow 0$:

$$\lim_{\phi \rightarrow 0} P_n^{(\frac{1}{2}\alpha+\frac{1}{2})}(-\frac{1}{2}\phi^{-1}x; \phi) = L_n^{(\alpha)}(x).$$

Jacobi \rightarrow Laguerre

The Laguerre polynomials are obtained from the Jacobi polynomials given by (9.8.1) if we set $x \rightarrow 1 - 2\beta^{-1}x$ and then take the limit $\beta \rightarrow \infty$:

$$\lim_{\beta \rightarrow \infty} P_n^{(\alpha, \beta)}(1 - 2\beta^{-1}x) = L_n^{(\alpha)}(x).$$

Meixner \rightarrow Laguerre

If we take $\beta = \alpha + 1$ and $x \rightarrow (1-c)^{-1}x$ in the definition (9.10.1) of the Meixner polynomials and let $c \rightarrow 1$ we obtain the Laguerre polynomials:

$$\lim_{c \rightarrow 1} M_n((1-c)^{-1}x; \alpha+1, c) = \frac{L_n^{(\alpha)}(x)}{L_n^{(\alpha)}(0)}.$$

$$\lim_{\alpha \rightarrow \infty} \left(\frac{2}{\alpha} \right)^{\frac{1}{2}n} L_n^{(\alpha)}((2\alpha)^{\frac{1}{2}}x + \alpha) = \frac{(-1)^n}{n!} H_n(x). \quad (9.12.13)$$
$$L_n^{(\alpha)}(x) = \frac{1}{n!} \sum_{k=0}^n \frac{(-n)_k}{k!} (\alpha + k + 1)_{n-k} x^k.$$
$$L_n^{(\alpha)}(x) = \frac{(-x)^n}{n!} y_n(2x^{-1}; -2n - \alpha - 1).$$
$$\frac{(-a)^n}{n!} C_n(x; a) = L_n^{(x-n)}(a).$$
$$H_{2n}(x) = (-1)^n n! 2^{2n} L_n^{(-\frac{1}{2})}(x^2)$$
$$H_{2n+1}(x) = (-1)^n n! 2^{2n+1} x L_n^{(\frac{1}{2})}(x^2).$$
[illegible]
$$\begin{aligned} y_n(x; a) &= {}_2F_0 \left(\begin{matrix} -n, n+a+1 \\ - \end{matrix}; -\frac{x}{2} \right) \\ &= (n+a+1)_n \left(\frac{x}{2} \right)^n {}_1F_1 \left(\begin{matrix} -n \\ -2n-a \end{matrix}; \frac{2}{x} \right), \quad n=0, 1, 2, \dots, N. \end{aligned} \quad (9.13.1)$$

Orthogonality relation

$$\begin{aligned}
& \int_0^\infty x^a e^{-\frac{2}{x}} y_m(x; a) y_n(x; a) dx \\
&= -\frac{2^{a+1}}{2n+a+1} \Gamma(-n-a) n! \delta_{mn}, \quad a < -2N-1.
\end{aligned} \tag{9.13.2}$$

Recurrence relation

$$\begin{aligned}
& 2(n+a+1)(2n+a)y_{n+1}(x; a) \\
&= (2n+a+1)[2a+(2n+a)(2n+a+2)x]y_n(x; a) \\
&\quad + 2n(2n+a+2)y_{n-1}(x; a).
\end{aligned} \tag{9.13.3}$$

Normalized recurrence relation

$$\begin{aligned}
xp_n(x) &= p_{n+1}(x) - \frac{2a}{(2n+a)(2n+a+2)} p_n(x) \\
&\quad - \frac{4n(n+a)}{(2n+a-1)(2n+a)^2(2n+a+1)} p_{n-1}(x),
\end{aligned} \tag{9.13.4}$$

where

$$y_n(x; a) = \frac{(n+a+1)_n}{2^n} p_n(x).$$

Differential equation

$$x^2 y''(x) + [(a+2)x+2] y'(x) - n(n+a+1)y(x) = 0, \quad y(x) = y_n(x; a). \tag{9.13.5}$$

Forward shift operator

$$\frac{d}{dx} y_n(x; a) = \frac{n(n+a+1)}{2} y_{n-1}(x; a+2). \tag{9.13.6}$$

Backward shift operator

$$x^2 \frac{d}{dx} y_n(x; a) + (ax+2)y_n(x; a) = 2y_{n+1}(x; a-2) \tag{9.13.7}$$

or equivalently

$$\frac{d}{dx} \left[x^a e^{-\frac{2}{x}} y_n(x; a) \right] = 2x^{a-2} e^{-\frac{2}{x}} y_{n+1}(x; a-2). \tag{9.13.8}$$

Rodrigues-type formula

$$y_n(x; a) = 2^{-n} x^{-a} e^{\frac{2}{x}} D^n \left(x^{2n+a} e^{-\frac{2}{x}} \right). \tag{9.13.9}$$

Generating function

$$(1-2xt)^{-\frac{1}{2}} \left(\frac{2}{1+\sqrt{1-2xt}} \right)^a \exp \left(\frac{2t}{1+\sqrt{1-2xt}} \right) = \sum_{n=0}^{\infty} y_n(x; a) \frac{t^n}{n!}. \tag{9.13.10}$$

Limit relation**Jacobi \rightarrow Bessel**

If we take $\beta = a - \alpha$ in the definition (9.8.1) of the Jacobi polynomials and let $\alpha \rightarrow -\infty$ we find the Bessel polynomials:

$$\lim_{\alpha \rightarrow -\infty} \frac{P_n^{(\alpha, a-\alpha)}(1 + \alpha x)}{P_n^{(\alpha, a-\alpha)}(1)} = y_n(x; a).$$

Remarks

The following notations are also used for the Bessel polynomials:

$$y_n(x; a, b) = y_n(2b^{-1}x; a) \quad \text{and} \quad \theta_n(x; a, b) = x^n y_n(x^{-1}; a, b).$$

However, the Bessel polynomials essentially depend on only one parameter.

The Bessel polynomials are related to the Laguerre polynomials given by (9.12.1) in the following way:

$$L_n^{(\alpha)}(x) = \frac{(-x)^n}{n!} y_n(2x^{-1}; -2n - \alpha - 1).$$

The special case $a = -2N - 2$ of the Bessel polynomials can be obtained from the pseudo Jacobi polynomials by setting $x \rightarrow vx$ in the definition (9.9.1) of the pseudo Jacobi polynomials and taking the limit $v \rightarrow \infty$ in the following way:

$$\lim_{v \rightarrow \infty} \frac{P_n(vx; v, N)}{v^n} = \frac{2^n}{(n - 2N - 1)_n} y_n(x; -2N - 2).$$

References

[?], [?], [?], [?], [?], [?], [?], [?], [?], [?], [?].

9.14 Charlier**Hypergeometric representation**

$$C_n(x; a) = {}_2F_0 \left(\begin{matrix} -n, -x \\ - \end{matrix} ; -\frac{1}{a} \right). \quad (9.14.1)$$

Orthogonality relation

$$\sum_{x=0}^{\infty} \frac{a^x}{x!} C_m(x; a) C_n(x; a) = a^{-n} e^a n! \delta_{mn}, \quad a > 0. \quad (9.14.2)$$

Recurrence relation

$$-xC_n(x; a) = aC_{n+1}(x; a) - (n + a)C_n(x; a) + nC_{n-1}(x; a). \quad (9.14.3)$$

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + (n + a)p_n(x) + nap_{n-1}(x), \quad (9.14.4)$$

where

$$C_n(x; a) = \left(-\frac{1}{a} \right)^n p_n(x).$$

Difference equation

$$-ny(x) = ay(x+1) - (x+a)y(x) + xy(x-1), \quad y(x) = C_n(x; a). \quad (9.14.5)$$

Forward shift operator

$$C_n(x+1; a) - C_n(x; a) = -\frac{n}{a}C_{n-1}(x; a) \quad (9.14.6)$$

or equivalently

$$\Delta C_n(x; a) = -\frac{n}{a}C_{n-1}(x; a). \quad (9.14.7)$$

Backward shift operator

$$C_n(x; a) - \frac{x}{a}C_n(x-1; a) = C_{n+1}(x; a) \quad (9.14.8)$$

or equivalently

$$\nabla \left[\frac{a^x}{x!} C_n(x; a) \right] = \frac{a^x}{x!} C_{n+1}(x; a). \quad (9.14.9)$$

Rodrigues-type formula

$$\frac{a^x}{x!} C_n(x; a) = \nabla^n \left[\frac{a^x}{x!} \right]. \quad (9.14.10)$$

Generating function

$$e^t \left(1 - \frac{t}{a} \right)^x = \sum_{n=0}^{\infty} \frac{C_n(x; a)}{n!} t^n. \quad (9.14.11)$$

Limit relations
Meixner \rightarrow Charlier

If we take $c = (a + \beta)^{-1}a$ in the definition (9.10.1) of the Meixner polynomials and let $\beta \rightarrow \infty$ we find the Charlier polynomials:

$$\lim_{\beta \rightarrow \infty} M_n(x; \beta, (a + \beta)^{-1}a) = C_n(x; a).$$

Krawtchouk \rightarrow Charlier

The Charlier polynomials can be found from the Krawtchouk polynomials given by (9.11.1) by taking $p = N^{-1}a$ and letting $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} K_n(x; N^{-1}a, N) = C_n(x; a).$$

Charlier \rightarrow Hermite

The Hermite polynomials given by (9.15.1) are obtained from the Charlier polynomials if we set $x \rightarrow (2a)^{1/2}x + a$ and let $a \rightarrow \infty$. In fact we have

$$\lim_{a \rightarrow \infty} (2a)^{\frac{1}{2}n} C_n((2a)^{\frac{1}{2}}x + a; a) = (-1)^n H_n(x). \quad (9.14.12)$$

$$\frac{d}{dx} \left[e^{-x^2} H_n(x) \right] = -e^{-x^2} H_{n+1}(x). \quad (9.15.8)$$

Rodrigues-type formula

$$e^{-x^2} H_n(x) = (-1)^n \left(\frac{d}{dx} \right)^n \left[e^{-x^2} \right]. \quad (9.15.9)$$

Generating functions

$$\exp(2xt - t^2) = \sum_{n=0}^{\infty} \frac{H_n(x)}{n!} t^n. \quad (9.15.10)$$

$$\begin{cases} e^t \cos(2x\sqrt{t}) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)!} H_{2n}(x) t^n \\ \frac{e^t}{\sqrt{t}} \sin(2x\sqrt{t}) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)!} H_{2n+1}(x) t^n. \end{cases} \quad (9.15.11)$$

$$\begin{cases} e^{-t^2} \cosh(2xt) = \sum_{n=0}^{\infty} \frac{H_{2n}(x)}{(2n)!} t^{2n} \\ e^{-t^2} \sinh(2xt) = \sum_{n=0}^{\infty} \frac{H_{2n+1}(x)}{(2n+1)!} t^{2n+1}. \end{cases} \quad (9.15.12)$$

$$\begin{cases} (1+t^2)^{-\gamma} {}_1F_1 \left(\frac{\gamma}{2}; \frac{x^2 t^2}{1+t^2} \right) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{(2n)!} H_{2n}(x) t^{2n} \\ \frac{xt}{\sqrt{1+t^2}} {}_1F_1 \left(\gamma + \frac{1}{2}; \frac{x^2 t^2}{1+t^2} \right) = \sum_{n=0}^{\infty} \frac{(\gamma + \frac{1}{2})_n}{(2n+1)!} H_{2n+1}(x) t^{2n+1} \end{cases} \quad (9.15.13)$$

with γ arbitrary.

$$\frac{1+2xt+4t^2}{(1+4t^2)^{\frac{3}{2}}} \exp \left(\frac{4x^2 t^2}{1+4t^2} \right) = \sum_{n=0}^{\infty} \frac{H_n(x)}{[n/2]!} t^n, \quad (9.15.14)$$

where $[n/2]$ denotes the largest integer smaller than or equal to $n/2$.

Limit relations**Meixner-Pollaczek \rightarrow Hermite**

If we take $x \rightarrow (\sin \phi)^{-1}(x\sqrt{\lambda} - \lambda \cos \phi)$ in the definition (9.7.1) of the Meixner-Pollaczek polynomials and then let $\lambda \rightarrow \infty$ we obtain the Hermite polynomials:

$$\lim_{\lambda \rightarrow \infty} \lambda^{-\frac{1}{2}n} P_n^{(\lambda)}((\sin \phi)^{-1}(x\sqrt{\lambda} - \lambda \cos \phi); \phi) = \frac{H_n(x)}{n!}.$$

Jacobi \rightarrow Hermite

The Hermite polynomials follow from the Jacobi polynomials given by (9.8.1) by taking $\beta = \alpha$ and letting $\alpha \rightarrow \infty$ in the following way:

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\frac{1}{2}n} P_n^{(\alpha, \alpha)}(\alpha^{-\frac{1}{2}}x) = \frac{H_n(x)}{2^n n!}.$$

Gegenbauer / Ultraspherical \rightarrow Hermite

The Hermite polynomials follow from the Gegenbauer (or ultraspherical) polynomials given by (9.8.19) by taking $\lambda = \alpha + \frac{1}{2}$ and letting $\alpha \rightarrow \infty$ in the following way:

$$\lim_{\alpha \rightarrow \infty} \alpha^{-\frac{1}{2}n} C_n^{(\alpha + \frac{1}{2})}(\alpha^{-\frac{1}{2}}x) = \frac{H_n(x)}{n!}.$$

Krawtchouk \rightarrow Hermite

The Hermite polynomials follow from the Krawtchouk polynomials given by (9.11.1) by setting $x \rightarrow pN + x\sqrt{2p(1-p)N}$ and then letting $N \rightarrow \infty$:

$$\lim_{N \rightarrow \infty} \sqrt{\binom{N}{n}} K_n(pN + x\sqrt{2p(1-p)N}; p, N) = \frac{(-1)^n H_n(x)}{\sqrt{2^n n! \left(\frac{p}{1-p}\right)^n}}.$$

Laguerre \rightarrow Hermite

The Hermite polynomials can be obtained from the Laguerre polynomials given by (9.12.1) by taking the limit $\alpha \rightarrow \infty$ in the following way:

$$\lim_{\alpha \rightarrow \infty} \left(\frac{2}{\alpha} \right)^{\frac{1}{2}n} L_n^{(\alpha)}((2\alpha)^{\frac{1}{2}}x + \alpha) = \frac{(-1)^n}{n!} H_n(x).$$

Charlier \rightarrow Hermite

If we set $x \rightarrow (2a)^{1/2}x + a$ in the definition (9.14.1) of the Charlier polynomials and let $a \rightarrow \infty$ we find the Hermite polynomials. In fact we have

$$\lim_{q \rightarrow \infty} (2a)^{\frac{1}{2}n} C_n((2a)^{\frac{1}{2}}x + a; a) = (-1)^n H_n(x).$$

Remarks

The Hermite polynomials can also be written as:

$$\frac{H_n(x)}{n!} = \sum_{k=0}^{\lfloor n/2 \rfloor} \frac{(-1)^k (2x)^{n-2k}}{k! (n-2k)!},$$

where $\lfloor n/2 \rfloor$ denotes the largest integer smaller than or equal to $n/2$.

The Hermite polynomials and the Laguerre polynomials given by (9.12.1) are also connected by the following quadratic transformations:

$$H_{2n}(x) = (-1)^n n! 2^{2n} L_n^{(-\frac{1}{2})}(x^2)$$

and

$$H_{2n+1}(x) = (-1)^n n! 2^{2n+1} x L_n^{(\frac{1}{2})}(x^2).$$

References

[illegible]