

Regression S	tatistics	Wei	ght = 6	$B_0 + B_1 N$	$1ale + \beta_2 H$	leight
Multiple R	0.78801776		,	0 . 71		
R Square	0.62097199					
Adjusted R Squar	0.62049342					
Standard Error	9.86271666					
Observations	1587					
ANOVA						
	df	SS	MS	F	Significance F	
Regression	2	252434.6717	126217.3	1297.555	0	
Residual	1584	154080.7171	97.27318			
Total	1586	406515.3888				-
	Coefficients :	Standard Error	t Stat	P-value	Lower 95%	Upper 95%
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793
Height(cm)	0.96697779	0.026071479	37.08949	2.8E-217	0.915839553	1.018116



Regression S	tatistics	Wei	ght = 6	$B_0 + B_1 N$	$Male + \beta_2 H$	leight
Multiple R	0.78801776		,	0 ,1	, -	· ·
R Square	0.62097199					
Adjusted R Squar	0.62049342					
Standard Error	9.86271666					
Observations	1587					
ANOVA						L.
	df	SS	MS	F	Significance F	
Regression	2	252434.6717	126217.3	1297.555	0	
Residual	1584	154080.7171	97.27318			
Total	1586	406515.3888				
	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793
Height(cm)	0.96697779	0.026071479	37.08949	2.8E-217	0.915839553	1.018116



Regression S	tatistics	Wei	ght = 6	$B_0 + B_1 N$	$Male + \beta_2 H$	leight
Multiple R	0.78801776		,	0 , 1	, -	
R Square	0.62097199					
Adjusted R Squar	0.62049342					
Standard Error	9.86271666					
Observations	1587					
ANOVA						
	df	SS	MS	F	Significance F	
Regression	2	252434.6717	126217.3	1297.555	0	
Residual	1584	154080.7171	97.27318			
Total	1586	406515.3888				
	Coefficients :	Standard Error	t Stat	P-value	Lower 95%	Upper 95%
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793
Height(cm)	0.96697779	0.026071479	37.08949	2.8E-217	0.915839553	1.018116



Regression S	tatistics	Wei	$Weight = \beta_0 + \beta_1 Male + \beta_2 Height$							
Multiple R	0.78801776		,	0 ,1	, -	0				
R Square	0.62097199	Effect	t of ' <i>Hei</i> ç	<i>ght</i> on '	<i>Weight</i> is s	ame acros				
Adjusted R Squar	0.62049342									
Standard Error	9.86271666									
Observations	1587									
ANOVA										
	df	SS	MS	F	Significance F					
Regression	2	252434.6717	126217.3	1297.555	0					
Residual	1584	154080.7171	97.27318							
Total	1586	406515.3888								
	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%				
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282				
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793				
Height(cm)	0.96697779	0.026071479	37.08949	2.8E-217	0.915839553	1.018116				



Regression S	tatistics	Wei	ght = f	$B_0 + \beta_1 N$	$Aale + \beta_2 H$	leight	
Multiple R	0.78801776					Ü	
R Square	0.62097199	Effect	t of ' <i>Hei</i>	<i>ght</i> on '	<i>Weight</i> is s	ame acros	s Gender
Adjusted R Squar	0.62049342	Dumi	my varia	ble ' <i>Mal</i> e	e' allows for	r some ove	rall differ
Standard Error	9.86271666		•				
Observations	1587						
ANOVA							
	df	SS	MS	F	Significance F		
Regression	2	252434.6717	126217.3	1297.555	0		
Residual	1584	154080.7171	97.27318				
Total	1586	406515.3888					
	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282	
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793	
Height(cm)	0.96697779	0.026071479	37.08949	2 8F-217	0.915839553	1.018116	



Interaction Effects in a Regression Model.

Regression S	tatistics	Wei	ght = f	$B_0 + B_1 N$	$1ale + \beta_2 H$	leight	
Multiple R	0.78801776		,	0 71	, -	Ü	
R Square	0.62097199	Effect	t of ' <i>Hei</i>	ghť on '	<i>Weight</i> is s	ame acros	ss Gender.
Adjusted R Squar	0.62049342	Dumi	my varia	ble ' <i>Mal</i> e	e' allows for	r some ove	erall difference
Standard Error	9.86271666				allowed ba		
Observations	1587	- 140 C	ciidei di	noronoo	anowed ba	300 OH 77	oigiit .
ANOVA							
	df	SS	MS	F	Significance F		
Regression	2	252434.6717	126217.3	1297.555	0		
Residual	1584	154080.7171	97.27318				
Total	1586	406515.3888					
	Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	
Intercept	-101.96468	4.466768794	-22.8274	6E-100	-110.726083	-93.203282	
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793	
Height(cm)	0.96697779	0.026071479	37.08949	2.8E-217	0.915839553	1.018116	



Regression S	tatistics	Wei	ght = 6	$B_0 + B_1 N$	$1ale + \beta_2 H$	leight	
Multiple R	0.78801776		,	0 . 71			
R Square	0.62097199	Effec	t of ' <i>Hei</i> ç	<i>ght</i> on '	<i>Weight</i> ' is s	ame acros	s Gender.
Adjusted R Squar	0.62049342	Dumi	mv varia	ble ' <i>Mal</i> e	e' allows for	r some ove	rall differe
Standard Error	9.86271666				allowed ba		
Observations	1587	1 110 0	chack at	Herence	anowed ba	360 OH 776	agric.
		How de	o we allo	w impac	t of 'Height	dased on	'Gender'?
ANOVA							
	df	SS	MS	F	Significance F		
Regression	2	252434.6717	126217.3	1297.555	0		
Residual	1584	154080.7171	97.27318				
Total	1586	406515.3888					
	Coefficients :	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	
Intercept	-101.96468	4.466768794		6E-100	-110.726083	1	
Male	5.52761923	0.591069119	9.3519	2.83E-20	4.368259169	6.6869793	
Height(cm)	0.96697779	0.026071479	37 08949	2 8F-217	0.915839553	1.018116	



$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height$$



$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height$$



$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height$$



$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height + \beta_3 \underline{Male * Height}$$
Interaction Variable

$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height + \beta_3 Male * Height$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height + \beta_3 Male * Height$$

Value of Y variable when all X variables are zero.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height + \beta_3 Male * Height$$

Value of Y variable when all X variables are zero.

No 'managerial' interpretation.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 Height + \beta_3 Male * Height$$

Value of Y variable when all X variables are zero.

No 'managerial' interpretation.

We could mean-center the variable 'Height'.



$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

$$\uparrow$$
Mean Centered
Mean Centered

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

$$Male = 0$$
,

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

$$Male = 0, \quad \overline{Height} = 0,$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

$$Male = 0$$
, $\overline{Height} = 0$, $Male * \overline{Height} = 0$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

'Weight' of an Olympian when all X variables are zero.

$$Male = 0$$
, $\overline{Height} = 0$, $Male * \overline{Height} = 0$

□ Olympian is a *Female*.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

$$Male = 0$$
, $\overline{Height} = 0$, $Male * \overline{Height} = 0$

- □ Olympian is a *Female*.
- □ *Actual height* is the average height observed in data.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

'Weight' of an Olympian when all X variables are zero.

$$Male = 0$$
, $\overline{Height} = 0$, $Male * \overline{Height} = 0$

- □ Olympian is a *Female*.
- □ *Actual height* is the average height observed in data.

 β_0 is the weight of a Female Olympian whose height is at a level equal to the average height observed in the data.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The *additional weight* of a Male Olympian as compared to a Female Olympian when the height is equal to the average height observed in the data.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The *additional weight* of a Male Olympian as compared to a Female Olympian when the height is equal to the average height observed in the data.

 $\beta_0 + \beta_1 = Total \ weight$ of a Male Olympian whose height is equal to the average height observed in the data.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The impact of one centimeter increase in height on the weight, all other variables held constant.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The impact of one centimeter increase in height on the weight, all other variables held constant.

To keep the interaction variable 'at the same level', in our interpretation, we need to consider a Female Olympian.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The impact of one centimeter increase in height on the weight, all other variables held constant.

To keep the interaction variable 'at the same level', in our interpretation, we need to consider a Female Olympian.

 β_2 is the impact of one centimeter increase in height on the weight of Female Olympians.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The impact of one centimeter increase in height on the weight, all other variables held constant.

To keep the interaction variable 'at the same level', in our interpretation, we need to consider a Female Olympian.

 β_2 is the impact of one centimeter increase in height on the weight of Female Olympians.

What about the impact of height on the weight of Male Olympians?



$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The additional impact of Height on weight, for Male Olympians.

Interaction Effects in a Regression Model.

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

The additional impact of Height on weight, for Male Olympians.

The *total impact* of Height on weight for Male Olympians is $\beta_2 + \beta_3$

$$Weight = \beta_0 + \beta_1 Male + \beta_2 \overline{Height} + \beta_3 Male * \overline{Height}$$

- \square β_0 is the weight of a Female Olympian whose height is at a level equal to the average height observed in the data.
- \square β_1 is the *additional weight* of a Male Olympian vis-à-vis a Female Olympian when the height is equal to the average height observed in the data.
- \Box $\beta_0 + \beta_1 = Total \ weight$ of a Male Olympian whose height is equal to the average height observed in the data.
- \square β_2 is the impact of one centimeter increase in height on the weight of Female Olympians.
- $\ \square$ $\ \beta_3$ is the *additional impact* of Height on weight, for Male Olympians.
- \Box $\beta_2 + \beta_3$ is the *total impact* of Height on weight for Male Olympians.