

$$Sales = \beta_0 + \beta_1 Price + \beta_2 AdExp + \beta_3 PromExp$$

$$Sales = \beta_0 + \beta_1 Price + \beta_2 AdExp + \beta_3 PromExp$$

Could the true value of  $\beta_2$  be 500?

Sales = 
$$\beta_0$$
 +  $\beta_1 Price$  +  $\beta_2 AdExp$  +  $\beta_3 PromExp$ 

Could the true value of  $\beta_2$  be 500?

$$H_0$$
:  $\beta_2 = 500$   
 $H_A$ :  $\beta_2 \neq 500$ 

$$Sales = \beta_0 + \beta_1 Price + \beta_2 AdExp + \beta_3 PromExp$$

Could the true value of  $\beta_2$  be 500?

$$H_0$$
:  $\beta_2 = 500$   
 $H_{\Delta}$ :  $\beta_2 \neq 500$ 

#### **Conclusion:**

- > Do not reject the Null hypothesis
- > True value of  $\beta_2$  could be 500
- We cannot reject the belief held by salespeople



Another approach to Hypothesis Testing



Another approach to Hypothesis Testing the p-value approach

#### **Step 1: Formulate Hypothesis**

 $H_0$ :  $\beta_2 = 500$ 

 $H_A$ :  $\beta_2 \neq 500$ 

#### Step 2: Calculate the t-statistic

t-statistic = 
$$\frac{b_2 - \beta_2}{s_{b_2}} = 0.711$$

Another approach to Hypothesis Testing the p-value approach

#### Step 1: Formulate Hypothesis



$$H_0$$
:  $\beta_2 = 500$ 

$$H_A$$
:  $\beta_2 \neq 500$ 

#### Step 2: Calculate the t-statistic

t-statistic = 
$$\frac{b_2 - \beta_2}{s_{b_2}} = 0.711$$

Another approach to Hypothesis Testing the p-value approach

#### **Step 1: Formulate Hypothesis**

 $H_0$ :  $\beta_2 = 500$ 

 $H_A$ :  $\beta_2 \neq 500$ 

#### Step 2: Calculate the t-statistic

$$\checkmark \qquad \text{t-statistic} = \frac{b_2 - \beta_2}{s_{b_2}} = 0.711$$

#### Another approach to Hypothesis Testing the p-value approach

Step 1: Formulate Hypothesis

$$H_0$$
:  $\beta_2 = 500$ 


$$H_A$$
:  $\beta_2 \neq 500$ 

Step 2: Calculate the t-statistic


t-statistic = 
$$\frac{b_2 - \beta_2}{s_{b_2}} = 0.711$$

#### Step 3: Calculate the p-value


































































```
p-value = 2*T.DIST(-|t-statistic|, residual df, TRUE)
= 0.4853
```



Another approach to Hypothesis Testing the p-value approach

```
p-value = 2*T.DIST(-|t-statistic|, residual df, TRUE)
= 0.4853
```

#### **Conclusion:**

- > Do not reject the Null hypothesis
- We cannot reject the belief held by salespeople



Another approach to Hypothesis Testing the p-value approach

```
p-value = 2*T.DIST(-|t-statistic|, residual df, TRUE)
= 0.4853
```

#### **Conclusion:**

- > Do not reject the Null hypothesis
- We cannot reject the belief held by salespeople



- > The t-cutoff approach
- > The p-value approach



- → > The t-cutoff approach
  - > The p-value approach



- > The t-cutoff approach
- → > The p-value approach



- > The t-cutoff approach
- > The p-value approach
- → > The confidence interval approach