$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = 35.027 - 0.0579 Displacement$$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = 35.027 - 0.0579 Displacement$$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement$$
 $MPG = 35.027 - 0.0579 Displacement$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$
(p-value 1.07x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Displacement$$
 $MPG = 35.027 - 0.0579 Displacement$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$

 $MPG = 42.509 - 3.29 Cylinders$
(p-value 1.07x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Displacement$$
 $MPG = 35.027 - 0.0579 Displacement$
(p-value 4.99x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = 42.509 - 3.29 Cylinders$$
(p-value 1.07x10⁻⁰⁹)

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

X variable 'significant'.

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

X variable 'significant'.

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

X variable 'significant'.

$$MPG = \beta_0 + \beta_1 Cylinders$$

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

X variable 'significant'.

$$MPG = \beta_0 + \beta_1 Cylinders$$



$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- Both X variables 'insignificant'
- Reasonable R-square fits

$$MPG = \beta_0 + \beta_1 Displacement$$

X variable 'significant'.

$$MPG = \beta_0 + \beta_1 Cylinders$$

X variable 'significant'.



- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.



- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.



- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.



- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.

- Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- □ Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- > This model may predict reasonably.
- However, individual impact of *Displacement* and *Cylinders* cannot be discerned.

- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- □ Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- > This model may predict reasonably.
- However, individual impact of *Displacement* and *Cylinders* cannot be discerned.

- □ Occurs when two or more X variables are highly correlated.
- □ Variation in Y cannot be apportioned individually across X variables.
- □ Model fits may be reasonable and prediction may be ok.
- □ Interpretation of individual X variable impact will be suspect.

$$MPG = \beta_0 + \beta_1 Displacement + \beta_2 Cylinders$$

- > This model may predict reasonably.
- > However, individual impact of *Displacement* and *Cylinders* cannot be discerned.



Detecting Multicollinearity

- Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.



Detecting Multicollinearity

- Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.



Detecting Multicollinearity

- □ Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.

- □ Do you *need* to correct for collinearity?
 - May not be a problem if using regression only to predict.
 - Collinearity is problematic when interpreting coefficient impacts.
- □ To remove multicollinearity you may drop variables causing high correlation.



Detecting Multicollinearity

- □ Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.

- □ Do you need to correct for collinearity?
 - May not be a problem if using regression only to predict.
 - Collinearity is problematic when interpreting coefficient impacts.
- □ To remove multicollinearity you may drop variables causing high correlation.



Detecting Multicollinearity

- □ Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.

- □ Do you need to correct for collinearity?
 - May not be a problem if using regression only to predict.
 - Collinearity is problematic when interpreting coefficient impacts.
- □ To remove multicollinearity you may drop variables causing high correlation.



Detecting Multicollinearity

- Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.

- □ Do you need to correct for collinearity?
 - May not be a problem if using regression only to predict.
 - Collinearity is problematic when interpreting coefficient impacts.
- □ To remove multicollinearity you may drop variables causing high correlation.



Detecting Multicollinearity

- Some software produce collinearity statistics such as VIF (Variance Inflation Factor).
- □ We can look at correlations across X variables.

- □ Do you need to correct for collinearity?
 - May not be a problem if using regression only to predict.
 - Collinearity is problematic when interpreting coefficient impacts.
- □ To remove multicollinearity you may drop variables causing high correlation.