

Incorporating 'Categorical Variables' in a regression model.



Incorporating 'Categorical Variables' in a regression model.

...the Dummy Variable regression technique

A parcel delivery service operates in two different regions, **region "A"** and **region "B"**. Delivery trucks leave the central warehouse and travel to region A and deliver parcels in that region. Similarly delivery trucks also leave the central warehouse and travel to region B and deliver parcels in that region.



Incorporating 'Categorical Variables' in a regression model.

...the Dummy Variable regression technique

Region




Incorporating 'Categorical Variables' in a regression model.





Incorporating 'Categorical Variables' in a regression model.





Incorporating 'Categorical Variables' in a regression model.

...the Dummy Variable regression technique



Region One 'dummy variable' needed (REGA)



Incorporating 'Categorical Variables' in a regression model.



	Α	В
REGA	1	0



Incorporating 'Categorical Variables' in a regression model.



	Α	В
REGA	1	0
,	<b>†</b>	



Incorporating 'Categorical Variables' in a regression model.



	Α	В
REGA	1	0
		1

Incorporating 'Categorical Variables' in a regression model.

$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 Parcels + \beta_3 TruckAge$$

Incorporating 'Categorical Variables' in a regression model.

$$\label{eq:minutes} \begin{aligned} \textit{Minutes} &= \beta_0 + \beta_1 \textit{REGA} + \beta_2 \textit{Parcels} + \beta_3 \textit{TruckAge} \\ \\ \textit{Minutes} &= -33.13 + 106.84 \textit{REGA} + 10.02 \textit{Parcels} + 3.21 \textit{TruckAge} \end{aligned}$$



Incorporating 'Categorical Variables' in a regression model.

Example (deliveries2.xlsx)

The parcel delivery service obtains some additional data, this time for a third region it delivers to, region C. So now, the data consists of deliveries across three regions, region A, region B and region C.



Incorporating 'Categorical Variables' in a regression model.

Example (deliveries2.xlsx)

The parcel delivery service obtains some additional data, this time for a third region it delivers to, <u>region C</u>. So now, the data consists of deliveries across three regions, region A, region B and region C.



Incorporating 'Categorical Variables' in a regression model.

Example (deliveries2.xlsx)

The parcel delivery service obtains some additional data, this time for a third region it delivers to, region C. So now, the data consists of deliveries across three regions, **region A**, **region B** and **region C**.



Incorporating 'Categorical Variables' in a regression model.


#### **Example** (deliveries2.xlsx)

The parcel delivery service obtains some additional data, this time for a third region it delivers to, region C. So now, the data consists of deliveries across three regions, **region A**, **region B** and **region C**.
























































$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$






$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$




$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$




$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$






$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$




$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$




$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$






$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$






$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$





$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$



Incorporating 'Categorical Variables' in a regression model.



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Difference in overall delivery time it takes to make deliveries to Region A as compared to Region C, all other variables kept at the same level.



Incorporating 'Categorical Variables' in a regression model.



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Difference in overall delivery time it takes to make deliveries to Region A as compared to Region C, all other variables kept at the same level.




Incorporating 'Categorical Variables' in a regression model.



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Difference in overall delivery time it takes to make deliveries to Region A as compared to Region C, all other variables kept at the same level.

Incorporating 'Categorical Variables' in a regression model.




$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Is the difference in the fixed time to make deliveries across the two regions.



Incorporating 'Categorical Variables' in a regression model.




p-value > 
$$\alpha$$

$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Is the difference in the fixed time to make deliveries across the two regions.



Incorporating 'Categorical Variables' in a regression model.



$$Minutes = \beta_0 + \beta_1 REGA + \beta_2 REGB + \beta_3 Parcels + \beta_4 TruckAge$$

Difference in overall delivery time it takes to make deliveries to Region B as compared to Region C, all other variables kept at the same level.