

# Math 300 NTI Lesson 15

## Multiple Regression - Two Numerical

Professor Bradley Warner

June, 2022

### Contents

Objectives . . . . .	1
Reading . . . . .	1
Lesson . . . . .	1
Documenting software . . . . .	7

### Objectives

1. For two numerical explanatory variables in a linear regression model, conduct exploratory analysis and explain the relationship between the variables.
2. Fit a linear regression model to two numerical explanatory variables using the `lm()` function and interpret the output.

### Reading

Chapter 6.2

### Lesson

*Remember that you will be running this more like a lab than a lecture. You want them using R and answering questions. Have them open the notes rmd and work through it together.*

Work through the learning checks LC6.2 - 6.3.

- We use `select()` to change the names of the variables as well as to select them.
- Collinearity (or multicollinearity) is a phenomenon where one explanatory variable in a multiple regression model is highly correlated with another. This course doesn't discuss multicollinearity but it impacts the inference portion of the analysis cycle. Math 378 is a course that presents methods to handle multicollinearity.
- We preface our interpretation with the statement, "taking into account all the other explanatory variables in our model" in this section. This means we have to treat the other variables as at a constant value even though collinearity in practice may not allow this. It is only from an interpretation point of view that we use that statement.

- A phenomenon known as Simpson's Paradox, whereby overall trends that exist in aggregate either disappear or reverse when the data are broken down into groups. The next lesson discusses this in more depth.

## Setup

```
library(tidyverse)
library(moderndive)
library(skimr)
library(ISLR)
```

*Recreate the analysis done in the book.*

```
my_skim <- skim_with(numeric = sfl(hist = NULL))
```

```
credit_ch6 <- Credit %>% as_tibble() %>%
  select(ID, debt = Balance, credit_limit = Limit,
         income = Income, credit_rating = Rating, age = Age)
```

Let's look at 5 random rows of data.

```
set.seed(507)
credit_ch6 %>%
  sample_n(size = 5)
```

```
## # A tibble: 5 x 6
##       ID debt credit_limit income credit_rating age
##   <int> <int>      <int>  <dbl>      <int> <int>
## 1   218   955      5395   12.5        392    65
## 2   160    0      3000   53.3        235    53
## 3   112    0      2959   28.6        231    60
## 4    77   532      3293   30.6        251    68
## 5   265   651      5107   28.0        380    55
```

```
glimpse(credit_ch6)
```

```
## Rows: 400
## Columns: 6
## $ ID      <int> 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 1~
## $ debt    <int> 333, 903, 580, 964, 331, 1151, 203, 872, 279, 1350, 1407~
## $ credit_limit <int> 3606, 6645, 7075, 9504, 4897, 8047, 3388, 7114, 3300, 68~
## $ income  <dbl> 14.891, 106.025, 104.593, 148.924, 55.882, 80.180, 20.99~
## $ credit_rating <int> 283, 483, 514, 681, 357, 569, 259, 512, 266, 491, 589, 1~
## $ age     <int> 34, 82, 71, 36, 68, 77, 37, 87, 66, 41, 30, 64, 57, 49, ~
```

```
credit_ch6 %>%
  select(debt, credit_limit, income) %>%
  my_skim() %>%
  print()
```

```
## -- Data Summary -----
##                               Values
## Name                         Piped data
## Number of rows                400
## Number of columns             3
## -----
## Column type frequency:
##   numeric                      3
## -----
## Group variables               None
##
## -- Variable type: numeric -----
##   skim_variable n_missing complete_rate   mean    sd    p0    p25    p50    p75
## 1 debt          0           1  520.    460.    0    68.8  460.   863
## 2 credit_limit  0           1 4736.  2308.  855  3088  4622.  5873.
## 3 income        0           1  45.2   35.2  10.4  21.0  33.1   57.5
##   p100
## 1  1999
## 2 13913
## 3   187.
```

```
## $numeric
##
## -- Variable type: numeric -----
##   skim_variable n_missing complete_rate   mean    sd    p0    p25    p50    p75
## 1 debt          0           1  520.    460.    0    68.8  460.   863
## 2 credit_limit  0           1 4736.  2308.  855  3088  4622.  5873.
## 3 income        0           1  45.2   35.2  10.4  21.0  33.1   57.5
## # ... with 1 more variable: p100 <dbl>
```

```
credit_ch6 %>%
  select(debt, credit_limit, income) %>%
  cor()
```

```
##           debt credit_limit  income
## debt      1.0000000    0.8616973 0.4636565
## credit_limit 0.8616973    1.0000000 0.7920883
## income      0.4636565    0.7920883 1.0000000
```

## LC 6.2 (Objective 1)

(LC6.2) Conduct a new exploratory data analysis with the same outcome variable  $y$  being `debt` but with `credit_rating` and `age` as the new explanatory variables  $x_1$  and  $x_2$ . Remember, this involves three things:

- Most crucially: Looking at the raw data values.
- Computing summary statistics, such as means, medians, and interquartile ranges.
- Creating data visualizations.

What can you say about the relationship between a credit card holder's debt and their credit rating and age?

**Solution:**

- Most crucially: Looking at the raw data values.

```
credit_ch6 %>%
  select(debt, credit_rating, age) %>%
  head()
```

```
## # A tibble: 6 x 3
##   debt credit_rating age
##   <int>      <int> <int>
## 1   333         283   34
## 2   903         483   82
## 3   580         514   71
## 4   964         681   36
## 5   331         357   68
## 6  1151         569   77
```

Computing summary statistics, such as means, medians, and interquartile ranges.

```
credit_ch6 %>%
  select(debt, credit_rating, age) %>%
  my_skim() %>%
  print()
```

```
## -- Data Summary -----
##                               Values
## Name                         Piped data
## Number of rows                400
## Number of columns              3
## -----
## Column type frequency:
##   numeric                      3
## -----
## Group variables                None
##
## -- Variable type: numeric -----
##   skim_variable n_missing complete_rate mean    sd p0   p25  p50  p75 p100
## 1 debt          0           1 520.  460.   0  68.8 460. 863 1999
## 2 credit_rating 0           1 355.  155.  93 247. 344 437. 982
## 3 age           0           1  55.7  17.2 23  41.8 56  70  98


## $numeric
##
## -- Variable type: numeric -----
##   skim_variable n_missing complete_rate mean    sd p0   p25  p50  p75 p100
## 1 debt          0           1 520.  460.   0  68.8 460. 863 1999
## 2 credit_rating 0           1 355.  155.  93 247. 344 437. 982
## 3 age           0           1  55.7  17.2 23  41.8 56  70  98
```

```
credit_ch6 %>%
  select(debt, credit_rating, age) %>%
  cor()
```


```
##           debt credit_rating    age
## debt      1.000000000    0.8636252 0.001835119
## credit_rating 0.863625161    1.0000000 0.103164996
## age         0.001835119    0.1031650 1.000000000
```

- Creating data visualizations.

```
ggplot(credit_ch6, aes(x = credit_rating, y = debt)) +
  geom_point() +
  labs(
    x = "Credit rating", y = "Credit card debt (in $)",
    title = "Debt and credit rating"
  ) +
  geom_smooth(method = "lm", se = FALSE)
```



```
ggplot(credit_ch6, aes(x = age, y = debt)) +
  geom_point() +
  labs(
    x = "Age (in year)", y = "Credit card debt (in $)",
    title = "Debt and age"
  ) +
  geom_smooth(method = "lm", se = FALSE)
```



It seems that there is a positive relationship between one's credit rating and their debt, and very little relationship between one's age and their debt. There is a slight linear relationship between `age` and `credit_rating`.

### LC 6.3 (Objective 2)

(LC6.3) Fit a new simple linear regression using `lm(debt ~ credit_rating + age, data = credit_ch6)` where `credit_rating` and `age` are the new numerical explanatory variables  $x_1$  and  $x_2$ . Get information about the “best-fitting” regression plane from the regression table by applying the `get_regression_table()` function. How do the regression results match up with the results from your previous exploratory data analysis?

```
# Fit regression model:
debt_model_2 <- lm(debt ~ credit_rating + age, data = credit_ch6)

# Get regression table:
get_regression_table(debt_model_2)
```

```
## # A tibble: 3 x 7
##   term          estimate std_error statistic p_value lower_ci upper_ci
##   <chr>          <dbl>    <dbl>    <dbl>   <dbl>   <dbl>   <dbl>
## 1 intercept     -270.      44.8     -6.02     0    -358.    -181.
## 2 credit_rating   2.59      0.074    34.8     0      2.45     2.74
## 3 age           -2.35      0.668    -3.52     0     -3.66    -1.04
```

The coefficients for both new numerical explanatory variables  $x_1$  and  $x_2$ , `credit_rating` and `age`, are 2.59 and  $-2.35$  respectively, which means that `debt` and `credit_rating` are positively correlated, which matches

up with our explanatory analysis. However, `debt` and `age` are negatively correlated but in our exploratory analysis we surmised there was no relationship. When we account for credit rating, the debt tends to decrease with age, for one year increase in age, the debt on average decreases -2.35 with a credit rating held constant.

## Documenting software

- File creation date: 2022-06-05
- R version 4.1.3 (2022-03-10)
- `tidyverse` package version: 1.3.1
- `skimr` package version: 2.1.4
- ISLR package version: 1.4
  
- `moderndive` package version: 0.5.4