

Reading Notes: Moduli Spaces of Curves

LIU XIAOLONG

November 5, 2022

Contents

I	The basic facts of curves	7
1	Basic facts of general curves	9
1.1	Standard results	9
1.2	Automorphisms of curves	10
2	Families of curves	13
2.1	Families of smooth curves	13
2.2	Families of elliptic curves	14
3	Singularities of curves	15
3.1	δ -invariant	15
4	The varieties associated to curves	17
4.1	Jacobian variety of curves	17
4.1.1	Analytic approach	17
4.1.2	Algebraic approach	18
4.2	Picard varieties of curves	18
4.3	Basic fact of the determinantal varieties	20
4.4	The varieties of special linear series on a curve	20
4.5	The basic results of the Brill-Noether theory	22
II	The basic theory of moduli space of curves	23
5	\mathcal{M}_g be a Deligne-Mumford Stack for $g \neq 1$	25
5.1	\mathcal{M}_g be a stack for $g \neq 1$	25
5.2	For $g \geq 2$, \mathcal{M}_g be a Deligne-Mumford stack	26
5.3	First properties of \mathcal{M}_g for $g \geq 2$	27
5.4	Smoothness and dimension of \mathcal{M}_g for $g \geq 2$	27
5.5	For $g = 0$	28
6	Nodal curves	29
6.1	Basic facts of nodal curves	29
6.2	Genus fomula	30
6.3	The dualizing sheaf	30
6.3.1	The first way	31
6.3.2	The second way	31

6.3.3	The third way	31
6.4	Local structure of nodes	33
7	Stable curves	35
7.1	Basic facts of stable curves	35
7.2	Positivity of the dualizing sheaf	36
7.3	Families of stable curves	38
7.4	Rational tails and bridges	39
7.5	The stable model	39
7.5.1	The stable model of a single curve	39
7.5.2	The stable model of a family of curves	40
8	Deformation theory of nodal and stable curves	41
8.1	Elementary deformation theory and smooth objects	41
8.2	Elementary deformations of nodal and stable curves	43
8.3	Basic concept of Kuranishi family	47
8.4	The Hilbert scheme of ν -canonical curves	48
8.5	Construction of Kuranishi families	48
9	The stack of all curves	49
9.1	Families of all arbitrary curves	49
9.2	Algebraicity of the stack of all curves	50
9.3	Algebraicity of several stacks and boundedness of stable curves	52
9.4	The family of elliptic curves $\mathcal{M}_{1,1}$	53
10	Stable reduction: why $\overline{\mathcal{M}}_{g,n}$ is proper?	55
10.1	Proof of stable reduction in characteristic 0	56
10.2	Explicit stable reduction	58
10.3	Separatedness of $\overline{\mathcal{M}}_{g,n}$	60
11	Gluing and forgetful morphisms	63
11.1	Gluing morphisms	63
11.2	Boundary divisors of $\overline{\mathcal{M}}_g$	64
11.3	Forgetful morphisms	64
11.4	Universal family $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$	65
12	Irreducibility	67
12.1	Preliminaries–Branched coverings	67
12.2	Irreducibility over characteristic 0 using admissible covers	69
13	Projectivity	71
13.1	Kollár’s Criteria	71
13.2	Nefness of pluri-canonical bundles	73
13.3	Positivity via positivity theory	74
13.4	Projectivity via GIT, a sketch	74

III	Some geometry properties of the moduli space of curves	77
14	Preliminaries	79
14.1	Graphs and dual graphs	79
14.2	More on gluing morphisms	80
14.2.1	Gluing via graphs	80
14.2.2	Gluing functors	81
15	Picard groups of the moduli space of curves	85
16	The kodaira dimension of moduli space of curves	87
IV	Analytic construction via Teichmüller space	89
V	Cellular decomposition and cohomology	93
VI	Alterations and the moduli space of stable curves	97
VII	Appendix	101
A	Some basic result in scheme theory	103
A.1	Some corollaries of semi-continuity theorem	103
A.2	Artin approximation and its corollaries	103
A.3	Miscellany	104
B	Some results of resolution of singularities for surfaces	105
C	Basic theory of algebraic spaces and stacks	107
C.1	Some basic facts	107
C.2	Miscellany	108
	Bibliography	110

Part I

The basic facts of curves

Chapter 1

Basic facts of general curves

1.1 Standard results

Definition 1.1.1. A curve over k is a pure one-dimensional scheme C of finite type over k . If C is proper, we define the arithmetic genus (simply the genus) of C as $g(C) := g_a(C) = 1 - \chi(C, \mathcal{O}_C)$. By Review 3, if C is geometrically connected and geometrically reduced, this is equal to $h^1(C, \mathcal{O}_C)$.

Theorem 1.1.2 (St 0B5Y). Let k be a field. Let C be a proper scheme of dimension ≤ 1 over k . Let L be an invertible \mathcal{O}_X -module. Let C_i be the irreducible components of dimension 1. Then L is ample if and only if $\deg(L|_{C_i}) > 0$ for all i .

Theorem 1.1.3 (Serre duality of smooth curves). Let C be a smooth projective curve over k with canonical bundle $\omega_C = \Omega_C$, then for any vector bundle F we get

$$H^0(C, F^\vee \otimes \omega_C) \cong H^1(C, F)^\vee.$$

If we define the geometrical genus $g_e(C) = h^0(C, \omega_C)$ and if C is smooth projective curve which is geometrically connected and geometrically reduced, then $h^0(C, \mathcal{O}_C) = 1$. Hence by serre-duality we get $g_e(C) = g_a(C)$.

Theorem 1.1.4 (Riemann-Roch for smooth curves). Let C be a smooth projective curve over k with a line bundle L , then

$$\chi(C, L) = h^0(C, L) - h^0(C, \omega_C \otimes L^\vee) = \deg L + 1 - g.$$

Theorem 1.1.5 (Positivity of divisors on smooth curves). Let C be a smooth projective curve over k of genus g with a line bundle L , then

- (a) if $\deg L \geq 2g$, then L is base-point-free;
- (b) if $\deg L \geq 2g + 1$, then L is very ample;
- (c) if $\deg L > 0$, then L is ample.
- (d) if $\deg L < 0$, then $h^0(C, L) = 0$.

Proof. See the section IV.3 of [23] for the proof when k is algebraic closed. This is also right when k is not algebraic closed, see section 20.2 in [34].

Here we use another method to show (d) as a special case of [23] Ex.III.7.1. We just consider the case C is integral. If $\deg L < 0$, then L^{-1} is ample. Let $h^0(C, L) > 0$ and take a nonzero

$s \in H^0(C, L)$. As $H^0(C, L) = \text{Hom}(\mathcal{O}_C, L)$, we can get $- \times s : \mathcal{O}_C \rightarrow L$. As C integral, s must nonzero at the generic point, hence $- \times s : \mathcal{O}_C \rightarrow L$ is injective. Hence we get $L^{-1} \subset \mathcal{O}_C$. Let n such that L^{-n} generated by global sections, we get $L^{-n} \subset \mathcal{O}_C$. Hence $H^0(C, L^{-n}) \subset H^0(C, \mathcal{O}_C)$. Consider hilbert polynomial $\chi(L^{-n}) = \alpha n + \beta$ as $\deg \chi(L^{-n}) = \dim \text{supp}(L^{-1}) = \dim C = 1$. By Serre's vanishing theorem, we get for $n \rightarrow \infty$, we have $\chi(L^{-n}) = h^0(C, L^{-n}) \rightarrow \infty$. This is impossible since $h^0(C, \mathcal{O}_C) < \infty$. \square

Theorem 1.1.6 (Riemann-Hurwitz Theorem, St 0C1B). *Let $f : X \rightarrow Y$ be a separable morphism of smooth proper curves over a field k and if $k = H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y)$ and X and Y have genus g_X and g_Y , then*

$$2g_X - 2 = (2g_Y - 2) \deg(f) + \deg R$$

where R be the ramified divisor. Moreover, $\deg R = \sum_x d_x [\kappa(x) : k]$ where $d_x = \text{length}_{\mathcal{O}_{X,x}} \Omega_{X/Y,x}$. Of course if $\mathcal{O}_{X,x}$ is tamely ramified over $\mathcal{O}_{Y,f(x)}$ then $d_x = e_x - 1$. If not, we only have $d_x > e_x - 1$ where e_x is the ramification index.

1.2 Automorphisms of curves

Here we only consider smooth connected projective curves of genus g over an algebraically closed field k .

Proposition 1.2.1. *For $g = 0$, we get $\text{Aut}(\mathbb{P}_k^1) \cong \text{PGL}_2$. Moreover, if we consider all automorphisms fixed n points, then this group is finite if and only if $n \geq 3$.*

Proof. See [23] Example II.7.1.1, we get $\text{Aut}(\mathbb{P}_k^1) \cong \text{PGL}_2$. Moreover, all automorphisms fixed n points is finite if and only if $n \geq 3$ by easy linear algebra. \square

Proposition 1.2.2. *For curve C with $g = 1$, we get $\text{Aut}(C)$ is infinite group. Moreover, if we consider all automorphisms fixed n points, then this group is finite if and only if $n \geq 1$.*

Proof. In this case C is actually a group scheme of dimension 1 (by Picard varieties) and C can then act on C . Hence $C \subset \text{Aut}(C)$, hence infinite. Moreover, by [23] Corollary IV.4.7, if we fixed one point P_0 , then $\text{Aut}(C; P_0)$ is finite. \square

Proposition 1.2.3 (Hurwitz). *For curve C with $g \geq 2$, the group $\text{Aut}(C)$ is finite. Moreover, if k has characteristic 0, we have $\#(\text{Aut}(C)) \leq 84g - 84$.*

Proof. See [23] Ex.V.1.11 and Hurwitz's Automorphism Theorem. \square

Lemma 1.2.4 (St 0E67). *Let X be a smooth, proper, connected curve over k of genus g .*

- (a) *If $g \geq 2$, then $\text{Der}_k(\mathcal{O}_X, \mathcal{O}_X) = 0$;*
- (b) *If $g = 1$ and $D \in \text{Der}_k(\mathcal{O}_X, \mathcal{O}_X)$ is nonzero, then D does not fix any closed point of X ;*
- (c) *If $g = 0$ and $D \in \text{Der}_k(\mathcal{O}_X, \mathcal{O}_X)$ is nonzero, then D can fix at most 2 closed points of X .*

Remark 1.2.5. *We will say an element $D \in \text{Der}_k(\mathcal{O}_X, \mathcal{O}_X)$ fixes x if $D(\mathcal{I}) \subset \mathcal{I}$ where \mathcal{I} is the ideal sheaf of x .*

Sketch. As we have the canonical derivation $d : \mathcal{O}_X \rightarrow \Omega_{X/k}$, taking any $D \in \text{Der}_k(\mathcal{O}_X, \mathcal{O}_X)$ we get $D = f \circ d$ where $f \in \text{Hom}_{\mathcal{O}_X}(\Omega_{X/k}, \mathcal{O}_X)$ and $\deg(\Omega_{X/k}) = 2g - 2$.

- (a) *If $g \geq 2$, then $\deg(\Omega_{X/k}) > 0$. Hence*

$$\text{Hom}_{\mathcal{O}_X}(\Omega_{X/k}, \mathcal{O}_X) = \text{Hom}_{\mathcal{O}_X}(\mathcal{O}_X, T_{X/k}) = \Gamma(X, T_{X/k}) = 0,$$

hence $f = 0$;

(b)(c) We claim that the vanishing of f at $x \in X$ is equivalent to the statement that D fixes x . Indeed, by St 0C1E we get for the uniformizer $z \in \mathcal{O}_{X,x}$, dz is a basis of $\Omega_{X,x}$. Since $D(z) = f(dz)$, we conclude the claim.

If $g = 1$, then a nonzero f does not vanish anywhere. Hence by the claim, D does not fix any closed point of X . If $g = 0$, then a nonzero f vanishes in a divisor of degree 2. Hence by the claim, D can fix at most 2 closed points of X . \square

Lemma 1.2.6. *Let X be a proper scheme over a field k of dimension ≤ 1 , then the following are equivalent*

- (i) $\underline{\text{Aut}}(X)$ is geometrically reduced over k and has dimension 0;
- (ii) $\underline{\text{Aut}}(X) \rightarrow \text{Spec}(k)$ is unramified;
- (iii) $\text{Der}_k(\mathcal{O}_X, \mathcal{O}_X) = 0$.

Proof. See St 0DSW and St 0E6G. Note that these two lemmas can also give the results about automorphism groups of smooth connected curves. \square

Proposition 1.2.7. *Let C be a curve of genus g over a field k of characteristic 0, then for any non-trivial automorphism of C fixed at most $2g + 2$ points.*

Proof. See I.F-4 in [3] for now. To add. \square

Chapter 2

Families of curves

2.1 Families of smooth curves

Lemma 2.1.1 (Families of smooth curves). *Let $(S, f : C \rightarrow S)$ in \mathcal{M}_g for $g \geq 2$.*

- (i) $f_*\mathcal{O}_C = \mathcal{O}_S$;
- (ii) For $k > 1$ the sheaf $f_*(\Omega_{C/S}^1)^{\otimes k}$ is locally free of rank $(2k-1)(g-1)$ on S , and for any $g : S' \rightarrow S$, we get an isomorphism $g^*f_*(\Omega_{C/S}^1)^{\otimes k} \cong f'_*(\Omega_{C'/S'}^1)^{\otimes k}$. Moreover, $R^i f_*(\Omega_{C/S}^1)^{\otimes k} = 0, i > 0$;
- (iii) The sheaf $f_*\Omega_{C/S}^1$ is locally free of rank g on S , and for any $g : S' \rightarrow S$, we get an isomorphism $g^*f_*\Omega_{C/S}^1 \cong f'_*\Omega_{C'/S'}^1$. Moreover, $R^1 f_*(\Omega_{C/S}^1) = \mathcal{O}_S$ and $R^i f_*(\Omega_{C/S}^1) = 0, i > 1$;
- (iv) For $k \geq 3$, $(\Omega_{C/S}^1)^{\otimes k}$ is relative very ample.

Proof. (i) By definition, for all $s \in S$ the C_s is proper geometrically connected and geometrically reduced, then by Review 3 we get $H^0(C_s, \mathcal{O}_s) = \kappa(s)$, hence $\phi_s^0 : f_*\mathcal{O}_C \otimes \kappa(s) \rightarrow H^0(C_s, \mathcal{O}_s)$ is surjective. By Review 1 with $i = 0$, we get ϕ_s^0 is an isomorphism and $f_*\mathcal{O}_C$ is a line bundle. Now consider the natural map $\mathcal{O}_S \rightarrow f_*\mathcal{O}_C$ induce a surjective fiber map $\kappa(s) \rightarrow f_*\mathcal{O}_C \otimes \kappa(s)$ by seen

$$\kappa(s) \rightarrow f_*\mathcal{O}_C \otimes \kappa(s) \rightarrow H^0(C_s, \mathcal{O}_s) = \kappa(s).$$

Thus $\mathcal{O}_S \rightarrow f_*\mathcal{O}_C$ is surjective, hence an isomorphism.

(ii) For all $s \in S$ and $k > 1$ we get $H^1(C_s, (\Omega_{C_s/\kappa(s)}^1)^{\otimes k}) = H^0(C_s, (\Omega_{C_s/\kappa(s)}^1)^{\otimes(1-k)})^\vee = 0$ as $(\Omega_{C_s/\kappa(s)}^1)^{\otimes(1-k)}$ is anti-ample. Hence $H^i(C_s, (\Omega_{C_s/\kappa(s)}^1)^{\otimes k}) = 0$ for $i > 0$. Now use Review 1 we get $R^i f_*(\Omega_{C/S}^1)^{\otimes k} = 0, i > 0$.

On the other hand, by Riemann-Roch theorem, we get

$$h^0(C_s, (\Omega_{C_s/\kappa(s)}^1)^{\otimes k}) = \deg((\Omega_{C_s/\kappa(s)}^1)^{\otimes k}) + 1 - g = (2k-1)(g-1).$$

Use Review 1 again, we get $f_*(\Omega_{C/S}^1)^{\otimes k}$ is locally free of rank $(2k-1)(g-1)$ on S .

(iii) By Review 1 and the fact $H^i(C_s, \Omega_{C/S}^1 \otimes \kappa(s)) = 0, i > 1$ implies $R^i f_*\Omega_{C/S}^1 = 0, i > 1$. Now we use the duality $f_*\mathcal{H}om(F, \Omega_{C/S}^1) \cong \mathcal{H}om(R^1 f_*F, \mathcal{O}_S)$, then let $F = \Omega_{C/S}^1$. We get $f_*\mathcal{O}_C \cong (R^1 f_*\Omega_{C/S}^1)^*$. Hence $R^1 f_*\Omega_{C/S}^1 \cong (f_*\mathcal{O}_C)^* = \mathcal{O}_S^* \cong \mathcal{O}_S$.

By Review 1(ii) with $i = 1$, we get $\phi_s^0 : f_*\Omega_{C/S}^1 \otimes \kappa(s) \rightarrow H^0(C_s, \Omega_{C_s/\kappa(s)}^1)$ is surjective, hence an isomorphism. Then apply Review 1(i)-(ii) with $i = 0$ to imply $f_*\Omega_{C/S}^1$ is locally free of rank $h^0(C_s, \Omega_{C_s/\kappa(s)}^1) = g$.

(iv) Easy to see for any $s \in S$ the fiber $(\Omega_{C_s/\kappa(s)}^1)^{\otimes k}$ is very ample as $\deg(\Omega_{C_s/\kappa(s)}^1)^{\otimes k} = k(2g - 2) \geq 2g + 1$. Using noetherian approximation, we may let S is noetherian. Then use Review 4 and well done. \square

Proposition 2.1.2 (Flatness Criterion over Smooth Curves). *Let C be an integral and regular scheme of dimension 1 (e.g. the spectrum of a DVR or a smooth connected curve over a field) and $X \rightarrow C$ a qcqs morphism of schemes. A quasi-coherent \mathcal{O}_X -module F is flat over C if and only if every associated point of F maps to the generic point of C .*

2.2 Families of elliptic curves

This section are some preliminaries of the coarse moduli space of $\mathcal{M}_{1,1}$. Here we follows [32] 13.1 and for the basic theory of single elliptic curves, we refer [23] IV.4.

Chapter 3

Singularities of curves

3.1 δ -invariant

The the more details, see St 0C3Q and St 0C3Z.

Lemma 3.1.1 (St 0C3S). *Let (A, \mathfrak{m}) be a reduced 1-dimensional local ring of finite type over a field k . Let A' be the integral closure of A in the total ring of fractions of A . Then A' is a normal with $A \rightarrow A'$ is finite, and A'/A has finite length as an A -module.*

Definition 3.1.2. *Let A be a reduced 1-dimensional local ring of finite type over a field k . The δ -invariant of A is $\text{length}_A(A'/A)$ where A' is as in Lemma.*

Let X be a scheme locally of finite type over k . Let $x \in X$ such that $\mathcal{O}_{X,x}$ is reduced with dimension 1. The δ -invariant of X at x is the δ -invariant of $\mathcal{O}_{X,x}$.

Proposition 3.1.3 (St 0C3V). *Let A be a reduced 1-dimensional local ring of finite type over a field k . Then \hat{A} has the same δ -invariant as A and $A' \otimes_A \hat{A}$ is the integral closure of \hat{A} in its total ring of fractions.*

Proposition 3.1.4 (St 0C1R). *Let X be a reduced scheme locally finite type over a field of dimension 1 with normalization $f : \tilde{X} \rightarrow X$. Then $\mathcal{O}_X \subset f_*\mathcal{O}_{\tilde{X}}$ and $f_*\mathcal{O}_{\tilde{X}}/\mathcal{O}_X$ is a direct sum of skyscraper sheaves \mathcal{Q}_x in the singular points x and $\mathcal{Q}_x = (f_*\mathcal{O}_{\tilde{X}})_x/\mathcal{O}_{X,x}$ has finite length equal to the δ -invariant of X at x .*

Chapter 4

The varieties associated to curves

In most of cases of this chapter we focus on the proper reduced curves over an algebraically closed field k (may not irreducible and smooth). But when we let C smooth, we will automatically let C irreducible. When we consider \mathbb{C} we can use the language of Riemann surfaces via Serre's GAGA-principle.

4.1 Jacobian variety of curves

4.1.1 Analytic approach

We let C a smooth projective curve of genus g over \mathbb{C} . We follows [3].

► **Approach 1.** If ω, ω' are holomorphic forms on C , then checking locally we get $\omega \wedge \omega' = 0$ and $\int_C \sqrt{-1} \omega \wedge \bar{\omega} > 0$ ($\omega \neq 0$). Moreover we have $d\omega = 0$, hence $[\omega] \in H_{DR}^1(C) \cong H^1(C, \mathbb{C})$.

Choose a basis $\omega_1, \dots, \omega_g \in H^0(C, K)$ and $\gamma_1, \dots, \gamma_{2g} \in H_1(C, \mathbb{Z})$. We can define the period matrix

$$\Omega = (\Omega_1, \dots, \Omega_{2g})_{g \times 2g}, \Omega_i = \begin{pmatrix} \int_{\gamma_i} \omega_1 \\ \vdots \\ \int_{\gamma_i} \omega_g \end{pmatrix}.$$

Hence by construction we can see that $\Omega_1, \dots, \Omega_{2g}$ generates a lattice Λ in \mathbb{C}^g . Hence we define $J(C) := \mathbb{C}^g / \Lambda$ as the Jacobian variety of C .

► **Approach 2.** Or equivalently, consider $H_1(C, \mathbb{Z}) \hookrightarrow H^0(C, K)^\vee$ by $\gamma \mapsto \int_\gamma$, then $J(C) = H^0(C, K)^\vee / H_1(C, \mathbb{Z})$. Sometimes we call this Albanese torus when X be a compact Kähler manifold of higher dimension (See [24], for example).

► **Approach 3.** Let $\text{Pic}^0(C)$ be the degree 0 line bundles (or the kernel of the first Chern class in higher dimension). We can consider the exponential sequence and since C is compact, then $H^1(C, \mathbb{Z}) \rightarrow H^1(C, \mathcal{O}_C)$ is injective, hence $\text{Pic}^0(C) \cong H^1(C, \mathcal{O}_C) / H^1(C, \mathbb{Z})$. By some easy argument of Hodge theory (as [24] Corollary 3.3.6), it is a complex torus.

Proposition 4.1.1. *These three approaches defined the same variety associated to C .*

Proof. The first two approaches are the same trivially.

Now we consider the second and the third approaches. Actually this is by Serre duality $H^0(C, K)^\vee \cong H^1(C, \mathcal{O}_C)$ and Poincaré duality $H^1(C, \mathbb{Z}) \cong H_1(C, \mathbb{Z})$ which is compatible by trivial reasons. \square

4.1.2 Algebraic approach

Here we follow [23] and let C a smooth projective curve over an algebraically closed field k . Actually this is just the special case of the Picard variety.

Let T be a scheme over k and we let $\text{Pic}^0(C \times_k T)$ be the subgroup of $\text{Pic}(C \times_k T)$ consisting of invertible sheaves whose restriction to any fibers C_t for $t \in T$ has degree 0. Hence we can define $\text{Pic}^0(C/T) := \text{Pic}^0(C \times_k T)/p^*\text{Pic}T$ where $p : C \times_k T \rightarrow T$.

We can show that (omitted here) the functor

$$J(C) : (\text{Sch}/\text{Spec}k)^{\text{opp}} \rightarrow (\text{Sets}), T \mapsto \text{Pic}^0(C/T)$$

is represented by a k -scheme, we also denote $J(C)$. Here we find that for any points $x \in C(k)$, we find that $x : \text{Spec}k \rightarrow J(C)$ correspond to an element of $\text{Pic}^0(C)$. Hence this notation make sense.

Proposition 4.1.2. *The Jacobian variety $J(C)$ is a group variety over k .*

Proof. We define $e : \text{Spec}k \rightarrow J(C)$ correspond to $0 \in \text{Pic}^0(C/k)$ be the identity. Let $i : J(C) \rightarrow J(C)$ correspond to $\mathcal{L}_{\text{univ}}^{-1} \in \text{Pic}^0(C/J(C))$ be the inverse. Let $\mu : J(C) \times_k J(C) \rightarrow J(C)$ correspond to $p_1^*\mathcal{L}_{\text{univ}} \otimes p_2^*\mathcal{L}_{\text{univ}} \in \text{Pic}^0(C/J(C) \times_k J(C))$ be the multiple. Then the axiom of the group varieties is easy to check. \square

Proposition 4.1.3. *The Zariski tangent space $T_{J(C),0} \cong H^1(C, \mathcal{O}_C)$.*

Proof. Consider the dual number $T = k[\varepsilon]/(\varepsilon^2) \rightarrow J(C)$ by sending $\text{Spec}k$ to 0. By [23] Ex.III.4.6, we get

$$0 \rightarrow H^1(C, \mathcal{O}_C) \rightarrow \text{Pic}C[\varepsilon] \rightarrow \text{Pic}C \rightarrow 0.$$

Hence we win. \square

Proposition 4.1.4. *The Jacobian variety $J(C)$ is proper and nonsingular over k .*

Proof. Using valuation criterion, we need to extend the line bundle at a codimension 2 point of $C \times \text{Spec}R$. This is trivial. It is nonsingular by [23] Remark IV.4.10.9. \square

Proposition 4.1.5. *When C is a elliptic curve, then $J(C) \cong C$. In particular, C has a group structure.*

Proof. Omitted, see [23] Theorem IV.4.11. \square

4.2 Picard varieties of curves

For more detail about the general Picard scheme, we refer [28]. Here we focus on the theory on curves over an algebraically closed field k . We follow [5] in St 0B92.

Definition 4.2.1 (Picard functor). *Let $f : X \rightarrow S$ be a morphism in the big fppf-site $(Sch)_{fppf}$. Consider the functor*

$$\mathrm{Pic}_{X/S}^{Psh} : (Sch/S)_{fppf} \rightarrow (Sets), T \mapsto \mathrm{Pic}(X_T).$$

Let $\mathrm{Pic}_{X/S} := (\mathrm{Pic}_{X/S}^{Psh})^{fppf}$ the sheafification in fppf-topology.

Proposition 4.2.2. *(St 0B9N) Let $f : X \rightarrow S$ as in the definition which admits a section σ . Assume that $\mathcal{O}_T \cong f_{T,*}\mathcal{O}_{X_T}$ for all $T \in \mathrm{Ob}((Sch/S)_{fppf})$, then we have*

$$0 \longrightarrow \mathrm{Pic}T \longrightarrow \mathrm{Pic}X_T \xrightarrow{\sigma_T^*} \mathrm{Pic}_{X/S}(T) \longrightarrow 0$$

is exact and split by σ_T^ .*

sketch. • The left-exactness don't need the σ : WLOG we let $S = T$. If $f^*N \cong \mathcal{O}_X$, then $f_*f^*N \cong f_*\mathcal{O}_X \cong \mathcal{O}_S$ by assumption. Since N is locally trivial, we see that the canonical map $N \rightarrow f_*f^*N$ is locally an isomorphism (because $\mathcal{O}_S \rightarrow f_*f^*\mathcal{O}_S$ is an isomorphism by assumption). Hence we conclude that $N \rightarrow f_*f^*N \rightarrow \mathcal{O}_S$ is an isomorphism and we see that N is trivial. This proves the first arrow is injective.

The exactness in the middle is easy by fppf-descent of quasi-coherent sheaves.

• The right-exactness need the σ : Let $K(T) := \ker(\sigma_T^*)$, hence $\mathrm{Pic}(X_T) \cong \mathrm{Pic}T \oplus K(T)$ and $K(T) \subset \mathrm{Pic}_{X/S}(T)$. As $\mathrm{Pic}_{X/S}$ is the sheafification of K , we just need to show that K is a fppf-sheaf. I omitted here. \square

Lemma 4.2.3. *If C be a smooth projective curve over an algebraically closed field k , then the hypotheses of the previous Proposition are satisfied.*

Proof. We of course have a k -rational point (hence a section). Moreover, as $H^0(C, \mathcal{O}_C) = k$, by cohomology and base change we get $\mathcal{O}_T \rightarrow f_{T,*}\mathcal{O}_{C_T}$ is an isomorphism. \square

If C be a smooth projective curve over an algebraically closed field k with a closed point σ . Consider the functor

$$\mathrm{Pic}_{C/k,\sigma} : (Sch/k)^{opp} \rightarrow (Sets), T \mapsto \ker(\sigma_T^* : \mathrm{Pic}(C_T) \rightarrow \mathrm{Pic}T),$$

which is isomorphic to $\mathrm{Pic}_{C/k}$ before by the previous propositions. Hence we denote it by $\mathrm{Pic}_{C/k}$.

Theorem 4.2.4. *(St 0B9Z, St 0BA0) Let C be a smooth projective curve of genus g over an algebraically closed field k .*

- (i) *The functor $\mathrm{Pic}_{C/k}$ is representable by a group scheme, denote it also by $\mathrm{Pic}_{C/k}$;*
- (ii) *There is the disjoint decomposition of g -dimensional smooth proper varieties*

$$\mathrm{Pic}_{C/k} = \coprod_{d \in \mathbb{Z}} \mathrm{Pic}_{C/k}^d;$$

- (iii) *The closed points of $\mathrm{Pic}_{C/k}^d$ correspond to invertible \mathcal{O}_C -modules of degree d ;*
- (iv) *$\mathrm{Pic}_{C/k}^0$ is an open and closed subgroup scheme.*

Sketch. (iv) follows from the fact (ii). (iii) is trivial by definition. For the disjoint decomposition in (ii), by St 0B9T (locally constant of Euler characteristic) that for all $d \in \mathbb{Z}$ there is an open subfunctor $\mathrm{Pic}_{C/k}^d \subset \mathrm{Pic}_{C/k}$ whose value on a scheme T over k consists of those $L \in \mathrm{Pic}_{C/k,\sigma}(T)$ such that $\chi(C_t, L_t) = d + 1 - g$ and moreover we have $\mathrm{Pic}_{C/k,\sigma} = \coprod_{d \in \mathbb{Z}} \mathrm{Pic}_{C/k,\sigma}^d$. For (i) and the smoothness and properness of $\mathrm{Pic}_{C/k}$ we omitted, we refer St 0BA0. \square

4.3 Basic fact of the determinantal varieties

We refer [3] Chapter II.

Let $M = M(m, n) := \mathbb{A}_{\mathbb{C}}^{mn}$ be the variety of $m \times n$ matrix. Let $M_k \subset M$ be a subvariety consist of matrixes at most rank k . This is called the generic determinantal variety.

We also can let $\widetilde{M}_k = \{(A, W) \in M \times \text{Gr}(n - k, n) : AW = 0\}$ be a smooth connected subvariety of $M \times \text{Gr}(n - k, n)$. If $\pi : \text{Gr}(n - k, n) \times M \rightarrow M$, then $\pi(\widetilde{M}_k) = M_k$. Hence we can get M_k is irreducible of codimension $(m - k)(n - k)$.

Proposition 4.3.1. *We have $\text{Sing}(M_k) = M_{k-1}$.*

Proof. This need some calculation, I omit it here. \square

Theorem 4.3.2 (The Second Fundamental Theorem of Invariant Theory). *The ideal of M_k in M generated by all $(k + 1) \times (k + 1)$ minors and is radical.*

Proof. The proof of this fact relies on a detailed analysis of the homogeneous coordinate ring of the Grassmannian $\text{Gr}(k, N)$ under the Plücker embedding. See [3] Page 71-76. \square

Theorem 4.3.3 (The First Fundamental Theorem of Invariant Theory). *Let $G = \text{GL}(n, \mathbb{C})$ and act on $M(m, k) \times M(k, n)$ as $g(A, B) = (Ag^{-1}, gB)$. Let the multiplication $\mu : M(m, k) \times M(k, n) \rightarrow M(m, n)$, hence $\text{Im} \mu = M_k$. If we let $M(m, k) \times M(k, n) = \text{Spec} S$, then $M_k = \text{Spec} S^G$. Moreover, M_k is normal.*

Proof. Just some linear algebra, see [3] Page 77-79. \square

Theorem 4.3.4. *M_k is Cohen-Macaulay.*

Proof. This proof is much complicated by showing the cone over a Schubert variety is Cohen-Macaulay. See [3] Page 80-82. \square

Now we consider the general case of the determinantal variety. If X be a scheme over \mathbb{C} and $\phi : E \rightarrow F$ be a morphism of vector bundles of rank n, m . Let open U be the trivialization, hence $\phi|_U$ be a $m \times n$ matrix. Hence this induce $f : U \rightarrow M(m, n)$. Let $U_k = f^{-1}(M_k)$ and glue it together, we get $X_k(\phi) \subset X$. We call this the k -th determinantal variety. (Similarly we can get $\widetilde{X}_k(\phi)$. Also we have an intrinsical definition, see [3] Page 84.)

Hence $\text{codim}_X X_k(\phi) \leq (m - k)(n - k)$. Combining the previous local theorem and some algebra, we can get:

Proposition 4.3.5. *Let X be a smooth projective variety over \mathbb{C} with $\phi : E \rightarrow F$ be a morphism of vector bundles of rank n, m . If $\text{codim}_X X_k(\phi) = (m - k)(n - k)$, then $X_k(\phi)$ is Cohen-Macaulay.*

4.4 The varieties of special linear series on a curve

Let C be a smooth projective curve of genus g over \mathbb{C} . Let $C_d := C^d / S_d$ and easy to see that C_d be the set of effective divisors of degree d on C .

Fixed $p_0 \in C$ we first consider $C \rightarrow J(C)$ as $p \mapsto \int_{p_0}^p$, then this can extend to $\text{Div}(C) \rightarrow J(C)$ as

$$\sum_i p_i - \sum_j q_j \mapsto \sum_i \int_{p_0}^{p_i} - \sum_j \int_{p_0}^{q_j}.$$

Hence we have $\nu : \text{Div}^d(C) \rightarrow J(C)$. Hence we also can restrict to $\mu : C_d \rightarrow J(C)$.

Now using the Abel's theorem (see [3] Page 18), we have the following factorization

$$\begin{array}{ccccc} C_d & \hookrightarrow & \text{Div}^d(C) & \xrightarrow{-/\sim} & \text{Pic}^d(C) \\ & \searrow \mu & \downarrow \nu & \swarrow u & \\ & & J(C) & & \end{array}$$

► **The variety C_d^r :** Now let

$$C_d^r = \{D \in C_d : \dim |D| \geq r\}$$

with the variety-structure by using Brill-Noether matrix (see [3] IV.1, omitted here).

► **The variety $W_d^r(C)$:** Roughly speaking, we can let

$$W_d^r(C) = \{\text{parametrizing complete series } |D| : \deg D = d, h^0(C, \mathcal{O}_C(D)) \geq r+1\} \subset \text{Pic}_C^d.$$

Hence if we consider $f : C_d \rightarrow \text{Pic}_C^d$, then $f(C_d^r) = W_d^r(C)$.

The precise argument coming from the representability of the Picard variety Pic_C^d as follows. Let the degree d universal line bundle (or they called Poincaré line bundle) $\mathcal{L} = \mathcal{L}_{\text{univ}}$ on $C \times \text{Pic}_C^d$. Let $v : \text{Pic}_C^d \times C \rightarrow \text{Pic}_C^d$ be the projection.

Take E be an effective divisor on C such that $m := \deg E \geq 2g - d - 1$. Let $\Gamma = E \times \text{Pic}_C^d$ be a divisor on $C \times \text{Pic}_C^d$, by some sheaf-theoric argument (kind of flat base-change, see [3] IV.2.6), we have $R^1 v_* \mathcal{L}(\Gamma) = 0$ and $v_* \mathcal{L}(\Gamma)$ is locally free of rank $n = d + m - g + 1$. Hence we have

$$0 \rightarrow v_* \mathcal{L} \rightarrow K^0 := v_* \mathcal{L}(\Gamma) \xrightarrow{\gamma} K^1 := v_*(\mathcal{L}(\Gamma)/\mathcal{L}) \rightarrow R^1 v_* \mathcal{L} \rightarrow 0.$$

Hence $\ker \gamma = v_* \mathcal{L}$, $\text{coker} \gamma = R^1 v_* \mathcal{L}$ and $\text{rank} K^0 = n$, $\text{rank} K^1 = m$. Now we let

$$W_d^r(C) := X_{m+d-g-r}(\gamma) \text{ where } X = \text{Pic}_C^d.$$

Note that $W_d^r(C)$ is independent of the choice of E , see [3] Page 179.

By [3] Lemma IV.3.1, we get

Proposition 4.4.1. *The variety $W_d^r(C)$ represented the functor*

$$S \mapsto \left\{ L \in \text{Pic}^d(C \times S) \text{ such that the fitting rank of } R^1 \phi_* L \text{ is at least } g - d + r \right\}.$$

(For $\mathcal{F} \in \text{Coh}(X)$ with presentation $\mathcal{O}_X^n \xrightarrow{\gamma} \mathcal{O}_X^m \rightarrow \mathcal{F}$ the fitting rank of \mathcal{F} is the largest integer h such that the ideal in \mathcal{O}_X generated by the $(m - h + 1) \times (m - h + 1)$ minors of A vanishes.)

Also by [3] Lemma IV.3.1, we get the set of all \mathbb{C} -valued points of $W_d^r(C)$ is just

$$\{L \in \text{Pic}^d(C) : h^0(C, L) \geq r + 1\}.$$

Proposition 4.4.2. *For $r \geq d - g$, the each component of $W_d^r(C)$ has dimension greater or equal to the Brill-Noether number*

$$\rho(g, r, d) = g - (r + 1)(g - d + r).$$

Proof. Trivial by the local analysis above. □

Proposition 4.4.3. *For $f : C_d \rightarrow \text{Pic}_C^d$, we get $f^{-1}(W_d^r(C)) = C_d^r$.*

Proof. See [3] Proposition IV.3.4. □

► **The variety $G_d^r(C)$:** Roughly speaking, we can let

$$G_d^r(C) = \{\text{parametrizing } \mathfrak{g}_d^r \text{ of degree } d \text{ and dimension } r\}.$$

For precise definition, we let

$$G_d^r(C) := \tilde{X}_{m+d-g-r}(\gamma)$$

as previous construction. For any \mathbb{C} -valued point (L, V) where $L \in \text{Pic}^d(C)$ and V be a $(r+1)$ -dimensional subspace of $\ker \gamma_L$. By [3] Lemma IV.3.1 we get $\ker \gamma_L = H^0(C, L)$, we get

$$G_d^r(C)(\mathbb{C}) = \{(L, V) : L \in \text{Pic}^d(C), V \in \text{Gr}(r+1, H^0(C, L))\}.$$

Hence parametrizing all \mathfrak{g}_d^r . Actually $G_d^r(C)$ can also defined as a representable functor, we refer [3] page 182-183 and omit it here.

Now we collect some conclusions of these three varieties in [3] section IV.4.

Proposition 4.4.4. (i) *Every component of $G_d^r(C)$ has dimension at least equal to the Brill-Noether number $\rho = g - (r+1)(g-d+r)$;*

(ii) *Let $w = (L, W \subset H^0(C, L))$ be a point in $G_d^r(C)$ and consider the cup product*

$$\mu_{0,W} : W \otimes H^0(C, K \otimes L^{-1}) \rightarrow H^0(C, K) = H^1(C, \mathcal{O}_C)^\vee,$$

then $\dim T_w G_d^r(C) = \rho + \dim \ker \mu_{0,W}$. In particular, $G_d^r(C)$ is smooth at w of dimension ρ if and only if $\ker \mu_{0,W} = 0$;

(iii) *Let $L \in W_d^r(C) \setminus W_d^{r+1}(C)$ (hence $r \geq d-g$), then $T_L W_d^r(C) \cong (\text{Im } \mu_0)^\perp$ where $\mu_0 : H^0(C, L) \otimes H^0(C, K \otimes L^{-1}) \rightarrow H^0(C, K)$ be the cup product;*

(iv) *Let $L \in W_d^{r+1}(C)$, then $T_L W_d^r(C) = T_L \text{Pic}_C^d$;*

(v) *If $G_d^r(C)$ is smooth of dimension ρ , then $W_d^r(C)$ is Cohn-Macaulay, reduced and normal. If $d < g+r$ then $\text{Sing}(W_d^r(C)) = W_d^{r+1}(C)$.*

Proof. For (i),(ii), we refer [3] IV.4.1; for (iii),(iv), we refer [3] IV.4.2; for (v) we refer [3] IV.4.4. □

4.5 The basic results of the Brill-Noether theory

Part II

The basic theory of moduli space of curves

Chapter 5

\mathcal{M}_g be a Deligne-Mumford Stack for $g \neq 1$

Here we mainly consider the $g \neq 1$ curves.

Definition 5.0.1. *Let \mathcal{M}_g be the fibered category over schemes with objects of form $(S, f : C \rightarrow S)$ where S be a scheme and f be a proper smooth morphism such that every geometric fiber of S is a connected genus g curve. The morphisms are base-change.*

Our main result of this section is to prove that \mathcal{M}_g is a Deligne-Mumford stack for $g \geq 2$. For $g = 0$ we can run the same argument and when we just consider the stack over Sch/k where k be algebraically closed, we can get $\mathcal{M}_0 \cong BPG L_2$. Here we follows [1].

5.1 \mathcal{M}_g be a stack for $g \neq 1$

Lemma 5.1.1 (Descent for polarized schemes). *Let $\mathcal{P}ol$ be the category whose objects are pairs $(f : X \rightarrow Y, L)$ where f is a proper flat morphism and L is a relatively ample invertible sheaf. The morphism are diagrams of cartesian with isomorphic pullback of line bundles. Consider the fibered category $\mathcal{P}ol \rightarrow (Sch)$, then it has effective fppf-descent.*

Proof. See [32] 4.4.10. □

Theorem 5.1.2. *For $g \neq 1$, the fibered category \mathcal{M}_g is a stack.*

Proof. Consider $\mathcal{M}_g \rightarrow \mathcal{P}ol$ sends $C \rightarrow S$ to $(C \rightarrow S, \Omega_{C/S}^1)$ when $g \geq 2$ and $(C \rightarrow S, \Omega_{C/S}^{1, \otimes -1})$ when $g = 0$.

For a fppf covering $S' \rightarrow S$, then we get

$$\begin{array}{ccc} \mathcal{M}_g(S) & \longrightarrow & \mathcal{P}ol(S) \\ \downarrow & & \downarrow \cong \\ \mathcal{M}_g(S' \rightarrow S) & \longrightarrow & \mathcal{P}ol(S' \rightarrow S) \end{array}$$

Hence every object of $\mathcal{M}_g(S' \rightarrow S)$ is in the essential image of $\mathcal{M}_g(S)$. By the descent of sheaves used in $h_- \rightarrow h_+$ making it fully faithful. □

5.2 For $g \geq 2$, \mathcal{M}_g be a Deligne-Mumford stack

Now let $L_{C/S} = (\Omega_{C/S}^1)^{\otimes 3}$. By Lemma 2.1.1 (iv), for any family of smooth curves $p : D \rightarrow S$ we get a closed immersion $D \hookrightarrow \mathbb{P}(p_* L_{D/S})$ where $p_* L_{D/S}$ is locally free of rank $5g - 5$. Let $H = \underline{\text{Hilb}}_{\mathbb{P}^{5g-6}}^P$ where $P(t) = \deg(L_{C/S}^{\otimes t}) + 1 - g = (6g - 6)t + 1 - g$ be the Hilbert polynomial of $D_s \hookrightarrow \mathbb{P}_{\kappa(s)}^{5g-6}$. Let the universal closed subscheme:

$$\begin{array}{ccc} C & \hookrightarrow & H \times \mathbb{P}^{5g-6} \\ & \searrow \pi & \downarrow \\ & & H \end{array}$$

► **Claim 1.** There is a unique subscheme $H' \subset H$ consist of $h \in H$ such that

- (a) $C_h \rightarrow \text{Spec}(\kappa(h))$ is smooth and geometrically connected;
- (b) $C_h \hookrightarrow \mathbb{P}_{\kappa(h)}^{5g-6}$ is embedded by complete linear system $|L_{C_h/\kappa(h)}|$;
- (c) the line bundles $L_{C_{H'}/H'}$ and $\mathcal{O}_{C_{H'}}(1)$ differ by a pullback of a line bundle from H' (that is, there exists a line bundle N over H' such that $L_{C_{H'}/H'} \otimes p^* N = \mathcal{O}_{C_{H'}}(1)$).

Moreover, if $T \rightarrow H$ be a morphism such that (a)-(c) hold for the family $C_T \rightarrow T$, then $T \rightarrow H$ factors through H' .

Since the condition that a fiber of a proper morphism (of finite presentation) is smooth is an open condition on the target, the condition on H that C_h is smooth is open. Consider the Stein factorization (St 03H0) $C \rightarrow \tilde{H} := \underline{\text{Spec}}_H \pi_* \mathcal{O}_C \rightarrow H$ where $C \rightarrow \tilde{H}$ is proper with geometrically connected fibres and $\tilde{H} \rightarrow H$ is finite. As $\mathcal{O}_H \rightarrow \pi_* \mathcal{O}_C$ is a morphism between coherent sheaves, then the kernel and cokernel of it have closed supports. Hence $\tilde{H} \rightarrow H$ is an isomorphism over an open subscheme of H , which is precisely where the fibers of $C \rightarrow H$ are geometrically connected. Hence the points satiefies (a) be a open subscheme of H , denoted by $H_1 \subset H$.

By Review 2, there exists a locally closed subscheme $H_2 \subset H_1$ such that a morphism $T \rightarrow H_1$ factor through H_2 if and only if $L_{C_T/T}$ and $\mathcal{O}_{C_T}(1)$ differ by a pullback of a line bundle from T . In particular, (c) holds and for all $h \in H_2$, $L_{C_h/\kappa(h)} \cong \mathcal{O}_{C_h}(1)$.

For (b), let $\pi_2 : C_2 := C_{H_2} \rightarrow H_2$. Consider $\alpha : H^0(\mathbb{P}_{\mathbb{Z}}^{5g-6}, \mathcal{O}(1)) \otimes \mathcal{O}_{H_2} \rightarrow \pi_{2,*} \mathcal{O}_{C_2}(1)$ of vector bundles of rank $5g - 5$ on H_2 with fiber $\alpha_h : H^0(\mathbb{P}_{\kappa(h)}^{5g-6}, \mathcal{O}(1)) \rightarrow H^0(C_h, \mathcal{O}_{C_h}(1)) \cong H^0(C_h, L_{C_h/\kappa(h)})$. As they have the same rank, α_h is an isomorphism if and only if h is not in $\text{supp}(\text{coker}(\alpha))$. Let $H' = H_2 \setminus (\text{supp}(\text{coker}(\alpha)))$ and it satisfies (a)-(c) with that universal property.

► **Claim 2.** The group scheme $\text{PGL}_{5g-5} = \underline{\text{Aut}}(\mathbb{P}_{\mathbb{Z}}^{5g-6})$ act on H as: for $g \in \text{Aut}(\mathbb{P}_S^{5g-6})$ and $[D \subset \mathbb{P}_S^{5g-6}] \in H(S)$, we let $g \cdot [D \subset \mathbb{P}_S^{5g-6}] = [g(D) \subset \mathbb{P}_S^{5g-6}]$. As H' is PGL_{5g-5} -invariant, we claim that $\mathcal{M}_g \cong [H'/\text{PGL}_{5g-5}]$ be an algebraic stack. (See St 044O, St 04UV for quot stacks)

Consider $H' \rightarrow \mathcal{M}_g$ as $[D \subset \mathbb{P}_S^{5g-6}] \mapsto (D \rightarrow \mathbb{P}_S^{5g-6} \rightarrow S)$ is well defined by **Claim 1**. This morphism is PGL_{5g-5} -invariant, hence descends to $[H'/\text{PGL}_{5g-5}]^{pre} \rightarrow \mathcal{M}_g$ (**Why?**). We claim that this map is fully faithful. Indeed, for a family $p : D \rightarrow S$ in H' given by $D \subset \mathbb{P}_S^{5g-6}$, we get $\mathcal{O}_D(1) \cong L_{D/S} \otimes p^* M$ for some line bundle M on S . Use (b) we get

$$H^0(\mathbb{P}_{\mathbb{Z}}^{5g-6}, \mathcal{O}(1)) \otimes \mathcal{O}_S \rightarrow p_* \mathcal{O}_D(1) \cong p_*(L_{D/S} \otimes p^* M) \cong p_* L_{D/S} \otimes M$$

be an isomorphism. Then any automorphism of $D \rightarrow S$ induces an automorphism of $L_{D/S}$ and thus an automorphism of $p_* L_{D/S} \otimes M$, which induce an automorphism of \mathbb{P}_S^{5g-6} preserving D . By Theorem 5.1.2, \mathcal{M}_g be a stack, hence induce $[H'/\text{PGL}_{5g-5}] \rightarrow \mathcal{M}_g$ which is fully faithful since stackification is fully faithful. Finally we check that $[H'/\text{PGL}_{5g-5}] \rightarrow \mathcal{M}_g$ is essentially

surjective. As these are all stacks, then they satisfied effective descent of étale covering. Hence we just need to show that for any $p : D \rightarrow S$, there exists an étale covering $\{S_i \rightarrow S\}$ such that each D_{S_i} is in the image of $H'(S_i) \rightarrow \mathcal{M}_g(S_i)$. Actually since $L_{D/S}$ defined $D \hookrightarrow \mathbb{P}(p_*L_{D/S})$ and $p_*L_{D/S}$ is locally free of rank $5g - 5$, we let $\{S_i\}$ be open (zariski, hence étale) covering of S such that $(p_*L_{D/S})|_{S_i}$ are all free. Well done.

► **Claim 3.** The algebraic stack \mathcal{M}_g is a Deligne-Mumford stack.

By Theorem C.1.1, we just need to show for any smooth connected proper curve C over a algebraic closed field k , the group scheme $G := \underline{\text{Aut}}_k(C) = \text{Aut}(C)$ is finite and reduced. We find that $T_{G,e}$ can be identified with the automorphism group of the trivial first order deformation of C . Hence by Proposition 8.1.6, we get $T_{G,e} = H^0(C, T_C) = 0$, well done.

5.3 First properties of \mathcal{M}_g for $g \geq 2$

Proposition 5.3.1. As $\mathcal{M}_g \cong [H'/\text{PGL}_{5g-5}]$ and H' is locally of finite type, then \mathcal{M}_g is locally of finite type over \mathbb{Z} . As H' is noetherian, so is \mathcal{M}_g . So it is finite type over \mathbb{Z} .

Proposition 5.3.2. \mathcal{M}_g have affine diagonal. Indeed, since we have $\mathcal{M}_g \cong [H'/\text{PGL}_{5g-5}]$ which is an algebraic stack, then we have cartesian square

$$\begin{array}{ccc} H' \times \text{PGL}_{5g-5} & \longrightarrow & H' \times H' \\ \downarrow & & \downarrow \\ \mathcal{M}_g & \longrightarrow & \mathcal{M}_g \times \mathcal{M}_g \end{array}$$

As PGL_{5g-5} affine, then $\text{PGL}_{5g-5} \times H' \rightarrow H' \times H' \rightarrow H'$ affine, so is $\text{PGL}_{5g-5} \times H' \rightarrow H' \times H'$.

5.4 Smoothness and dimension of \mathcal{M}_g for $g \geq 2$

Proposition 5.4.1. If C is a smooth connected projective curve of genus $g \geq 2$ over k , then $\dim T_{\mathcal{M}_g, [C]} = 3g - 3$.

Proof. By Proposition 8.1.2, we get $T_{\mathcal{M}_g, [C]} = H^1(C, T_C)$. As $\deg T_C < 0$, we get $H^0(C, T_C) = 0$. So by Riemann-Roch we get

$$\dim T_{\mathcal{M}_g, [C]} = \dim H^1(C, T_C) = -\chi(T_C) = -\deg T_C + g - 1 = 3g - 3,$$

well done. □

Theorem 5.4.2. For $g \geq 2$, the Deligne-Mumford stack \mathcal{M}_g is smooth over \mathbb{Z} of relative dimension $3g - 3$.

Proof. Let a field k and a smooth projective connected curve $C \rightarrow \text{Spec}(k)$. Consider the following 2-diagram:

$$\begin{array}{ccccc} & & [C] & & \\ & \curvearrowright & & \searrow & \\ \text{Spec}(k) & \longrightarrow & \text{Spec}(A_0) & \longrightarrow & \mathcal{M}_g \\ & & \downarrow & & \downarrow f \\ & & \text{Spec}(A) & \longrightarrow & \text{Spec}(\mathbb{Z}) \end{array}$$

where $A \rightarrow A_0$ be a surjective maps of artinian local rings with residue field k with $k = \ker(A \rightarrow A_0)$. The map $\mathrm{Spec}(A_0) \rightarrow \mathcal{M}_g$ corresponds to a family of curves $C_0 \rightarrow \mathrm{Spec}(A_0)$ and a cartesian:

$$\begin{array}{ccccc} C & \longrightarrow & C_0 & \dashrightarrow & C' \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{Spec}(k) & \longrightarrow & \mathrm{Spec}(A_0) & \hookrightarrow & \mathrm{Spec}(A) \end{array}$$

of solid arrows. So to find the lifting, we just need to find the dashed arrows, that is, deformation of C along A . By Proposition 8.1.6(iii), there exists a cohomology class $ob_C \in H^2(C, T_C)$ such that this happens if and only if $ob_C = 0$. Hence this is right as C be a curve. Hence \mathcal{M}_g is smooth. By Theorem C.1.2, we get

$$\dim_{[C]} \mathcal{M}_g = \dim T_{\mathcal{M}_g, [C]} - \dim \mathrm{Aut}(C).$$

By the final step of the proof of the DM-ness of \mathcal{M}_g , we get $\dim \mathrm{Aut}(C) = 0$. Hence $\dim_{[C]} \mathcal{M}_g = \dim T_{\mathcal{M}_g, [C]} = 3g - 3$, well done. \square

5.5 For $g = 0$

By the same argument we can get \mathcal{M}_0 be a Deligne-Mumford stack.

Proposition 5.5.1. \mathcal{M}_0 may (not) be a stack over \mathbb{Z} isomorphic to $BPGL_2$.

Analysis. We should repeat the proof of the case $g \geq 2$. But when we consider $L_{C/S} = (\Omega_{C/S}^1)^{\otimes(-1)}$ for $f : C \rightarrow S$, we get $\deg(L_{C_s/\kappa(s)}) = 2$ and $f_* L_{C/S}$ is locally free of rank 3, which induce $C \hookrightarrow \mathbb{P}_S^2$. And with the Hilbert polynomial $p(t) = 2t + 1$. So we will use PGL_3 and get $\mathcal{M}_0 \cong [H'/PGL_3]$ for some subscheme $H' \subset \underline{\mathrm{Hilb}}_{\mathbb{P}^2}^{p(t)}$!

Actually by St 0C6U, we can not identify C_s over some $\kappa(s)$ with $\mathbb{P}_{\kappa(s)}^1$ since we may have **no invertible sheaves of odd degree!** If we can assume this, we have M of degree 1. The Hilbert polynomial $p(t) = t + 1$ in \mathbb{P}^1 and can show that it is \mathbb{P}^1 by some tricks (actually this is right for all linear subspaces). Hence

$$\mathcal{M}_0 \cong [\underline{\mathrm{Hilb}}_{\mathbb{P}^1}^{p(t)}/PGL_2] = [\mathrm{Grass}(2, 2)/PGL_2] = BPGL_2,$$

well done. \square

Corollary 5.5.2. When we consider the stack over Sch/k for any algebraically closed field k , we have $\mathcal{M}_0 \cong BPGL_2$.

Proof. Actually here we have invertible sheaves of degree 1. Hence any proper smooth curve of genus 0 be \mathbb{P}_k^1 . The Hilbert polynomial $p(t) = t + 1$ in \mathbb{P}_k^1 and can show that it is \mathbb{P}_k^1 by some tricks. Hence

$$\mathcal{M}_0 \cong [\underline{\mathrm{Hilb}}_{\mathbb{P}_k^1}^{p(t)}/PGL_2] = [\mathrm{Grass}_k(2, 2)/PGL_2] = BPGL_2,$$

well done. \square

Chapter 6

Nodal curves

6.1 Basic facts of nodal curves

The the more details, see St 0C46.

Definition 6.1.1 (Nodes). *Let C be a curve over k .*

- (a) *If k algebraically closed, we say that $p \in C(k)$ is a node is we have $\widehat{\mathcal{O}}_{C,p} \cong k[[x, y]]/(xy)$;*
- (b) *If k need not be algebraically closed, we say a closed point $p \in C$ is a node if there exists a node $p' \in C_{\bar{k}}$ over p .*

We say C be a nodal curve if every closed point is either smooth or nodal.

Proposition 6.1.2. *Let C be a curve over k . Consider the following statments.*

- (a) *$p \in C$ is a node;*
- (b) *$\kappa(p)/k$ is separable, $\widehat{\mathcal{O}}_{C,p} \cong \kappa(p)[[x, y]]/(ax^2 + bxy + cy^2)$ as a k -algebra where $ax^2 + bxy + cy^2$ is a nondegenerate quadratic form over $\kappa(p)$;*
- (c) *$\kappa(p)/k$ is separable and $\mathcal{O}_{C,p}$ is reduced, has δ -invariant 1.*

Then we have $(a) \Leftrightarrow (b) \Rightarrow (c)$.

Proof. See St 0C49 and St 0C4D.

We assume (a) \Leftrightarrow (b). Here by Lemma 3.1.3, we just need to consider the case $\mathcal{O}_{C,p} \cong \kappa(p)[[x, y]]/(ax^2 + bxy + cy^2)$ where $Q = ax^2 + bxy + cy^2$ is a nondegenerate quadratic form.

Case (I): If Q is split, we may let $\mathcal{O}_{C,p} \cong \kappa(p)[[x, y]]/(xy)$ after some coordinate transformation. Then we get

$$\widetilde{\mathcal{O}}_{C,p} \cong \kappa(p)[[x, y]]/(x) \times \kappa(p)[[x, y]]/(y);$$

Case (II): If not, in this case $c \neq 0$ and nondegenerate means $b^2 - 4ac \neq 0$. Hence $\kappa' = \kappa(p)[t]/(a + bt + ct^2)$ is a degree 2 separable extension of $\kappa(p)$. Then $t = y/x$ is integral over $\mathcal{O}_{C,p}$. and we conclude that

$$\widetilde{\mathcal{O}}_{C,p} = \kappa'[[x]]$$

with y mapping to tx on the right hand side.

In both cases one verifies by hand that the δ -invariant is 1, well done. □

Remark 6.1.3. (i) *As for a node $p \in C$ in a nodal curve C , we have $\kappa(p)/k$ is separable. As the two cases above, if p is of case (I), then $f^{-1}(p)$ has two points with residue fields $\kappa(p)$. If p is*

of case (II), then $f^{-1}(p)$ has only one point with residue field κ' , a degree 2 separable extension of $\kappa(p)$;

(ii) As in (i), all closed points of \tilde{C} is regular with separable residue fields over k . Hence \tilde{C} is smooth over k by St 00TV.

Proposition 6.1.4. *If C is a curve over k and $p \in C$ be a node. Then exists a finite separable field extension K/k , a point $P \in C_K$ over p and $\hat{\mathcal{O}}_{C_K, P} \cong K[[x, y]]/(xy)$.*

Proof. By Proposition 6.1.2(b), we get $\kappa(p)/k$ is separable, $\hat{\mathcal{O}}_{C, p} \cong \kappa(p)[[x, y]]/(ax^2 + bxy + cy^2)$ as a k -algebra where $Q = ax^2 + bxy + cy^2$ is a nondegenerate quadratic form over $\kappa(p)$. If Q is split, well done. If not, let $K = k[t]/(at^2 + bt + c)$ be a separable extension over k with Q split, well done. \square

6.2 Genus fomula

Let k be algebraically closed field now. Let C be a connected nodal projective curve over k . Let z_1, \dots, z_s be its nodes and C_1, \dots, C_t be its irreducible components.

By Proposition A.3.1(1) and (4), we get $\tilde{C} = \coprod_{i=1}^t \tilde{C}_i$ where \tilde{C}, \tilde{C}_i are normalizations. Let $f : \tilde{C} \rightarrow C$. By Proposition 3.1.4, we get a exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow f_* \mathcal{O}_{\tilde{C}} \rightarrow \bigoplus_{i=1}^s \mathcal{Q}_i \rightarrow 0$$

where \mathcal{Q}_i supported over z_i . Since by Proposition 6.1.2(c), we get $\mathcal{Q}_i = \kappa(z_i)$ as the δ -invariant are all 1, hence we get

$$0 \rightarrow \mathcal{O}_C \rightarrow f_* \mathcal{O}_{\tilde{C}} \rightarrow \bigoplus_{i=1}^s \kappa(z_i) \rightarrow 0.$$

Hence we get long exact sequence

$$0 \rightarrow \underbrace{H^0(C, \mathcal{O}_C)}_1 \rightarrow \underbrace{H^0(\tilde{C}, \mathcal{O}_{\tilde{C}})}_t \rightarrow \underbrace{\bigoplus_{i=1}^s \kappa(z_i)}_s \rightarrow \underbrace{H^1(C, \mathcal{O}_C)}_{g(C)} \rightarrow \underbrace{H^1(\tilde{C}, \mathcal{O}_{\tilde{C}})}_{\sum_{i=1}^t g(\tilde{C}_i)} \rightarrow 0$$

with the labels underneath indicating the dimensions.

Theorem 6.2.1 (Genus fomula). *With the situation as above, we get*

$$g(C) = \sum_{i=1}^t g(\tilde{C}_i) + s - t + 1.$$

Proof. Trivial by the argument above. \square

6.3 The dualizing sheaf

We have three way to see this. Consider C be a fixed nodal curve over k .

6.3.1 The first way

We find that C is locally complete intersection as we can check locally. As for a node $p \in C$, we have $\widehat{\mathcal{O}}_{C,p} \cong \kappa(p)[[x, y]]/(ax^2 + bxy + cy^2)$ for some nondegenerate quadratic form. By [31] Theorem 21.2(iii), we get $\mathcal{O}_{C,p}$ is a complete intersection over k . Hence by [23] Theorem III.7.11 (adjunction formula for l.c.i), if we embed it into \mathbb{P}^N , then we have $\omega_C \cong \omega_{\mathbb{P}^N} \otimes \bigwedge^{N-1}(\mathcal{I}/\mathcal{I}^2)$ where \mathcal{I} be the ideal sheaf. As this is locally complete intersection, this is a line bundle.

6.3.2 The second way

This is an abstract way of duality theory, see St 0E31 for more details. As C is locally complete intersection, then by St 0BVA we get C is Gorenstein. By St 0BS2, C must have a dualizing complex ω_C^* . By 0BFQ, as C is Gorenstein, ω_C^* is invertible. By C Cohen-Macaulay, $\omega_C^* = \omega_C[0]$. Hence we win.

6.3.3 The third way

We can explicit ω_C precisely. Let Σ be the set of nodes of C and let $U = C \setminus \Sigma$. Let the normalization $f : \tilde{C} \rightarrow C$ and $\tilde{\Sigma} := f^{-1}(\Sigma)$, $\tilde{U} := f^{-1}(U)$. Now \tilde{C} is smooth, then we have the dualizing sheaf (line bundle) $\Omega_{\tilde{C}}$. We get

$$0 \rightarrow \Omega_{\tilde{C}} \rightarrow \Omega_{\tilde{C}}(\tilde{\Sigma}) \rightarrow \mathcal{O}_{\tilde{\Sigma}} \rightarrow 0.$$

Actually the sections of $\Omega_{\tilde{C}}(\tilde{\Sigma})$ is the rational sections of $\Omega_{\tilde{C}}$ with at worst simple poles in $\tilde{\Sigma}$. Hence for any open $V \subset \tilde{C}$ and $y \in V \cap \tilde{\Sigma}$ we have the residue $\text{res}_y : \Gamma(V, \Omega_{\tilde{C}}(\tilde{\Sigma})) \rightarrow \kappa(y)$.

Definition 6.3.1. We define the subsheaf $\omega_C \subset f_*\Omega_{\tilde{C}}(\tilde{\Sigma})$ as for any open $V \subset C$ we have

$$\Gamma(V, \omega_C) = \left\{ s \in \Gamma(f^{-1}(V), \Omega_{\tilde{C}}(\tilde{\Sigma})) : \begin{array}{l} \text{for any } z_i \in V \cap \Sigma \\ \text{and } f^{-1}(z_i) = \{p_i, q_i\} \text{ with } \text{res}_{p_i}(s) + \text{res}_{q_i}(s) = 0 \end{array} \right\}.$$

Hence we get two exact sequences

$$\begin{aligned} 0 \rightarrow \omega_C \rightarrow f_*\Omega_{\tilde{C}}(\tilde{\Sigma}) &\longrightarrow \bigoplus_{z_i \in \Sigma} \kappa(z_i) \longrightarrow 0 \\ s &\longmapsto (\text{res}_{p_i}(s) - \text{res}_{q_i}(s)) \end{aligned}$$

and

$$\begin{aligned} 0 \rightarrow f_*\Omega_{\tilde{C}} \rightarrow \omega_C \rightarrow \bigoplus_{z_i \in \Sigma} \kappa(z_i) &\rightarrow 0 \\ s &\longmapsto (\text{res}_{p_i}(s)) \end{aligned}$$

Proposition 6.3.2. Let C be a nodal curve C over k .

- (a) If $g : C' \rightarrow C$ be an étale morphism, then $g^*\omega_C \cong \omega_{C'}$;
- (b) Conclude that ω_C be a line bundle.

Proof. (a) As the normalization commutes with étale base change (see St 03GE), we have the cartesian with normalizations

$$\begin{array}{ccc} \tilde{C}' & \xrightarrow{g'} & \tilde{C} \\ \downarrow f' & & \downarrow f \\ C' & \xrightarrow{g} & C \end{array}$$

By flat base change, we have the process $g^*\omega_C \subset g^*f_*\Omega_{\tilde{C}}(\tilde{\Sigma}) \cong f'_*(g')^*\Omega_{\tilde{C}}(\tilde{\Sigma}) = f'_*\Omega_{\tilde{C}'}(\tilde{\Sigma}')$. By definition and this process, we get $g^*\omega_C \cong \omega_{C'}$.

(b) Use Corollary A.2.3 and Proposition 6.1.4, there exists a separable extension K/k such that we get the common étale neighborhood as

$$\begin{array}{ccc} & (U, u) & \\ F \swarrow & & \searrow G \\ (C, p) & & (\text{Spec} K[x, y]/(xy), 0) \end{array}$$

Let $D = \text{Spec} K[x, y]/(xy)$ and normalization $\tilde{D} \cong \mathbb{A}_K^1 \sqcup \mathbb{A}_K^1$. Then $\Gamma(\tilde{D}, \Omega_{\tilde{D}}) = \Gamma(\mathbb{A}_K^1, \omega_{\mathbb{A}_K^1}) \times \Gamma(\mathbb{A}_K^1, \omega_{\mathbb{A}_K^1})$ and $(\frac{dx}{x}, -\frac{dy}{y})$ be a section of ω_D . As any section is of form $(f(x)\frac{dx}{x}, -g(y)\frac{dy}{y})$ where $f(0) = g(0)$, which is precisely the condition for $(f, g) \in \Gamma(\tilde{D}, \mathcal{O}_{\tilde{D}})$ to descend to a global function on D . In other words, $\omega_D \cong \mathcal{O}_D$ with generator $(\frac{dx}{x}, -\frac{dy}{y})$. By (a), we get $\omega_U = G^*\omega_D$, hence ω_U is a line bundle. As $F^*\omega_C = \omega_U$ be a line bundle, we use the descent theory and we win. \square

Proposition 6.3.3. *Let C be a proper nodal curve C over k , then ω_C be the dualizing line bundle of C .*

Proof. (See [2]) We may assume that k is algebraic closed. Choose a divisor $D = r_1 + \dots + r_h$ consisting of distinct smooth points of C , with the property that any component of C contains at least one of the r_i 's. We first claim that $H^1(\omega_C(D)) = 0$. Indeed, we get an exact sequence

$$0 \rightarrow (f_*\omega_{\tilde{C}}) \otimes \mathcal{O}_C(D) = f_*(\omega_{\tilde{C}}(D)) \rightarrow \omega_C(D) \rightarrow \bigoplus_{\text{nodes}} k \rightarrow 0.$$

Hence deduce a surjection

$$H^1(C, f_*(\omega_{\tilde{C}}(D))) = H^1(\tilde{C}, \omega_{\tilde{C}}(D)) \rightarrow H^1(C, \omega_C(D)).$$

As \tilde{C} is smooth, we get for any irreducible components and Serre duality in smooth case, we get $H^1(\tilde{C}, \omega_{\tilde{C}}(D)) = 0$ as D meets every irreducible components. Hence $H^1(\omega_C(D)) = 0$.

Next we deduce an exact sequence by using the claim as

$$H^0(C, \omega_C(D)) \rightarrow H^0(C, \omega_C(D)/\omega_C) \rightarrow H^1(C, \omega_C) \rightarrow H^1(C, \omega_C(D)) = 0.$$

For any $\phi \in H^1(C, \omega_C)$ we have some lifts $\phi' \in H^0(C, \omega_C(D)/\omega_C)$. We define the trace map as

$$\text{tr}_C : H^1(C, \omega_C(D)) \rightarrow k, \phi \mapsto 2\pi\sqrt{-1} \sum_{i=1}^l \text{res}_{r_i} \phi'$$

and this is well defined by using residue theorem (of definition). Perfect pairing is omitted. \square

Proposition 6.3.4. *Let C be a nodal curve C over k and $T \subset C$ be an irreducible component and D_T be the union of the intersections of T and another irreducible components, then $\omega_C|_T = \omega_T(D_T)$.*

Proof. Trivial by definition of the dualizing sheaves. \square

6.4 Local structure of nodes

Theorem 6.4.1 (Local structure of nodes). *Let $\pi : C \rightarrow S$ be a flat and finitely presented morphism such that every geometric fiber is a curve. Let $p \in C$ be a node in C_s . Then we have a following diagram*

$$\begin{array}{ccccc} (C, p) & \xleftarrow{\text{ét}} & (U, u) & \xrightarrow{\text{ét}} & (\text{Spec} A[x, y]/(xy - f), 0) \\ \downarrow & & \downarrow & \swarrow & \\ (S, s) & \xleftarrow{\text{ét}} & (\text{Spec} A, s') & & \end{array}$$

where each horizontal arrow is a residually-trivial pointed étale morphism and $f \in A$ is a function vanishing at s' .

Sketch. See [1] 5.2.12 or St 0CBY for more details.

Step 1. Reduce to S of finite type over \mathbb{Z} . Using noetherian approximation.

Step 2. Reduce to the case where $\widehat{\mathcal{O}}_{C_s, p} \cong \kappa(s)[[x, y]]/(xy)$. Just need to use Proposition 6.1.4 and since separable, we can find a étale neighborhood (S', s') such that $\kappa(s') = K$.

Step 3. Show that $\widehat{\mathcal{O}}_{C, p} \cong \widehat{\mathcal{O}}_{S, s}[[x, y]]/(xy - f)$ where $f \in \widehat{\mathfrak{m}}_s$. Using the Schlessinger's theorem in formal deformation theory to deduce a diagram similar as what we want at the completion level.

Step 4. Apply Artin approximation (Theorem A.2.2). Using Artin approximation to deduce our diagram from the completion level. \square

Corollary 6.4.2. *Let $\pi : C \rightarrow S$ be a flat and finitely presented morphism such that every geometric fiber is a curve, then the locus $C^{\leq \text{nod}} = \{p \in C : p \in C_{\pi(p)} \text{ either smooth or node}\} \subset C$ is open.*

Proof. First we know the smooth locus is open. If $p \in C_{\pi(p)}$ is a node, then by Theorem 6.4.1 we get an étale morphism $g : (U, u) \rightarrow (C, p)$. Then $p \in g(U) \subset C^{\leq \text{nod}}$ is open. \square


Corollary 6.4.3. *Let $\pi : C \rightarrow S$ be a proper flat and finitely presented morphism such that every geometric fiber is a curve, then the locus $S^{\leq \text{nod}} = \{s \in S : C_s \text{ is nodal}\} \subset S$ is open.*

Proof. As we find that

$$S^{\leq \text{nod}} = S \setminus \pi(C \setminus C^{\leq \text{nod}}).$$

By the previous Corollary and π is proper, then $S^{\leq \text{nod}}$ is open. \square

Remark 6.4.4. *Actually later we can prove that the stack $\mathcal{M}_g^{\leq \text{nod}}$ is a algebraic stack. But the main problem of $\mathcal{M}_g^{\leq \text{nod}}$ is that it is not separated and not of finite type. We can see the figure below for intuitive understanding:*



Corollary 6.4.5 (Comparison). *Let's compare $\omega_{C/Y}$ and $\Omega_{C/Y}^1$ where $\phi : C \rightarrow Y$ are a family of complex nodal curves. We will follow [2] X.2 and more general we can see [30] 6.4.2. Also, we will work on the complex topology.*

Pick a node p in some fiber, then by Theorem 6.4.1 we get near p we have the composition $\phi|_U : U \hookrightarrow \mathbb{C}^2 \times Y \rightarrow Y$ where U defined by $F := xy - f$. By adjunction formula we get the local generator of $\omega_{C/Y}$ is $F^{-1}dx \wedge dy \pmod{F}$. Using [30] Lemma 6.4.12, we get a homomorphism

$$\rho : \Omega_{C/Y}^1 \rightarrow \omega_{C/Y}$$

given by id if it near smooth points and $\rho(\alpha) = F^{-1}\alpha' \wedge dF \pmod{F}$ if near the nodes where α' is on $\mathbb{C}^2 \times Y \rightarrow Y$ restriction is α . Actually near nodes we have $\rho(dx) = xF^{-1}dx \wedge dy$ and $\rho(dy) = -yF^{-1}dx \wedge dy$. Now we consider

$$0 \rightarrow \ker \rho \rightarrow \Omega_{C/Y}^1 \xrightarrow{\rho} \omega_{C/Y} \rightarrow \text{coker} \rho \rightarrow 0.$$

• **Claim 1.** $\rho(\Omega_{C/Y}^1) = \mathcal{I}\omega_{C/Y}$ where \mathcal{I} be the ideal locally generated by x, y (locally ideal of that node).

Let S be the subspace correspond to \mathcal{I} , then for now $\text{coker} \rho = \omega_{C/Y} \otimes \mathcal{O}_S$. As locally near nodes we get $xy = f$ and $\rho(\Omega_{C/Y}^1)$ generated by $xF^{-1}dx \wedge dy$ and $yF^{-1}dx \wedge dy$, then \mathcal{I} be the ideal locally generated by x, y .

• **Claim 2.** *When Y be a single point, then $\ker \rho$ is the one-dimensional complex vector space generated by the class of $xdy = -ydx$.*


This is trivial by this construction.

Chapter 7

Stable curves

7.1 Basic facts of stable curves

An n -pointed curve is a curve C over a field k together with an ordered collection of k -rational points $p_1, \dots, p_n \in C$ which we call the marked points. A point $q \in C$ of an n -pointed curve is called special if q is a node or a marked point.



Definition 7.1.1. A n -pointed curve (C, p_1, \dots, p_n) over k is *prestable* if it is a geometrically connected, nodal and projective curve, and $p_1, \dots, p_n \in C(k)$ are distinct smooth points.

A n -pointed curve (C, p_1, \dots, p_n) over k is *semistable* if

- (a) it is prestable;
- (b) every smooth rational subcurve $\mathbb{P}^1 \subset C$ contains at least 2 special points;
- (c) C is not of genus 1 without marked points.

A n -pointed curve (C, p_1, \dots, p_n) over k is *stable* if

- (a) it is semistable;
- (b) every smooth rational subcurve $\mathbb{P}^1 \subset C$ contains at least 3 special points.

Remark 7.1.2. (1) Note that there are no n -pointed stable curve of genus g if $2g - 2 + n \leq 0$ by Proposition 7.1.4. We will often impose the condition that $2g - 2 + n > 0$ in order to exclude these special cases;

(2) An automorphism of a stable curve (C, p_1, \dots, p_n) is an automorphism $\alpha : C \rightarrow C$ such that $\alpha(p_i) = p_i$. We denote by $\text{Aut}(C, p_1, \dots, p_n)$ the group of automorphisms;

(3) For some general Riemann Roch theorem (such as OBS6) and the fact that the prestable curves are proper geometrically connected and geometrically reduced, then $\deg(\omega_C) = 2g - 2$.

Proposition 7.1.3. *Let (C, p_1, \dots, p_n) be an n -pointed nodal projective curve such that the points p_i are distinct and smooth. Let $\pi : \tilde{C} \rightarrow C$ be the normalization and $\tilde{p}_i \in \tilde{C}$ be the unique preimage of p_i and $\tilde{q}_1, \dots, \tilde{q}_m \in \tilde{C}$ be an ordering of the preimages of nodes. Then*

- (a) *(C, p_1, \dots, p_n) is stable if and only if every connected component of $(\tilde{C}, \{\tilde{p}_i\}, \{\tilde{q}_j\})$ is stable.*
- (b) *The group scheme $\underline{\text{Aut}}(C, \{p_i\})$ is an algebraic group.*
- (c) *$\underline{\text{Aut}}(C, \{p_i\})$ is naturally a closed scheme of $\underline{\text{Aut}}(\tilde{C}, \{\tilde{p}_i\}, \{\tilde{q}_j\})$ with the same connected component of identity.*

Proof. (a) Easy to see that we just need to verify that every smooth rational subcurve $\mathbb{P}^1 \subset C$ contains at least 3 special points if and only if every connected component of $(\tilde{C}, \{\tilde{p}_i\}, \{\tilde{q}_j\})$ have the same property. This is also trivial as we just need to consider the rational component of $(\tilde{C}, \{\tilde{p}_i\}, \{\tilde{q}_j\})$ and using the genus formula.

- (b)
- (c)

□

Proposition 7.1.4. *Let (C, p_1, \dots, p_n) be an n -pointed prestable curve. The following are equivalent*

- (i) *(C, p_1, \dots, p_n) is stable;*
- (ii) *$\text{Aut}(C, p_1, \dots, p_n)$ is finite; and*
- (iii) *$\omega_C(p_1 + \dots + p_n)$ is ample.*

Proof. (i) \Leftrightarrow (ii). By the results in Section 1.2 we get for smooth connected projective curve, its automorphism group is finite if and only if $\Leftrightarrow 2g - 2 + n > 0$ for (g, n) . Now consider the normalization $f : (\tilde{C}, \{\tilde{p}_i\}_{i=1}^n, \{\tilde{q}_j\}_{j=1}^{2s}) \rightarrow (C, p_1, \dots, p_n)$ with $\tilde{C} = \coprod_{j=1}^t \tilde{C}_j$. By Proposition 7.1.3 (a), we have (i) \Leftrightarrow for all j , $(\tilde{C}_j, \{\tilde{p}_i \in \tilde{C}_j\}_{i=1}^n, \{\tilde{q}_k \in \tilde{C}_j\}_{k=1}^{2s})$ is stable. As all \tilde{C}_j have marked points and use Proposition 7.1.3 (c), (ii) \Leftrightarrow for all j , $\text{Aut}(\tilde{C}_j, \{\tilde{p}_i \in \tilde{C}_j\}_{i=1}^n, \{\tilde{q}_k \in \tilde{C}_j\}_{k=1}^{2s})$ are finite. Hence by the case of smooth case, we win.

(i) \Leftrightarrow (iii). By Proposition A.3.2, 6.3.4 and consider the normalization $\pi : \tilde{C} \rightarrow C$, we get $\omega_C(p_1 + \dots + p_n)$ is ample if and only if $\pi^*\omega_C(p_1 + \dots + p_n)$ is ample if and only if for any irreducible components $T \subset \tilde{C}$, $\omega_C(p_1 + \dots + p_n)|_T = \omega_T(\sum_{p_i \in T} p_i + D_T)$ is ample. This latter condition holds precisely if each $\mathbb{P}^1 \subset \tilde{C}$ contains at least three points that lie over nodes or marked points (using Theorem 1.1.2) and we win. □

7.2 Positivity of the dualizing sheaf

Theorem 7.2.1. *For any n -pointed stable curve (C, p_1, \dots, p_n) , the bundle $(\omega_C(p_1 + \dots + p_n))^{\otimes k}$ is very ample for $k \geq 3$.*

Proof. We just prove the case of k is algebraically closed and no marked points. In this case we just need to show that its sections separates points and tangent vectors. We just need to show

- (i) for all closed points $x \neq y$, we have surjection

$$H^0(C, \omega_C^{\otimes k}) \twoheadrightarrow H^0(C, (\omega_C^{\otimes k} \otimes \kappa(x)) \oplus (\omega_C^{\otimes k} \otimes \kappa(y)));$$

- (ii) for all closed point x , we have surjection

$$H^0(C, \omega_C^{\otimes k}) \twoheadrightarrow H^0(C, \omega_C^{\otimes k} \otimes \mathcal{O}_C/I_x^2).$$

Hence for $x, y \in C(k)$ (maybe the same points) and their ideal I_x, I_y , we have

$$0 \rightarrow \omega_C^{\otimes k} \otimes I_x I_y \rightarrow \omega_C^{\otimes k} \rightarrow \omega_C^{\otimes k} \otimes \mathcal{O}_C / I_x I_y \rightarrow 0.$$

So we just need to show that $H^1(C, \omega_C^{\otimes k} \otimes I_x I_y) = 0$. By Serre duality, we need to show

$$\begin{aligned} H^1(C, \omega_C^{\otimes k} \otimes I_x I_y) &= H^0(C, (\omega_C^{\otimes k} \otimes I_x I_y)^\vee \otimes \omega_C) \\ &= H^0(C, \mathcal{H}om(\omega_C^{\otimes k} \otimes I_x I_y, \omega_C)) = \text{Hom}(I_x I_y, \omega_C^{\otimes(1-k)}) = 0. \end{aligned}$$

We need a case analysis on whether x, y are smooth or nodal.

If $x \in C$ is smooth, then $I_x = \mathcal{O}_C(-x)$ is invertible. If $x \in C$ is a node, consider the blowing up $\pi : C' \rightarrow C$ along x with $\pi^{-1}(x) = \{x_1, x_2\}$. Then for any line bundle L on C we claim that

$$\text{Hom}(I_x, L) \cong H^0(C', \pi^* L), \text{Hom}(I_x^2, L) \cong H^0(C', \pi^* L(x_1 + x_2)).$$

We just prove the first statement, the second is similar. First we have

$$0 \rightarrow \mathcal{O}_C \rightarrow \pi_* \mathcal{O}_{C'} \rightarrow \kappa(x) \rightarrow 0,$$

tensoring L we get

$$0 \rightarrow L \rightarrow \pi_* \mathcal{O}_{C'} \otimes L = \pi_* \pi^* L \rightarrow L(x) \rightarrow 0.$$

Hence we have

$$0 \rightarrow \text{Hom}(I_x, L) \rightarrow \text{Hom}(\pi^* I_x, \pi^* L) \xrightarrow{f} \text{Hom}(I_x, L(x)) = \text{Hom}(I_x / I_x^2, L(x)).$$

On the other hand, we have a short exact sequence

$$0 \rightarrow \pi^* L \rightarrow \pi^* L(x_1 + x_2) \rightarrow \pi^* L(x_1) \oplus \pi^* L(x_2) \rightarrow 0$$

inducing

$$0 \rightarrow H^0(C', \pi^* L) \rightarrow H^0(C', \pi^* L(x_1 + x_2)) \xrightarrow{g} \pi^* L(x_1) \oplus \pi^* L(x_2).$$

Let $J = \mathcal{O}_{C'}(-x_1 - x_2) \subset \mathcal{O}_{C'}$, we get

$$0 \rightarrow K \rightarrow \pi^* I_x \rightarrow J \rightarrow 0$$

where $\text{supp}(K) = \{x_1, x_2\}$ by checking locally. Since $\pi^* L$ is torsion free at x_1, x_2 , we have $\text{Hom}(K, \pi^* L) = 0$, so this defines an isomorphism

$$\text{Hom}(\pi^* I_x, \pi^* L) \cong \text{Hom}(J, \pi^* L) \cong H^0(C', \pi^* L(x_1 + x_2)).$$

We also have isomorphisms $I_x / I_x^2 \cong \pi_*(J / J^2)$ with $\text{Hom}(I_x / I_x^2, L(x)) \cong \text{Hom}(\pi_*(J / J^2), L(x))$. Actually $\text{Hom}(\pi_*(J / J^2), L(x)) \cong \pi^* L(x_1) \oplus \pi^* L(x_2)$ (**Why?**). Hence conclude these, we get

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}(I_x, L) & \longrightarrow & \text{Hom}(\pi^* I_x, \pi^* L) & \longrightarrow & \text{Hom}(I_x / I_x^2, L(x)) \\ & & \downarrow & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & H^0(C', \pi^* L) & \longrightarrow & H^0(C', \pi^* L(x_1 + x_2)) & \longrightarrow & \pi^* L(x_1) \oplus \pi^* L(x_2) \end{array}$$

Hence the claim is right.

Case (I). If x, y are all smooth points, then $\deg(\omega_C^{\otimes(1-k)}(x+y)) = (1-k)(2g-2) + 2 < 0$ for $k \geq 3$. Hence

$$\mathrm{Hom}(I_x I_y, \omega_C^{\otimes(1-k)}) = H^0(C, \omega_C^{\otimes(1-k)}(x+y)) = 0.$$

Case (II). If x is a node and y is a smooth point, then by the claim, we win.

Case (III.1). If $x = y$ is a node, then by the claim we get

$$\mathrm{Hom}(I_x^2, \omega_C^{\otimes(1-k)}) \cong H^0(C', \pi^* \omega_C^{\otimes(1-k)}(x_1 + x_2)).$$

Consider the normalization \tilde{C} of C (and C' , also), we consider an irreducible component $E \subset \tilde{C}$. Then $\pi^* \omega_C^{\otimes(1-k)}(x_1 + x_2)$ restrict to E has degree

$$(1-k)(2g_E - 2 + \#\{E \cap \tilde{\Sigma}\}) + \#(\{x_1, x_2\} \cap E)$$

is negative unless $k = 3$, $\{x_1, x_2\} \subset E$, E is a rational curve meeting the other components of C in exactly one other point. In this case the degree on E is zero. So this global section is determined by its value at the point of E meeting the other components of C . Since not every component of \tilde{C} , we win.

Case (III.2). If $x \neq y$ are all nodes, the blowing up $\varpi : C'' \rightarrow C$ along $\{x, y\}$. We can get similar conclusion

$$\mathrm{Hom}(I_x I_y, \omega_C^{\otimes(1-k)}) \cong H^0(C', \varpi^* \omega_C^{\otimes(1-k)}).$$

This is zero since in any irreducible of normalization has negative degree. \square

7.3 Families of stable curves

Definition 7.3.1. (1) A family of n -pointed nodal curves is a flat, proper and finitely presented morphism $C \rightarrow S$ of schemes with n sections $\sigma_1, \dots, \sigma_n : S \rightarrow C$ such that every geometric fiber C_s is a (reduced) connected nodal curve.

(2) A family of n -pointed stable curves (resp. semistable curves, prestable curves) is a family $C \rightarrow S$ of n -pointed nodal curves such that every geometric fiber $(C_s, \sigma_1(s), \dots, \sigma_n(s))$ is stable (resp. semistable, prestable).

Remark 7.3.2. (1) We can define the fibered category of groupoid $\overline{\mathcal{M}}_{g,n}$ as for any scheme S , define $\overline{\mathcal{M}}_{g,n}(S) = \{(C, \sigma_1, \dots, \sigma_n) \rightarrow S : \text{is a family of stable curves of genus } g\}$. Note also that since the geometric fibers are stable curves, the image of each σ_i is a divisor contained in the smooth locus and we can form the line bundle $\omega_{C/S}(\sum_i \sigma_i)$.

(2) We can define relative dualizing line bundle $\omega_{C/S}$ as $C \rightarrow S$ is l.c.i. By [22], we can get the following properties: (2.a) $\omega_{C/S}|_{C_s} = \omega_{C_s}$; (2.b) for any $f : T \rightarrow S$ we have $f^* \omega_{C/S} = \omega_{C \times_S T/T}$.

Proposition 7.3.3. Let $\pi : (C, \sigma_1, \dots, \sigma_n) \rightarrow S$ be a family of n -pointed stable curves of genus g . Let $L = \omega_{C/S}(\sum_i \sigma_i)$. If $k \geq 3$, then $L^{\otimes k}$ is relatively very ample and $\pi_* L^{\otimes k}$ is a vector bundle of rank $(2k-1)(g-1) + kn$.

Proof. Similar as the smooth case by using Riemann Roch and cohomology and base change. Omitted here. \square

Proposition 7.3.4 (Openness of stability). Let $\pi : (C, \sigma_1, \dots, \sigma_n) \rightarrow S$ be a family of n -pointed nodal curves. The locus of S such that $(C_s, \sigma_i(s))$ is stable is open.

Proof. As the locus such that $\sigma_i(s)$ is smooth is open, we just need to let this family is prestable. Using 7.1.4, we have two arguments:

Argument 1. Group scheme $\underline{\text{Aut}}(C/S, \sigma_i) \rightarrow S$ has upper semicontinuous dimension of fibers, then as stable locus is the locus such that it is dimension 0 locus. Hence open.


Argument 2. Using the openness of ample locus. \square

7.4 Rational tails and bridges

Definition 7.4.1. Let (C, p_1, \dots, p_n) be a n -pointed prestable curve. We say a smooth rational subcurve $E \cong \mathbb{P}^1 \subset C$ is


(i) a rational tail if E meets other irreducible components at exactly 1 time, and E contains no marked points;

(ii) a rational bridge if either E meets other irreducible components at exactly 2 time and contains no marked points, or E meets other irreducible components at exactly 1 time and contains exactly 1 marked point.



Remark 7.4.2. (1) C is stable if and only if it is prestable and has no rational tails and bridges;
 (2) C is semistable if and only if it is prestable and has no rational tails.


Or we can also have chain of rational tails and bridges like:



7.5 The stable model

7.5.1 The stable model of a single curve

Let (C, p_1, \dots, p_n) be a n -pointed prestable curve. Let $E_i \subset C$ are its rational tails and bridges. We let $C^{st} := C \setminus \bigcup_i E_i$ and let $\pi : C \rightarrow C^{st}$ be the induced map. Let $p'_i = \pi(p_i)$, then $(C^{st}, \{p'_i\})$ is a stable curve, which we call the stable model of $(C, \{p_i\})$ and $\pi : C \rightarrow C^{st}$ the stabilization morphism. Like this:



For the serious argument of contraction to the stable curves, we refer [5]: contracting rational tails (St 0E3G), contracting rational bridges (St 0E7M), contracting to a stable curve (St 0E7N). We omitted here.

7.5.2 The stable model of a family of curves

For a family of nodal curves, we also have the following conclusion.

Proposition 7.5.1. *Let $(C \rightarrow S, \sigma_i)$ be a family of n -pointed prestable curves. Then there exists a unique (up to isomorphism) morphism $\pi : C \rightarrow C^{st}$ such that*

- (a) $(C^{st} \rightarrow S, \{\sigma'_i\})$ is a n -pointed family of stable curves where $\sigma'_i = \pi \circ \sigma_i$;
- (b) for each $s \in S$, $(C_s, \{\sigma_i(s)\}) \rightarrow (C_s^{st}, \{\sigma'_i(s)\})$ is the stable model;
- (c) $\mathcal{O}_{C^{st}} = \pi_* \mathcal{O}_C$ and $R^1 \pi_* \mathcal{O}_C = 0$ and this remains true after base change by a morphism $S' \rightarrow S$ of schemes;
- (d) If $C \rightarrow S$ is a family of semistable curves, then $\omega_{C/S}(\sum_i \sigma_i)$ is the pullback of the relatively ample line bundle $\omega_{C^{st}/S}(\sum_i \sigma'_i)$.

Proof. See St 0E7B. □

Chapter 8

Deformation theory of nodal and stable curves

After some basic results over arbitrary fields, we will focus on the curves over \mathbb{C} . We mainly follows [2] chapter XI (all results over \mathbb{C}) and some results over arbitrary fields we follows [1]. Some basic result and proofs we follows [33]. Here we let $k[\varepsilon] := k[x]/(x^2)$.

8.1 Elementary deformation theory and smooth objects

Definition 8.1.1. *Let X be a scheme over k . A first order deformation of X is a scheme \mathcal{X} flat over $k[\varepsilon] = k[\varepsilon]/(\varepsilon^2)$ with $X \cong \mathcal{X} \times_{k[\varepsilon]} k$.*

We say \mathcal{X} is trivial if \mathcal{X} is isomorphic as first deformations to $\mathcal{X} \times_k k[\varepsilon]$, and locally trivial if there exists a Zariski-cover $X = \bigcup_i U_i$ such that $\mathcal{X}|_{U_i}$ is a trivial first order deformation of U_i , that is, $U_i \times_k k[\varepsilon] \cong \mathcal{X}|_{U_i}$ where $\mathcal{X}|_{U_i} \subset \mathcal{X}$ be a open subscheme with the same topology of U_i .

We let $\text{Def}(X)$ be the isomorphism classes of first order deformations of X and $\text{Def}^{lt}(X)$ be the isomorphism classes of locally trivial first order deformations of X .

Proposition 8.1.2 (See [1] D.1.11). *For a scheme X of finite type over k with affine diagonal, there is a bijection*

$$\text{Def}^{lt}(X) \leftrightarrow H^1(X, T_X).$$

In particular, if X_0 is smooth, then we have bijection

$$\text{Def}(X) \leftrightarrow H^1(X, T_X),$$

as every first order deformations of smooth affine schemes is trivial.

Sketch. For a locally trivial first order deformation

$$\begin{array}{ccc} X & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow \\ \text{Spec } k & \hookrightarrow & \text{Spec } k[\varepsilon] \end{array}$$

let affine covering $\{U_i\}$ of X such that $\mathcal{X}|_{U_i}$ be a trivial first order deformation. Hence we get isomorphisms $\phi_i : U_i \times_k k[\varepsilon] \cong \mathcal{X}|_{U_i}$. Let $\phi_{ij} := \phi_j^{-1}|_{U_{ij} \times_k k[\varepsilon]} \circ \phi_i|_{U_{ij} \times_k k[\varepsilon]}$ are automorphisms of first order defs, hence we get $\phi_{ij} \in \text{Hom}_{\mathcal{O}_{U_{ij}}}(\Omega_{U_{ij}/k}, \mathcal{O}_{U_{ij}})$. As they satisfies cocycle condition, we get $\{\phi_{ij}\} \in H^1(X, T_X)$ by Čech theory (this is independent on the choice of covering, see [33] Proposition 1.2.9). Converse is trivial. \square

Remark 8.1.3. For a locally trivial first order deformations ξ of X , we gives a class $\kappa(\xi) \in H^1(X, T_X)$ is called the Kodaira-Spencer class of ξ .

Definition 8.1.4. Consider a family of deformation of a smooth algebraic variety X over k

$$\begin{array}{ccccc} X & \longrightarrow & \mathcal{X} & \longleftarrow & \mathcal{X}_f \\ \downarrow & & \downarrow & & \downarrow \\ \text{Spec } k & \xleftarrow{s} & S & \xleftarrow{f} & \text{Spec } k[\varepsilon] \end{array}$$

\curvearrowright

hence we get

$$\kappa_{\mathcal{X}/S, s} : T_{S, s} \rightarrow H^1(X, T_X),$$

we called it Kodaira-Spencer map.

Definition 8.1.5. Let $A' \twoheadrightarrow A$ has square-free kernel and $X \rightarrow \text{Spec}(A)$ is flat. A deformation of $X \rightarrow \text{Spec}(A)$ over A' is $X' \rightarrow \text{Spec}(A')$ with $X' \times_{A'} A \cong X$. A morphism of deformations over A' is a morphism of schemes over A' restricting to the identity on X .

Proposition 8.1.6. Let $A' \twoheadrightarrow A$ has square-free kernel J . If $X \rightarrow \text{Spec}(A)$ is a smooth morphism of schemes where X has affine diagonal, then

- (a) the group of automorphisms of a deformation $X' \rightarrow \text{Spec}(A')$ of $X \rightarrow \text{Spec}(A)$ over A' is bijective to $H^0(X, T_{X/A} \otimes_A J)$;
- (b) If there exists a deformation of $X \rightarrow \text{Spec}(A)$ over A' , then the set of isomorphism classes of all such deformations is a torsor under $H^1(X, T_{X/A} \otimes_A J)$;
- (c) There is an element $ob_X \in H^2(X, T_{X/A} \otimes_A J)$ with the property that there exists a deformation of $X \rightarrow \text{Spec}(A)$ over A' if and only if $ob_X = 0$.

Proof. See [1] Proposition D.2.6. To add. \square

Back to smooth curves over \mathbb{C} pf genus g .

Theorem 8.1.7. Let $(C; q_1, \dots, q_n)$ be a n -pointed smooth n -pointed genus g curve over \mathbb{C} .

(i) We have

$$\text{Def}(C; q_1, \dots, q_n) \leftrightarrow H^1(C, T_C(-\sum_{i=1}^n q_i));$$

(ii) There exists a deformation

$$\phi : C \rightarrow (B, b_0), \sigma_i : B \rightarrow C \text{ such that } \chi : (C; q_1, \dots, q_n) \cong (\phi^{-1}(b_0), \sigma_i(b_0))$$

of $(C; q_1, \dots, q_n)$ such that the Kodaira-Spencer map

$$\kappa : T_{b_0} B \rightarrow H^1(C, T_C(-\sum_{i=1}^n q_i))$$

is an isomorphism and B is a polydisc of dimension $3g - 3 + n + h^0(C, T_C(-\sum_{i=1}^n q_i))$.

Proof. See [2] Theorem XI.2.12. To add. \square

8.2 Elementary deformations of nodal and stable curves

Lemma 8.2.1. *Let (C, p_1, \dots, p_n) be an n -pointed nodal, connected and projective curve over k with each $p_i \in C$ smooth. Let $\{q_1, \dots, q_s\}$ be the nodes of C . Let $(\tilde{C}, p_i, q'_j, q''_j)$ be the pointed normalization $\pi : \tilde{C} \rightarrow C$ and $\pi^{-1}(q_j) = \{q'_j, q''_j\}$. Then we have the spectral sequence*

$$E_2^{p,q} = H^p(C, \mathcal{E}xt_{\mathcal{O}_C}^q(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)) \Rightarrow \text{Ext}_{\mathcal{O}_C}^{p+q}(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)$$

such that induce the following exact sequence

$$\begin{array}{ccc} 0 & \longrightarrow & H^1(C, \mathcal{H}om_{\mathcal{O}_C}(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)) \\ & & \downarrow \\ 0 & \longleftarrow & \bigoplus_j \text{Ext}_{\mathcal{O}_{C,q_j}}^1(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}) \longleftarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C) \end{array}$$

Moreover, $\forall j$ we have $\text{Ext}_{\hat{\mathcal{O}}_{C,q_j}}^1(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}) = k$ and $\text{Ext}_{\mathcal{O}_C}^2(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C) = 0$.

Proof. By Grothendieck spectral sequence, we have

$$E_2^{p,q} = H^p(C, \mathcal{E}xt_{\mathcal{O}_C}^q(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)) \Rightarrow \text{Ext}_{\mathcal{O}_C}^{p+q}(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C).$$

As C is a curve, $E_2^{p,q} = 0$ for $p \geq 2$.

By [14] Propostion 2.3, we get an exact sequence

$$0 \rightarrow E_2^{1,0} \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C) \rightarrow E_2^{0,1} \rightarrow E_2^{2,0} \rightarrow 0.$$

As Ω_C is locally free away from nodes, $\mathcal{E}xt_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)$ is zero-dimensional sheaf supported only at nodes. Hence $E_2^{1,1} = 0$ and

$$\begin{aligned} E_2^{0,1} &= H^0(C, \mathcal{E}xt_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)) \\ &= \bigoplus_j \text{Ext}_{\mathcal{O}_{C,q_j}}^1(\Omega_{C,q_j}, \mathcal{O}_{C,q_j}) = \bigoplus_j \text{Ext}_{\hat{\mathcal{O}}_{C,q_j}}^1(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}). \end{aligned}$$

where $\hat{\Omega}_{C,q_j} = \Omega_{\hat{\mathcal{O}}_{C,q_j}}$. Hence we get that exact sequence.

Similarly $\mathcal{E}xt_{\mathcal{O}_C}^2(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)$ is zero-dimensional sheaf supported only at nodes, then

$$E_2^{0,2} = H^0(C, \mathcal{E}xt_{\mathcal{O}_C}^2(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)) = \bigoplus_j \text{Ext}_{\hat{\mathcal{O}}_{C,q_j}}^2(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}).$$

Write $\hat{\mathcal{O}}_{C,q_j} = k[[x, y]]/(xy)$ and consider the locally free resolution

$$0 \longrightarrow \hat{\mathcal{O}}_{C,q_j} \xrightarrow{\begin{pmatrix} y \\ x \end{pmatrix}} \hat{\mathcal{O}}_{C,q_j}^{\oplus 2} \xrightarrow{(dx, dy)} \Omega_{\hat{\mathcal{O}}_{C,q_j}} \longrightarrow 0$$

Hence we get $\text{Ext}_{\hat{\mathcal{O}}_{C,q_j}}^1(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}) = k$ and $\text{Ext}_{\hat{\mathcal{O}}_{C,q_j}}^2(\Omega_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}) = 0$. Hence $E_2^{0,2} = E_2^{1,1} = E_2^{2,0} = 0$ and $\text{Ext}_{\mathcal{O}_C}^2(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C) = 0$. \square

Proposition 8.2.2. *Let (C, p_1, \dots, p_n) be an n -pointed nodal, connected and projective curve over k with each $p_i \in C$ smooth. Let $\{q_1, \dots, q_s\}$ be the nodes of C . Let $(\tilde{C}, p_i, q'_j, q''_j)$ be the*

pointed normalization $\pi : \tilde{C} \rightarrow C$ and $\pi^{-1}(q_j) = \{q'_j, q''_j\}$. Then we have the following exact sequence

$$0 \rightarrow \text{Def}^{dt}(C) \rightarrow \text{Def}(C) \rightarrow \bigoplus_j \text{Def}(\hat{\mathcal{O}}_{C,q_j}) \rightarrow 0$$

and

$$\text{Def}^{dt}(C) \cong \text{Def}(\tilde{C}, p_i, q'_j, q''_j) \cong H^1(\tilde{C}, T_{\tilde{C}}(-\sum_i p_i - \sum_j (q'_j + q''_j))),$$

$$\text{Def}(C) \cong \text{Ext}^1_{\mathcal{O}_C}(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C),$$

$$\text{Def}(\hat{\mathcal{O}}_{C,q_j}) \cong \text{Ext}^1_{\hat{\mathcal{O}}_{C,q_j}}(\Omega^1_{\hat{\mathcal{O}}_{C,q_j}}, \hat{\mathcal{O}}_{C,q_j}) \cong k.$$

Under these identifications, this exact sequence corresponds to the exact sequence in the Lemma.

Sketch. WLOG again we let $n = 0$. If $\mathcal{C} \rightarrow \text{Spec}k[\varepsilon]$ is a locally trivial first order deformation of C , each node q_j extend to a section $\tilde{q}_j : \text{Spec}k[\varepsilon] \rightarrow \mathcal{C}$. The pointed normalization of \mathcal{C} along the sections \tilde{q}_j is a first order deformation of the (possible disconnected) pointed normalization $(\tilde{C}, p_i, q'_j, q''_j)$. This gives a map $\text{Def}^{dt}(C) \rightarrow \text{Def}(\tilde{C}, p_i, q'_j, q''_j)$. The inverse is provided by gluing the sections of a first order deformation of $(\tilde{C}, p_i, q'_j, q''_j)$ along nodes.

If $\mathcal{C} \rightarrow \text{Spec}k[\varepsilon]$ is a first order deformation of C , then ideal sheaf I of $C \rightarrow \mathcal{C}$ is $I = I/I^2 \cong \mathcal{O}_C$. The right exact sequence

$$I/I^2 \rightarrow \Omega_{\mathcal{C}/k} \rightarrow \Omega_{C/k} \rightarrow 0$$

is left exact at every smooth point of C . As $C \rightarrow \text{Spec}k$ is generically smooth and it follows that $\mathcal{O}_C \cong I/I^2 \rightarrow \Omega_{\mathcal{C}/k}$ is generically injective, hence injective. Hence this defines $\text{Ext}^1_{\mathcal{O}_C}(\Omega_C, \mathcal{O}_C)$. This is bijective (one can see [2] section XI.3). \square

Remark 8.2.3. Hence we also have the Kodaira-Spencer map for some $\mathcal{C} \rightarrow (S, s)$ as

$$\kappa_{S,s} : T_{S,s} \rightarrow \text{Ext}^1_{\mathcal{O}_C}(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C).$$

Remark 8.2.4. Let C be a nodal curve over \mathbb{C} and let $p \in C$ be a node with normalization N and preimages $\{p_1, p_2\}$.

• **Claim 1.** $\text{Ext}^1(\Omega_{C,p}, \mathcal{O}_{C,p}) \cong \bigwedge^2(\mathfrak{m}_p/\mathfrak{m}_p^2) \otimes_{\mu_2} \tau$ where $\mu_2 = \{\pm 1\}$ and τ be the set consisting of the two possible orderings of the branches of C at p .

I omit this, see [2] page 180.

• **Claim 2.** $\text{Ext}^1(\Omega_{C,p}, \mathcal{O}_{C,p}) \cong T_{N,p_1} \otimes T_{N,p_2}$.

Trivial by Claim 1 and $\mathfrak{m}_p/\mathfrak{m}_p^2 = T_{C,p} = T_{N,p_1} \oplus T_{N,p_2}$ and $\bigwedge^2 T_{N,p_1} \oplus T_{N,p_2}$ identify with $T_{N,p_1} \otimes T_{N,p_2}$ depends on the choice of an ordering of the two summands, we win.

Remark 8.2.5. Let $(C; p_1, \dots, p_n)$ be an n -pointed nodal curve over \mathbb{C} for simplicity, and let $W = \{w_1, \dots, w_l\}$ be some set of nodes of C . Let $f : N \rightarrow C$ be the partial normalization at these nodes with $f^{-1}(w_i) = \{r_i, q_i\}$. Let $D = \sum_i p_i$ with inverse \tilde{D} and $E = \sum(r_i + q_i)$.

• **Claim 1.** $\mathcal{H}om(\Omega^1_C, \mathcal{O}_C(-D)) \cong f_* \mathcal{H}om(\Omega^1_N, \mathcal{O}_N(-\tilde{D} - E))$.

This is trivially true at points away from W , so we just need to consider the points in W . Pick any $w_i \in W$, we get $\text{Hom}(\Omega^1_{C,w_i}, \mathcal{O}_{C,w_i}) = \text{Hom}(\mathcal{I}_{w_i} \omega_{C,w_i}, \mathcal{O}_{C,w_i})$ by Corollary 6.4.5. As $\mathcal{I}_{w_i} \omega_{C,w_i} = \omega_{N,r_i} \oplus \omega_{N,q_i}$ and $\mathcal{I}_{w_i} = \mathcal{O}_{N,r_i}(-r_i) \oplus \mathcal{O}_{N,q_i}(-q_i)$ (*Why?*), we get

$$\text{Hom}(\Omega^1_{C,w_i}, \mathcal{O}_{C,w_i}) = \bigoplus_{p=r_i, q_i} \text{Hom}(\omega_{N,p}, \mathcal{O}_{N,p}(-p)),$$

hence $\mathcal{H}om(\Omega_C^1, \mathcal{O}_C(-D)) \cong f_* \mathcal{H}om(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E))$.

•**Claim 2.** We have

$$\begin{aligned} 0 \rightarrow \text{Ext}^1(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E)) &\rightarrow \text{Ext}^1(\Omega_C^1, \mathcal{O}_C(-D)) \rightarrow \\ &\bigoplus_{w_i \in W} \text{Ext}^1(\Omega_{C, w_i}^1, \mathcal{O}_{C, w_i}) \rightarrow 0. \end{aligned}$$

By Claim 1, we get $H^1(N, \mathcal{H}om(\Omega_C^1, \mathcal{O}_C(-D))) \cong H^1(N, \mathcal{H}om(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E)))$. Hence by Lemma 8.2.1, we get

$$\begin{array}{ccccc} 0 & \longrightarrow & H^1(N, \mathcal{H}om(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E))) & \longrightarrow & \text{Ext}^1(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E)) \\ & & \parallel & & \downarrow \\ 0 & \longrightarrow & H^1(N, \mathcal{H}om(\Omega_C^1, \mathcal{O}_C(-D))) & \longrightarrow & \text{Ext}^1(\Omega_C^1, \mathcal{O}_C(-D)) \\ & & & & \downarrow \\ & \longrightarrow & \bigoplus_{w \in \text{Sing}(C), w \notin W} \text{Ext}^1(\Omega_{C, w}^1, \mathcal{O}_{C, w}) & \longrightarrow & 0 \\ & & \downarrow & & \\ & \longrightarrow & \bigoplus_{w \in \text{Sing}(C)} \text{Ext}^1(\Omega_{C, w}^1, \mathcal{O}_{C, w}) & \longrightarrow & 0 \end{array}$$

hence we get

$$\begin{aligned} 0 \rightarrow \text{Ext}^1(\Omega_N^1, \mathcal{O}_N(-\tilde{D} - E)) &\rightarrow \text{Ext}^1(\Omega_C^1, \mathcal{O}_C(-D)) \rightarrow \\ &\bigoplus_{w_i \in W} \text{Ext}^1(\Omega_{C, w_i}^1, \mathcal{O}_{C, w_i}) \rightarrow 0. \end{aligned}$$

Note that the term on the left classifies first-order deformations which are locally trivial at the nodes belonging to W , and the one on the right classifies first-order smoothings of these nodes.

•**Claim 3.** $\bigoplus_{w_i \in W} \text{Ext}^1(\Omega_{C, w_i}^1, \mathcal{O}_{C, w_i}) = \bigoplus_{i=1}^l T_{N, r_i} \otimes T_{N, q_i}$.

By claims in Remark 8.2.4, this is trivial.

Here is a similar result as before over \mathbb{C} via analytic GAGA.

Theorem 8.2.6. Let $(C; p_1, \dots, p_n)$ be an n -pointed nodal curve of genus g over \mathbb{C} . There exists a deformation

$$\phi : \mathcal{C} \rightarrow (B, b_0), \sigma_i : B \rightarrow \mathcal{C} \text{ such that } \chi : (C; p_1, \dots, p_n) \cong (\phi^{-1}(b_0), \sigma_i(b_0))$$

of $(C; p_1, \dots, p_n)$ such that the Kodaira-Spencer map

$$\kappa : T_{b_0} B \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)$$

is an isomorphism and B is a polydisc of dimension $3g - 3 + n + \dim \text{Hom}(\Omega_C, \mathcal{O}_C)$.

Finally, if s is the number of nodes of C , one can choose coordinates t_1, \dots, t_s, \dots on B , vanishing at b_0 , in such a way that the locus parameterizing deformations which are locally trivial at the i -th node is $t_i = 0$; in particular, the locus parameterizing singular curves is $t_1 \cdots t_s = 0$.

Proof. See [2] Theorem XI.3.17. To add. □

Back to the general case.

Proposition 8.2.7. *Let (C, p_1, \dots, p_n) be an n -pointed nodal, connected and projective curve over k with each $p_i \in C$ smooth. Let $A' \rightarrow A$ be a surjection of artinian local k -algebras with residue field k such that $J = \ker(A' \rightarrow A)$ satisfies $\mathfrak{m}_{A'} J = 0$. If $C_A \rightarrow \operatorname{Spec}(A)$ be a family of nodal curves such that $C \cong C_A \times_A k$, then*

(a) *The group of automorphisms of a deformation $C_{A'} \rightarrow \operatorname{Spec}(A')$ of $C_A \rightarrow \operatorname{Spec}(A)$ over A' is bijective to $\operatorname{Ext}_{\mathcal{O}_C}^0(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C \otimes_k J)$;*

(b) *If there exists a deformation of $C_A \rightarrow \operatorname{Spec}(A)$ over A' , then the set of isomorphism classes of all such deformations is a torsor under $\operatorname{Ext}_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C \otimes_k J)$;*

(c) *There is an element $ob_{C_A} \in \operatorname{Ext}_{\mathcal{O}_C}^2(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C \otimes_k J)$ with the property that there exists a deformation of $C_A \rightarrow \operatorname{Spec}(A)$ over A' if and only if $ob_{C_A} = 0$.*

Proof. To add. □

Lemma 8.2.8 (St 0E68). *Let k be an algebraically closed field. Let X be an at-worst-nodal, proper, connected 1-dimensional scheme over k . Let $f : \tilde{X} \rightarrow X$ be the normalization. Let $S \subset \tilde{X}$ be the set of points where f is not an isomorphism.*

$$\operatorname{Der}_k(\mathcal{O}_X, \mathcal{O}_X) = \{D' \in \operatorname{Der}_k(\mathcal{O}_{\tilde{X}}, \mathcal{O}_{\tilde{X}}) : D' \text{ fixed every } x' \in S\}.$$

Proof. Let $x \in X$ be a node with the preimage $x', x'' \in \tilde{X}$. Pick two uniformizers u, v in $\mathcal{O}_{\tilde{X}, x'}$ and $\mathcal{O}_{\tilde{X}, x''}$, respectively. Hence we have

$$0 \rightarrow \mathcal{O}_{X, x} \rightarrow \mathcal{O}_{\tilde{X}, x'} \times \mathcal{O}_{\tilde{X}, x''} \rightarrow k \rightarrow 0,$$

thus we can view u, v as elements of $\mathcal{O}_{X, x}$ with $uv = 0$.

Since (u) is annihilator of v in $\mathcal{O}_{C, x}$ and vice versa, we see that $D(u) \in (u)$ and $D(v) \in (v)$. As $\mathcal{O}_{\tilde{C}, x'} = k + (u)$ we conclude that we can extend D to $\mathcal{O}_{\tilde{C}, x'}$ and moreover the extension fixes x' . This produces a D' in the right hand side of the equality. Conversely, given a D' fixing x' and x'' we find that D' preserves the subring $\mathcal{O}_{C, x} \subset \mathcal{O}_{\tilde{C}, x'} \times \mathcal{O}_{\tilde{C}, x''}$ and this is how we go from right to left in the equality. □

Proposition 8.2.9. *Let (C, p_1, \dots, p_n) be an n -pointed stable curve of genus g over k . Then*

$$\dim_k \operatorname{Ext}_{\mathcal{O}_C}^i \left(\Omega_C \left(\sum_i p_i \right), \mathcal{O}_C \right) = \begin{cases} 0, & i = 0, 2; \\ 3g - 3 + n, & i = 1. \end{cases}$$

Proof. We let k is algebraically closed and has no marked point. Let $\pi : \tilde{C} \rightarrow C$ be a normalization and let $\Sigma \subset C$ be the set of nodes. Let $\tilde{\Sigma} = \pi^{-1}(\Sigma) \subset \tilde{C}$.

By Lemma 8.2.1, we get $\dim_k \operatorname{Ext}_{\mathcal{O}_C}^2(\Omega_C, \mathcal{O}_C) = 0$. For Ext^0 , we first claim that

$$\operatorname{Hom}_{\mathcal{O}_{\tilde{C}}}(\Omega_{\tilde{C}}(\tilde{\Sigma}), \mathcal{O}_{\tilde{C}}) \cong \operatorname{Hom}_{\mathcal{O}_C}(\Omega_C, \mathcal{O}_C).$$

This is equivalent to show

$$\operatorname{Der}_k(\mathcal{O}_C, \mathcal{O}_C) \cong \{D' \in \operatorname{Der}_k(\mathcal{O}_{\tilde{C}}, \mathcal{O}_{\tilde{C}}) : D' \text{ fixes every points in } \tilde{\Sigma}\}.$$

Actually this is just Lemma 8.2.8. This finish the claim. Hence we get

$$\operatorname{Hom}_{\mathcal{O}_C}(\Omega_C, \mathcal{O}_C) \cong \operatorname{Hom}_{\mathcal{O}_{\tilde{C}}}(\Omega_{\tilde{C}}(\tilde{\Sigma}), \mathcal{O}_{\tilde{C}}) \cong H^0(\tilde{C}, T_{\tilde{C}}(-\tilde{\Sigma})) = 0.$$

For Ext^1 , by Lemma 8.2.1 we have

$$\begin{array}{ccc} 0 \rightarrow H^1(C, \mathcal{H}om_{\mathcal{O}_C}(\Omega_C, \mathcal{O}_C)) \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C, \mathcal{O}_C) \rightarrow \bigoplus_j \text{Ext}_{\widehat{\mathcal{O}}_{C, q_j}}^1(\Omega_{\widehat{\mathcal{O}}_{C, q_j}}, \widehat{\mathcal{O}}_{C, q_j}) \\ \parallel \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \downarrow \\ H^1(\widetilde{C}, T_{\widetilde{C}}(-\widetilde{\Sigma})) \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad 0 \end{array}$$

and $\text{Ext}_{\widehat{\mathcal{O}}_{C, q_j}}^1(\Omega_{\widehat{\mathcal{O}}_{C, q_j}}, \widehat{\mathcal{O}}_{C, q_j}) = k$. This equality in this exact sequence is because

$$H^1(C, \mathcal{H}om_{\mathcal{O}_C}(\Omega_C, \mathcal{O}_C))$$

be the set of locally trivial first order deformation of C preserving nodes and this is equivalent to the set of locally trivial first order deformation of \widetilde{C} fixed $\widetilde{\Sigma}$, which is $H^1(\widetilde{C}, T_{\widetilde{C}}(-\widetilde{\Sigma}))$.

Now let $\widetilde{C} = \coprod_{i=1}^t \widetilde{C}_i$ are connected components and $\widetilde{\Sigma}_i = \widetilde{C}_i \cap \widetilde{\Sigma}$. First we have

$$h^1(\widetilde{C}_i, T_{\widetilde{C}_i}(-\widetilde{\Sigma}_i)) = h^0(\widetilde{C}_i, \Omega_{\widetilde{C}_i}^{\otimes 2}(\widetilde{\Sigma}_i)) = 3g(\widetilde{C}_i) - 3 + \#(\widetilde{\Sigma}_i),$$

hence

$$\begin{aligned} \dim_k \text{Ext}_{\mathcal{O}_C}^1(\Omega_C, \mathcal{O}_C) &= h^1(\widetilde{C}, T_{\widetilde{C}}(-\widetilde{\Sigma})) + \#(\Sigma) \\ &= \sum_{i=1}^t (3g(\widetilde{C}_i) - 3 + \#(\widetilde{\Sigma}_i)) + \#(\Sigma) = 3 \sum_{i=1}^t g(\widetilde{C}_i) - 3t + 3\#(\Sigma) = 3g - 3 \end{aligned}$$

by the genus formula, we win. \square

8.3 Basic concept of Kuranishi family

We will work on analytic category over \mathbb{C} via Serre's GAGA.

Definition 8.3.1. Let $(C; p_1, \dots, p_n)$ be an n -pointed connected nodal curve of genus g over \mathbb{C} . A deformation

$$\phi : \mathcal{C} \rightarrow (B, b_0), \sigma_i : B \rightarrow \mathcal{C} \text{ such that } \chi : (C; p_1, \dots, p_n) \cong (\phi^{-1}(b_0), \sigma_i(b_0))$$

of $(C; p_1, \dots, p_n)$ is said to be a Kuranishi family for $(C; p_1, \dots, p_n)$ if it satisfies the following condition:

(Condition K). For any deformation $\psi : \mathcal{D} \rightarrow (E, e_0)$ of $(C; p_1, \dots, p_n)$ and for any sufficiently small connected neighborhood U of e_0 , there is a unique morphism of deformations of n -pointed curves

$$\begin{array}{ccc} \mathcal{D}|_U & \xrightarrow{F} & \mathcal{C} \\ \downarrow \psi|_U & & \downarrow \phi \\ (U, e_0) & \xrightarrow{f} & (B, b_0) \end{array}$$

We will show the following theorem at next two sections:

Theorem 8.3.2. A Kuranishi family for $(C; p_1, \dots, p_n)$ exists if and only if $(C; p_1, \dots, p_n)$ is stable.

Corollary 8.3.3. *The Kodaira-Spencer map of a Kuranishi family at the base point is an isomorphism.*

Proof. This is trivial as the family $\mathcal{C}_\varepsilon \rightarrow \text{Spec}\mathbb{C}[\varepsilon]$ just has and has unique map to $\phi : \mathcal{C} \rightarrow (B, b_0)$. This defines a bijection between T_{B, b_0} and $\text{Def}(C) = \text{Ext}_{\mathcal{O}_C}^1(\Omega_C(p_1 + \dots + p_n), \mathcal{O}_C)$ via κ_{B, b_0} . \square

Corollary 8.3.4. *Let there be given a deformation of a stable n -pointed curve $(C; p_1, \dots, p_n)$ over the pointed analytic space (E, e_0) . Suppose that its Kodaira-Spencer map at e_0 is an isomorphism and that E is smooth at e_0 . Then the deformation is a Kuranishi family for $(C; p_1, \dots, p_n)$.*

Proof. To add. \square

Corollary 8.3.5. *The base of the Kuranishi family of a stable n -pointed curve $(C; p_1, \dots, p_n)$ of genus g is smooth of dimension $3g - 3 + n$.*

Proof. By the previous corollary and Theorem 8.2.6 and the uniqueness of the Kuranishi family by the universal property. \square

Corollary 8.3.6. *Let $X \rightarrow S$ be a family of stable n -pointed curves, and s_0 a point of S . If $X \rightarrow S$ is a Kuranishi family for X_{s_0} , then it is a Kuranishi family for X_s , for all s in an open neighborhood U of s_0 .*

Proof. From the previous results that $X \rightarrow S$ is Kuranishi for X_s if and only if s is a smooth point of S and the Kodaira-Spencer map at s is an isomorphism. The first of these conditions is clearly open.

Since the dimension of $\text{Ext}^1(\Omega_{X_s}^1, \mathcal{O}_{X_s}(-\sum \sigma_i(s)))$ is independent of s , the second condition translates into a rank condition for a map between vector bundles and hence is open and We win. \square

8.4 The Hilbert scheme of ν -canonical curves

8.5 Construction of Kuranishi families


Chapter 9

The stack of all curves

9.1 Families of all arbitrary curves

Definition 9.1.1. Here we redefine a curve over k is a scheme C of finite type over k of dimension 1 (rather than pure dimension 1). The genus of C is defined as $g(C) = 1 - \chi(C, \mathcal{O}_C)$.

Remark 9.1.2. Why we not allow pure dimension 1? Since they may arise as deformations of connected pure one-dimensional curves; without this relaxation, the stack of all curves would fail to be algebraic. For example in [23] Example III.9.8.4, a flat family of rational curves defined by $\mathbb{P}^1 \rightarrow \mathbb{P}^3$ via $[x : y] \mapsto [x^3 : x^2y : xy^2 : ty^3]$ for any $t \neq 0$. As $t \rightarrow 0$, we may get a singular non-reduced curve C_0 with an embedded point at that node, but C_0 can deform to the disjoint union of a plane nodal curve and a point in \mathbb{P}^3 .



Definition 9.1.3. (i) A family of curves over a scheme S is a flat, proper and finitely presented morphism $C \rightarrow S$ of algebraic spaces such that every fiber is a curve.

(ii) A family of n -pointed curves is a family of curves $C \rightarrow S$ together with n sections $\sigma_1, \dots, \sigma_n : S \rightarrow C$ (with no condition on whether they are distinct or land in the relative smooth locus of C over S).

Remark 9.1.4. (i) When we consider a family of stable curve, since the relative dualizing sheaf is ample, we can get it is projective, hence must be a scheme;

(ii) There are some examples such that C are not a scheme.

Proposition 9.1.5. If $C \rightarrow S$ is a family of curves over S , there exists an étale cover $S' \rightarrow S$ such that $C_{S'} \rightarrow S'$ is projective.

Sketch-Local to global. Consider cartesian

$$\begin{array}{ccccccc}
 C_0 := C_s & \hookrightarrow & C_1 & \hookrightarrow & \cdots & \hookrightarrow & \widehat{C} \hookrightarrow C \\
 \downarrow & & \downarrow & & & & \downarrow \\
 S_0 := \mathrm{Spec} \kappa(s) & \hookrightarrow & S_1 := \mathrm{Spec} \mathcal{O}_{S,s}/\mathfrak{m}_s^2 & \hookrightarrow & \cdots & \hookrightarrow & \widehat{S} := \mathrm{Spec} \widehat{\mathcal{O}}_{S,s} \longrightarrow S
 \end{array}$$

Step 1. $C_0 \rightarrow \mathrm{Spec} \kappa(s)$. By St 0ADD, every separated algebraic space of dimension one is a scheme. Any one-dimensional proper $\kappa(s)$ -scheme is projective by St 0A26. In particular we get a ample line bundle L_0 on C_0 .

Step 2. $C_n \rightarrow S_n$. The obstruction to deforming a line bundle L_n on C_n to L_{n+1} on C_{n+1} lives in $H^2(C_0, \mathcal{O}_{C_0})$ and thus vanishes as $\dim C_0 = 1$. Thus there exists a compatible sequence of line bundles L_n on C_n . Since ampleness is an open condition in families and L_0 is ample, L_n is also ample.

Step 3. $\widehat{C} \rightarrow \widehat{S}$ with \widehat{S} noetherian. Use Grothendieck's Existence Theorem we get an equivalence $\mathrm{Coh}(\widehat{C}) \rightarrow \varprojlim \mathrm{Coh}(C_n)$. As $\widehat{C} \rightarrow \widehat{S}$ is proper, then by Chow's lemma there exists a projective birational morphism $C' \rightarrow \widehat{C}$ of algebraic spaces such that $C' \rightarrow S$ is projective. This allows one to reduce Grothendieck's Existence Theorem for $\widehat{C} \rightarrow \widehat{S}$ to $C' \rightarrow \widehat{S}$ using devissage. As a result, using again that ampleness is an open condition in families we can extend the sequence of line bundle L_n to a line bundle \widehat{L} on \widehat{C} which is ample.

Step 4. S is of finite type over \mathbb{Z} . For every closed point $s \in S$, apply Artin Approximation to the functor

$$(Sch/S) \rightarrow (Sets), (T \rightarrow S) \mapsto \mathrm{Pic}(C_T)$$

to obtain an étale neighborhood $(S', s') \rightarrow (S, s)$ of s and a line bundle L' on $C_{S'}$ extending L_0 . By openness of ampleness, we can replace S' with an open neighborhood of s' such that L' is relatively ample over S' .

Step 5. General S . Use noetherian approximation. □

9.2 Algebraicity of the stack of all curves

Definition 9.2.1. Let $\mathcal{M}_{g,n}^{\mathrm{all}}$ denote the category over Sch_{et} whose objects over S consists of families of curves $C \rightarrow S$ and n sections $\sigma_i : S \rightarrow C$. A morphism $(C' \rightarrow S', \sigma'_i) \rightarrow (C \rightarrow S, \sigma_i)$ is the data of the cartesian

$$\begin{array}{ccc}
 C' & \xrightarrow{g} & C \\
 \sigma'_i \uparrow \downarrow & & \sigma_i \uparrow \downarrow \\
 S' & \xrightarrow{f} & S
 \end{array}$$

with $g \circ \sigma'_i \rightarrow \sigma_i \circ f$.

Lemma 9.2.2. $\mathcal{M}_{g,n}^{\mathrm{all}}$ is a stack over Sch_{et} .

Proof. Handle $n = 0$. Let $S' \rightarrow S$ be an étale cover with $C' \rightarrow S'$. And $\alpha : p_1^* C' \rightarrow p_2^* C'$ is an isomorphism over $S' \times_S S'$ satisfying the cocycle condition. The quotient of the étale equivalence relation

$$\begin{array}{ccccc}
 R := p_1^* C' & \xrightarrow[p_2 \circ \alpha]{p_1} & C' & \dashrightarrow & C := C'/R \\
 \downarrow & & \downarrow & & \downarrow \\
 S' \times_S S' & \xrightarrow[p_2]{p_1} & S' & \longrightarrow & S
 \end{array}$$

Well done. \square

Lemma 9.2.3. $\Delta : \mathcal{M}_{g,n}^{\text{all}} \rightarrow \mathcal{M}_{g,n}^{\text{all}} \times \mathcal{M}_{g,n}^{\text{all}}$ is representable.

Proof. Handle $n = 0$. Consider the cartesian

$$\begin{array}{ccc} \text{Isom}_T(C_1, C_2) & \longrightarrow & T \\ \downarrow & & \downarrow (C_1, C_2) \\ \mathcal{M}_{g,n}^{\text{all}} & \xrightarrow{\Delta} & \mathcal{M}_{g,n}^{\text{all}} \times \mathcal{M}_{g,n}^{\text{all}} \end{array}$$

We need to show $\text{Isom}_T(C_1, C_2)$ is an algebraic space. By Proposition 9.1.5, there exists an étale cover $T' \rightarrow T$ such that $C_{i,T'} \rightarrow T'$ is projective. Hence we may let C_1, C_2 are projective over T . Indeed, as

$$\text{Isom}_T(C_1, C_2) \times_T T' = \text{Isom}_{T'}(C_{1,T'}, C_{2,T'}),$$

we get $\text{Isom}_{T'}(C_{1,T'}, C_{2,T'}) \rightarrow \text{Isom}_T(C_1, C_2)$ is representable, surjective and étale. Hence if $\text{Isom}_{T'}(C_{1,T'}, C_{2,T'})$ is an algebraic space, so is $\text{Isom}_T(C_1, C_2)$.

Fact. (St 05XD) If $f : X \rightarrow Y$ are T -morphism such that X, Y are proper, flat and locally of finite presentation over T , then for any $U \rightarrow T$ such that $X_U \cong Y_U$ if and only if $U \rightarrow T$ factor through an open subscheme $T_0 \subset T$.

Now we get the inclusions

$$\text{Isom}_T(C_1, C_2) \subset \text{Mor}_T(C_1, C_2) \subset \text{Hilb}(C_1 \times_T C_2 / T)$$

where the second inclusion is $(g : C_1 \rightarrow C_2) \mapsto (\Gamma_g : C_1 \rightarrow C_1 \times_T C_2)$. The first inclusion is representable open immersion by the above fact. The second inclusion, we find that a subspace $[Z \subset C_1 \times_T C_2] \in \text{Hilb}(C_1 \times_T C_2 / T)$ is in the image of the inclusion if and only if $Z \rightarrow C_1 \times_T C_2 \rightarrow C_1$ is an isomorphism (and similarly for other valued points). Therefore by the above fact we win. \square

Theorem 9.2.4. $\mathcal{M}_{g,n}^{\text{all}}$ is an algebraic stack locally of finite type over \mathbb{Z} .

Sketch. Step 1. Reduce to $n = 0$. Since $\mathcal{M}_{g,n+1}^{\text{all}}$ is the universal family over $\mathcal{M}_{g,n}^{\text{all}}$, we can prove the conclusion at the case $\mathcal{M}_g^{\text{all}}$. (Why?)

Step 2. Look for possible smooth cover H' of $\mathcal{M}_g^{\text{all}}$. Let C_0 be any projective curves C_0 over k . Choosing an embedding $C_0 \subset \mathbb{P}_k^N$ such that $h^1(C_0, \mathcal{O}(1)) = 0$ by Serre's vanishing theorem. Let $P(t)$ be its Hilbert polynomial. Let $H := \text{Hilb}^P(\mathbb{P}_{\mathbb{Z}}^N / \mathbb{Z})$ be the Hilbert scheme which is projective over \mathbb{Z} . Consider the universal family

$$\begin{array}{ccc} \mathcal{C} & \hookrightarrow & \mathbb{P}_H^N \\ \downarrow & \swarrow & \\ H & & \end{array}$$

there is a point $h_0 \in H(k)$ such that $\mathcal{C}_{h_0} = C_0$. By Review 1 we can find an open neighborhood $H' \subset H$ of h_0 such that for any $s \in H'$ we have $h^1(\mathcal{C}_s, \mathcal{O}_{\mathcal{C}_s}(1)) = 0$. Now consider

$$H' \rightarrow \mathcal{M}_g^{\text{all}}, [C \hookrightarrow \mathbb{P}^N] \mapsto [C],$$

and by the representability of the diagonal, this map is representable as H' is a scheme.

Step 3. Show that $H' \rightarrow \mathcal{M}_g^{all}$ is smooth. Using Infinitesimal Lifting Criterion such that for all surjections $A \rightarrow A_0$ of artinian local rings with residue field k such that $k = \ker(A \rightarrow A_0)$ and for all diagrams

$$\begin{array}{ccc}
 & \text{Spec } k & \\
 \swarrow & & \searrow \\
 \text{Spec } A_0 & \xrightarrow{[\mathcal{C}_0 \subset \mathbb{P}_{A_0}^N]} & H' \\
 \downarrow & \nearrow [\mathcal{C} \subset \mathbb{P}_A^N] & \downarrow \\
 \text{Spec } A & \xrightarrow{\mathcal{C}} & \mathcal{M}_g^{all}
 \end{array}$$

(Note: The diagram also includes a solid arrow from $\text{Spec } k$ to $\text{Spec } A_0$ labeled $[C \subset \mathbb{P}_k^N]$ and a solid arrow from $\text{Spec } k$ to H' labeled $[C \subset \mathbb{P}_k^N]$.)

We need to find that dotted arrow. This diagram is equivalent to

$$\begin{array}{ccccc}
 & & \mathbb{P}_k^N & & \mathbb{P}_{A_0}^N & & \mathbb{P}_A^N \\
 & \nearrow & & \nearrow & & \nearrow & \\
 C & \xrightarrow{\quad} & \mathcal{C}_0 & \xrightarrow{\quad} & C & \xrightarrow{\quad} & C \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 \text{Spec } k & \hookrightarrow & \text{Spec } A_0 & \hookrightarrow & \text{Spec } A & &
 \end{array}$$

For simplifying, we let C is of locally complete intersection (general case see [19] and [18]). Let \mathcal{J} be the ideal sheaf of $C \rightarrow \mathbb{P}_k^N$ generated by regular sequence locally and that $\mathcal{J}/\mathcal{J}^2$ is a vector bundle on C with

$$0 \rightarrow \mathcal{J}/\mathcal{J}^2 \rightarrow \Omega_{\mathbb{P}_k^N}|_C \rightarrow \Omega_C \rightarrow 0.$$

By long exact sequence we get

$$\text{Hom}_{\mathcal{O}_C}(\mathcal{J}/\mathcal{J}^2, \mathcal{O}_C) \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C, \mathcal{O}_C) \rightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_{\mathbb{P}_k^N}|_C, \mathcal{O}_C) = H^1(C, T_{\mathbb{P}_k^N}|_C).$$

Consider the canonical sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow \mathcal{O}_C(1)^{\oplus N+1} \rightarrow T_{\mathbb{P}^N}|_C \rightarrow 0.$$

Since $H^2(C, \mathcal{O}_C) = 0$ and $H^1(C, \mathcal{O}_C(1)) = 0$ by $[C] \in H'$ we get $H^1(C, T_{\mathbb{P}_k^N}|_C) = 0$. Hence we get a surjection

$$\text{Hom}_{\mathcal{O}_C}(\mathcal{J}/\mathcal{J}^2, \mathcal{O}_C) \twoheadrightarrow \text{Ext}_{\mathcal{O}_C}^1(\Omega_C, \mathcal{O}_C).$$

Use some deformation theory (But I don't know! May be use something about cotangent complex which is the reason we let C is of locally complete intersection!) we get $\text{Hom}_{\mathcal{O}_C}(\mathcal{J}/\mathcal{J}^2, \mathcal{O}_C)$ classifies embedded deformations of $\mathcal{C}_0 \rightarrow \mathbb{P}_{A_0}^N$ to $C' \rightarrow \mathbb{P}_A^N$ and $\text{Ext}_{\mathcal{O}_C}^1(\Omega_C, \mathcal{O}_C)$ classifies deformations of \mathcal{C}_0 over A_0 to C' over A . As the map is $[\mathcal{C}' \rightarrow \mathbb{P}_A^N] \mapsto C'$ and is surjective, we win. \square

9.3 Algebraicity of several stacks and boundedness of stable curves

Proposition 9.3.1 (Several stacks). *We have inclusions*

$$\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n} \subset \mathcal{M}_{g,n}^{ss} \subset \mathcal{M}_{g,n}^{\text{pre}} \subset \mathcal{M}_{g,n}^{\leq \text{nodal}} \subset \mathcal{M}_{g,n}^{all}$$

of prestacks. Then all of these are open substacks, hence all of these are algebraic stacks locally of finite type over \mathbb{Z} .

Proof. • By Theorem 9.2.4, $\mathcal{M}_{g,n}^{all}$ is an algebraic stack locally of finite type over \mathbb{Z} .

• $\mathcal{M}_{g,n}^{\leq nodal} \subset \mathcal{M}_{g,n}^{all}$ is an open substack. Actually by Corollary 6.4.3 we get the nodal locus is open when C is a scheme. In general for an étale cover $g : C' \rightarrow C$ by a scheme, we find that a point $p \in C'$ is a node in its fiber if and only if $g(p)$ is a node in its fiber. We win.

• $\mathcal{M}_{g,n}^{pre} \subset \mathcal{M}_{g,n}^{\leq nodal}$ is an open substack. This is because for a family $(C \rightarrow S, \{\sigma_i\})$ of nodal curves, the locus $\{s \in S : \sigma_i(s) \text{ are disjoint and smooth}\}$ is open.

• $\mathcal{M}_{g,n}^{ss} \subset \mathcal{M}_{g,n}^{pre}$ is an open substack. This is because the nef locus is open (**But I don't know the relation between nefness and semistable!**).

• $\overline{\mathcal{M}}_{g,n} \subset \mathcal{M}_{g,n}^{ss}$ is an open substack. Indeed the stable locus is open by Proposition 7.3.4.

• $\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$ is an open substack. Indeed this is by the fact that smooth locus is open. \square

Proposition 9.3.2. $\overline{\mathcal{M}}_{g,n}$ is a quasi-compact smooth Deligne-Mumford stack of dimension $3g - 3 + n$ over \mathbb{Z} .

Proof. • $\overline{\mathcal{M}}_{g,n}$ is quasi-compact. Let (C, p_1, \dots, p_n) be a n -pointed stable curve. By Theorem 7.2.1, we get $(\omega_C(p_1 + \dots + p_n))^{\otimes 3}$ is very ample, we get $C \hookrightarrow \mathbb{P}^N$ with Hilbert polynomial $P(t)$. This is independent of C . Hence consider closed subscheme $H \subset \underline{\text{Hilb}}^P(\mathbb{P}_{\mathbb{Z}}^N/\mathbb{Z}) \times (\mathbb{P}^N)^n$ of embedded curve and n points $(C \hookrightarrow \mathbb{P}^N, p_i \in C)$. Consider a forgetful functor

$$H \rightarrow \mathcal{M}_{g,n}^{all}, (C \hookrightarrow \mathbb{P}^N, p_i \in C) \mapsto (C, \{p_i\}).$$

Then the image of $|H| \rightarrow |\mathcal{M}_{g,n}^{all}|$ contains $\overline{\mathcal{M}}_{g,n}$. As $\underline{\text{Hilb}}^P(\mathbb{P}_{\mathbb{Z}}^N/\mathbb{Z})$ is projective, then H is quasi-compact. Hence $\overline{\mathcal{M}}_{g,n}$ is quasi-compact.

• $\mathcal{M}_{g,n}$ is Deligne-Mumford stack. By Proposition 8.2.9 for $i = 0$ and Proposition 8.2.7 (a), we get a n -pointed stable curve (C, p_1, \dots, p_n) has no infinitesimal automorphisms, i.e. that the Lie algebra $T_e \text{Aut}(C, p_1, \dots, p_n)$ is trivial. Since the automorphism group scheme $\text{Aut}(C, p_1, \dots, p_n)$ is of finite type, this implies that $\text{Aut}(C, p_1, \dots, p_n)$ is finite and discrete, hence $\overline{\mathcal{M}}_{g,n}$ is a quasi-compact Deligne-Mumford stack.

• $\mathcal{M}_{g,n}$ is smooth over $\text{Spec} \mathbb{Z}$. Proposition 8.2.9 for $i = 2$ and Proposition 8.2.7 (c) implies that there are no obstructions to deforming C . As the algebraicity of $\overline{\mathcal{M}}_{g,n}$, this will allow us to invoke the Infinitesimal Lifting Criterion to establish that $\overline{\mathcal{M}}_{g,n}$ is smooth over $\text{Spec} \mathbb{Z}$.

• $\overline{\mathcal{M}}_{g,n}$ has relative dimension $3g - 3 + n$ over $\text{Spec} \mathbb{Z}$. Proposition 8.2.9 for $i = 1$ and Proposition 8.2.7 (b) implies that isomorphism classes of deformations of (C, p_1, \dots, p_n) , it is identified with the Zariski tangent space of $\overline{\mathcal{M}}_{g,n}$ at the point corresponding to (C, p_1, \dots, p_n) . This will allow us to conclude that $\overline{\mathcal{M}}_{g,n}$ has relative dimension $3g - 3 + n$ over \mathbb{Z} . \square

9.4 The family of elliptic curves $\mathcal{M}_{1,1}$

Proposition 9.4.1. \mathcal{M}_1 is not a stack.

Proof. See [32] Remark 8.4.15 for the References. \square

Remark 9.4.2. If we let \mathcal{M}'_1 as morphisms of algebraic spaces, then this will be a stack. This follows the Picard functor and the stack $\mathcal{M}_{1,1}$. In fact if we consider the universal elliptic curve $\mathcal{E} \rightarrow \mathcal{M}_{1,1}$, then Picard functor gives $\mathcal{M}'_1 \rightarrow \mathcal{M}_{1,1}$ which induce $\mathcal{M}'_1 \cong B\mathcal{E}$. We omitted here.

Proposition 9.4.3. $\mathcal{M}_{1,1}$ is a smooth Deligne-Mumford stack.

Proof. By Proposition 9.3.1 we get that $\mathcal{M}_{1,1}$ is an open substack of $\overline{\mathcal{M}}_{1,1}$. Hence it is a smooth Deligne-Mumford stack. \square

Proposition 9.4.4. $\mathcal{M}_{1,1}$ has a coarse moduli space $M_{1,1} \cong \mathbb{A}^1$.

Proof.

□

Chapter 10

Stable reduction: why $\overline{\mathcal{M}}_{g,n}$ is proper?

In this section we will use the Valuative Criterion (Theorem C.1.3 (1)) to show that $\overline{\mathcal{M}}_{g,n}$ is proper. The existence of extension is called stable reduction, which is our main theorem:

Lemma 10.0.1. *The diagonal of the stack $\mathcal{M}_{g,n}^{all}$ is separated. In particular, $\mathcal{M}_{g,n}^{all} \rightarrow \text{Spec}(\mathbb{Z})$ is quasi-separated.*

Proof. Omitted. See St 0DSQ. □

Theorem 10.0.2 (Stable reduction). *Let R be a DVR with fraction field K and $\Delta = \text{Spec}(R)$, $\Delta^* = \text{Spec}(K)$. If $(\mathcal{C}^* \rightarrow \Delta^*, s_1^*, \dots, s_n^*)$ is a family of n -pointed stable curves of genus g , then there exists a finite cover $\Delta' \rightarrow \Delta$ of spectrums of DVRs and a family $(\mathcal{C}' \rightarrow \Delta', s_1', \dots, s_n')$ of stable curves extending $\mathcal{C}^* \times_{\Delta^*} \Delta'^* \rightarrow \Delta'^*$. As*

$$\begin{array}{ccccc}
 & & \mathcal{C}^* & & \\
 & \nearrow & \downarrow & \dashrightarrow & \mathcal{C}' \\
 \mathcal{C}^* \times_{\Delta^*} \Delta'^* & \xrightarrow{\quad} & \Delta'^* & \xrightarrow{\quad} & \Delta^* \\
 \downarrow & \searrow & \downarrow & \searrow & \downarrow \\
 \Delta'^* & \xrightarrow{\quad} & \Delta^* & \xrightarrow{\quad} & \Delta
 \end{array}$$

given by

$$\begin{array}{ccc}
 \Delta'^* & \xrightarrow{\quad} & \Delta^* \\
 \downarrow & & \downarrow \\
 \Delta' & \xrightarrow{\quad} & \Delta
 \end{array}
 \begin{array}{c}
 \xrightarrow{(C^* \rightarrow \Delta^*, \{s_i^*\})} \overline{\mathcal{M}}_{g,n} \\
 \xrightarrow{(C' \rightarrow \Delta', \{s_1'\})} \overline{\mathcal{M}}_{g,n}
 \end{array}$$


After proving this and the uniqueness, we can get the following conclusion:

Theorem 10.0.3. *If $2g - 2 + n > 0$, then $\overline{\mathcal{M}}_{g,n}$ is a proper smooth Deligne-Mumford stack of dimension $3g - 3 + n$ over \mathbb{Z} .*

By using the Keel-Mori Theorem, we get

Corollary 10.0.4. *If $2g - 2 + n > 0$, there exists a coarse moduli space $\overline{\mathcal{M}}_{g,n} \rightarrow \overline{M}_{g,n}$ where $\overline{M}_{g,n}$ is a proper algebraic space over \mathbb{Z} .*

Example 1. *Let $\Delta = \text{Spec}(R)$ where R be a DVR with uniformizer t . Let C be a smooth curve and consider $\mathcal{C} = C \times \Delta$ with sections $(\sigma_1, \sigma_2, \sigma_3) = (t^2, -t^2, 4t)$ as following diagram. The first*



two arrows are blowing up and the third is contracting E_1 .

Remark 10.0.5. *Actually there are several methods to prove this. The first proof due to the original paper [9] by consider the Jacobians of curves and reduce the case into the semistable reduction of abelian varieties. Our method follows [20] by using some birational geometry of surfaces to prove the case of characteristic 0. There is another method can deal the positive or mixed characteristic case by [4], for this we refer St 0E8C.*

10.1 Proof of stable reduction in characteristic 0

Lemma 10.1.1. *Let R be a DVR with uniforming t and $0 := (t)$. Let $\mathcal{C} \rightarrow \Delta = \text{Spec}(R)$ be a flat, proper and finitely presented morphisms such that each geometric fiber is a curve. Assume that \mathcal{C} is regular. Let $p \in \mathcal{C}_0$.*

(a) *If p is a smooth point in the reduced fiber $(\mathcal{C}_0)_{\text{red}}$. Show that after possibly an extension of DVRs, there exists an étale neighborhood of p (defined over R)*

$$\text{Spec}R[x, y]/(x^a - t) \rightarrow \mathcal{C}.$$

(b) *If p is a node in the reduced fiber $(\mathcal{C}_0)_{\text{red}}$. Show that there exists an étale neighborhood of p (defined over R)*

$$\text{Spec}R[x, y]/(x^a y^b - t) \rightarrow \mathcal{C}.$$

Proof.

□

Lemma 10.1.2. *Let a, b, m be positive integers such that both a and b divide m .*


(a) *Let $X = \text{Spec}k[x, t]/(t^m - x^a)$ and normalization $\tilde{X} \rightarrow X$. Then each preimage of the origin is locally defined by $x = t^k$ for some k .*

(b) *Let $X = \text{Spec}k[x, y, t]/(t^m - x^a y^b)$ and normalization $\tilde{X} \rightarrow X$. Then each preimage of the origin is locally defined by $t^k = xy$. In particular is a reduced and nodal point in the fiber over $t = 0$.*

Proof. (a) We have $x^a - t^m = \prod_{i=0}^{a-1} (x - \zeta^i t^{m/a})$ where ζ be a primitive a -th root of unity. Hence the origin has a preimages in \tilde{X} locally defined by $x - \zeta^i t^{m/a}$, respectively.

(b)

□



Lemma 10.1.3. *Let $\mathcal{C} \rightarrow \Delta = \text{Spec}(R)$ be a family of nodal curves where R be a DVR such that the general fiber \mathcal{C}^* is smooth. Then if E is a rational tail (rational bridge with out marked points) of \mathcal{C}_0 , then $E^2 = -1$ ($E^2 = -2$). As*

Proof. For any $E \cong \mathbb{P}^1 \subset \mathcal{C}_0$, then $0 = E \cdot \mathcal{C}_0 = E^2 + E \cdot E^c$. We win. (Actually I'm not familiar with surface and the intersection numbers!) \square

For simplicity of notation, we assume that there are no marked points, i.e. $n = 0$. Fix a spectrum of DVR $\Delta = \text{Spec}(R)$, $\Delta^* = \text{Spec}(K)$ and $t \in R$ is the uniformizer, and $0 = (t) \in \text{Spec} R$ the unique closed point. Consider $\mathcal{C}^* \rightarrow \Delta^*$ be a family of stable curve.

STEP 1. Reduce to the case where $\mathcal{C}^* \rightarrow \Delta^*$ is smooth. If \mathcal{C}^* has k nodes, then after a finite extension of K we can arrange that each node is given by K -points $p_i \in \mathcal{C}^*(K)$. Let the pointed normalization $(\tilde{\mathcal{C}}^*, q_1, \dots, q_{2k})$ of it. By induction on the genus g , we perform stable reduction on each connected component and then take the nodal union along sections. After possibly an extension of K (and R), this produces a family of curves $\mathcal{C} \rightarrow \Delta$ extending $\mathcal{C}^* \rightarrow \Delta^*$.

STEP 2. Find some flat extension $\mathcal{C} \rightarrow \Delta$. As $\omega_{\mathcal{C}^*/\Delta^*}^{\otimes 3}$ is very ample, we can get an embedding as follows

$$\begin{array}{ccc}
 \mathbb{P}^{5g-6} \times \Delta^* & \hookrightarrow & \mathbb{P}^{5g-6} \times \Delta \\
 \uparrow |\omega_{\mathcal{C}^*/\Delta^*}^{\otimes 3}| & & \uparrow \\
 \mathcal{C}^* & \hookrightarrow & \mathcal{C} := \overline{\mathcal{C}^*} \\
 \downarrow & & \downarrow f \\
 \Delta^* & \hookrightarrow & \Delta
 \end{array}$$

where $\mathcal{C} := \overline{\mathcal{C}}$ be the scheme-theoretic image of $\mathcal{C}^* \hookrightarrow \mathbb{P}^{5g-6} \times \Delta$. Now we focus on f . Actually the scheme-theoretic closure does not bring more embedded points. Hence by Proposition 2.1.2 we get f is flat.

STEP 3. Use embedded resolutions to find a resolution of singularities $\tilde{\mathcal{C}} \rightarrow \mathcal{C}$ so that the reduced central fiber $(\tilde{\mathcal{C}}_0)_{red}$ is nodal. By Theorem B.0.2, there exists a finite sequence of blow-ups at closed points of \mathcal{C}_0 yielding a projective birational morphism

$$\begin{array}{c}
 \tilde{\mathcal{C}}_0 \subset \tilde{\mathcal{C}} \longrightarrow \cdots \longrightarrow \mathcal{C} \supset \mathcal{C}_0 \\
 \searrow \hspace{10em} \downarrow \\
 \hspace{10em} \Delta
 \end{array}$$

such that $\tilde{\mathcal{C}}$ is regular flat family of curves and such that the reduced central fiber $(\tilde{\mathcal{C}}_0)_{red}$ is nodal. Now replace \mathcal{C} by $\tilde{\mathcal{C}}$.

STEP 4. Perform a base change $\Delta' \rightarrow \Delta$ such that the normalization of the total family $\mathcal{C} \times_{\Delta} \Delta'$ has a reduced nodal central fiber with many rational tails and bridges.

By Lemma 10.1.1, we choose local coordinates x, y around $p \in \mathcal{C}_0$ (étale locally and formally locally) such that $\mathcal{C} \rightarrow \Delta$ can be described as follows:

- (i) If $p \in (\mathcal{C}_0)_{red}$ is a smooth point, then $(x, y) \mapsto x^a$ and the multiplicity of the irreducible component of \mathcal{C}_0 containing p is a ;
- (ii) If $p \in (\mathcal{C}_0)_{red}$ is a (separated) node, then $(x, y) \mapsto x^a y^b$ and the two components of \mathcal{C}_0 containing p have multiplicities a and b .

Let m be the least common multiple of the multiplicities of the irreducible components of \mathcal{C}_0 . Let the ramified morphism $\Delta' = \text{Spec}(R) \rightarrow \Delta$ given by $t \mapsto t^m$. Hence we get

$$\begin{array}{ccccc} \tilde{\mathcal{C}}' & \longrightarrow & \mathcal{C}' & \longrightarrow & \mathcal{C} \\ & & \downarrow & & \downarrow \\ & & \Delta' & \longrightarrow & \Delta \end{array}$$

where $\mathcal{C}' = \mathcal{C} \times_{\Delta} \Delta'$ and $\tilde{\mathcal{C}}' \rightarrow \mathcal{C}'$ be the normalization. Consider $p \in (\mathcal{C}_0)_{red}$.

(a) If p is a smooth point, then the unique preimage of p in \mathcal{C}' defined locally by $x^a - t^m$. By Lemma 10.1.2 (a), we get each preimage of p in $\tilde{\mathcal{C}}'$ is locally defined by $x = t^k$ which are the smooth points in $\tilde{\mathcal{C}}'_0$;

(b) If p is a node, then the unique preimage of p in \mathcal{C}' defined locally by $x^a y^b - t^m$. By Lemma 10.1.2 (a), we get each preimage of p in $\tilde{\mathcal{C}}'$ is locally defined by $xy = t^k$ which are reduced and nodal points in $\tilde{\mathcal{C}}'_0$. If $k > 1$, $\tilde{\mathcal{C}}'$ have A_{k-1} -singularity.

Hence now we replace \mathcal{C} by $\tilde{\mathcal{C}}'$, which has a reduced central fiber with many rational tails and bridges.

STEP 5. After taking the minimal model, contract all rational tails and bridges in the central fiber. Using Theorem B.0.1 we get a minimal resolution $\mathcal{C}' \rightarrow \mathcal{C}$ and we get a family of prestable curves $\mathcal{C}' \rightarrow \Delta$ where \mathcal{C}' is regular. By Lemma 10.1.3 and Corollary B.0.4, we can get a projective birational map $\mathcal{C}' \rightarrow \mathcal{C}'_{min}$ where \mathcal{C}'_{min} is semistable. So we replace \mathcal{C} by \mathcal{C}'_{min} . (This is the semistable reduction!) Using Proposition 7.5.1, we can get a relative canonical stabel model $\mathcal{C}' \rightarrow \mathcal{C}^{st}$.

10.2 Explicit stable reduction

Proposition 10.2.1. *Let $\mathcal{C} \rightarrow \Delta$ be a generically smooth, proper and flat family such that $(\mathcal{C}_0)_{red}$ is nodal. Let $\mathcal{C}_0 = \sum_i a_i D_i$ where a_i is the multiplicity of D_i . Let $\Delta' \rightarrow \Delta$ defined by $t \mapsto t^p$ where p prime and set $\mathcal{C}' = \mathcal{C} \times_{\Delta} \Delta'$. Then after taking normalization $\tilde{\mathcal{C}}' \rightarrow \mathcal{C}$ is branched cover ramified over $\sum_i (a_i \pmod{p}) D_i$.*

Example 2 (Stable reduction of A_{2k+1} -singularity). *Let $\mathcal{C} \rightarrow \Delta = \text{Spec}(R)$ be a generically smooth family degenerating to a A_{2k+1} -singularity in the central fiber where have local equation around the singular point is $y^2 = x^{2k+1} + t$. Now we will work through the steps in the proof of stable reduction. The first two steps have already finished, now we start at step 3.*

► **STEP 3. Use embedded resolutions to find a resolution of singularities $\tilde{\mathcal{C}} \rightarrow \mathcal{C}$ so that the reduced central fiber $(\tilde{\mathcal{C}}_0)_{red}$ is nodal.** We consider two charts in blowing up with coordinates x', y' where the original coordinates are x, y , as:

$$\begin{array}{ccc} E|_{U_1} = V(x') \hookrightarrow U_1 \hookrightarrow \tilde{\mathcal{C}} = \text{Bl}_p \mathcal{C} & (x', y') & E|_{U_2} = V(y') \hookrightarrow U_2 \hookrightarrow \tilde{\mathcal{C}} = \text{Bl}_p \mathcal{C} & (x', y') \\ \downarrow \swarrow & \downarrow & \downarrow \swarrow & \downarrow \\ \mathcal{C} & (x', x'y') & \mathcal{C} & (x'y', y') \end{array}$$

•**The first blowing up.** In the first chart, the preimage of $y^2 - x^{2k+1}$ is $x'^2 y'^2 - x'^{2k+1} = x'^2(y'^2 - x'^{2k-1})$; in the second chart it is $y'^2 - (x'y')^{2k+1} = y'^2(1 - x'^{2k+1}y'^{2k-1})$. Hence the exceptional divisor E_1 has multiplicity 2.

•**The second blowing up.** In the first chart, the preimage of $x^2(y^2 - x^{2k-1})$ is $x'^4(y'^2 - x'^{2k-3})$; in the second chart it is $x'^2 y'^4(1 - x'^{2k-1}y'^{2k-3})$. Hence the exceptional divisor E_2 has multiplicity 4.

•**After k blowing ups.** We get $x^{2k}(y^2 - x)$ with the exceptional divisors E_i has multiplicity $2i$.

•**One more blowing up.** We get the preimage of $x^{2k}(y^2 - x)$ in the second chart is

$$x'^{2k} y'^{2k+1} (y' - x')$$


with the exceptional divisor F has multiplicity $2k + 1$.

•**The final blowing up.** We get the preimage of $x^{2k} y^{2k+1} (y - x)$ in the first chart is

$$x'^{4k+2} y'^{2k+1} (y' - 1)$$

with the exceptional divisor G has multiplicity $4k + 2$.

The process as follows:




►**STEP 4. Perform a base change $\Delta' \rightarrow \Delta$ such that the normalization of the total family $\mathcal{C} \times_{\Delta} \Delta'$ has a reduced nodal central fiber with many rational tails and bridges.**

•**The first base change.** First consider $\Delta' \rightarrow \Delta, t \mapsto t^{2k+1}$ and normalizing. After inductively apply to the prime factorization $2k + 1$ and normalization, we will use the proposition to analyze the preimage of these irreducible component. Actually we get this $2k + 1$ -degree cover ramified over $C'_0 + \sum_i E_i$ and we just need to consider F, G . For G , its preimage G' ramified at two points (intersects E_k, C'_0) with index $2k$. By Riemann-Hurwitz Theorem, we get $2g(G') - 2 = (2k + 1)(2g(G) - 2) + 4k = -2$. Hence $g(G') = 0$ and $G' \cong \mathbb{P}^1$. For F , its preimage F' is unramified at all points, hence $F' = \coprod_{j=1}^{2k+1} F_j$ are copies of F . Hence replace Δ by Δ' , we get the central fiber as $C_0 = C'_0 + 2G' + \sum_j F_j + \sum_i 2iE_i$.


•**The second base change.** Consider $\Delta' \rightarrow \Delta, t \mapsto t^2$ and normalizing. Actually we get this 2-degree cover ramified over $C'_0 + \sum_j F_j$ and we just need to consider G', E_i . For G' , the preimage H ramified at $2k + 2$ points (C'_0, F_j). Hence we get $g(H) = k$ by Riemann-Hurwitz Theorem. For E_i , the things become more complicated as follows:

But we can easy to see that after this process, these things are just plenty of rational bridges and rational tails.




•**The final base changes.** Here we just need to consider E_i and these have multiplicity i . Consider $t \mapsto t^k$, then (many) E_k has two ramified points, hence by Riemann-Hurwitz Theorem $g(E'_k) = 0$, hence rational. Then consider $t \mapsto t^{k-1}, \dots, t \mapsto t^2$, we have the same results. Hence we also get plenty of rational tails and bridges, which are all multiplicity 1.

The whole process as follows:



►**STEP 5. Contract all rational tails and bridges in the central fiber.** Now we kill all -1 -curves (many E_1 and all F_j), and then (every) E_2 become -1 -curves. Inductively, we kill all E_i and F_j and get a stable central fiber as follows and we win.



10.3 Separatedness of $\overline{\mathcal{M}}_{g,n}$

Proposition 10.3.1. Let R be a DVR with fraction field K with $\Delta = \text{Spec}(R)$, $\Delta^* = \text{Spec}(K)$. Let $(\mathcal{C} \rightarrow \Delta, \sigma_1^*, \dots, \sigma_n^*)$ and $(\mathcal{D} \rightarrow \Delta, \tau_1^*, \dots, \tau_n^*)$ are n -pointed stable curves, then for any $\alpha^* : \mathcal{C}^* \rightarrow \mathcal{D}^*$ with $\tau_i^* = \alpha^* \circ \sigma_i^*$ over generic fiber can extend to a unique isomorphism $\alpha : \mathcal{C} \rightarrow \mathcal{D}$

with $\tau_i = \alpha \circ \sigma_i$.

$$\begin{array}{ccc}
 \mathcal{C}^* & \xrightarrow{\alpha^*} & \mathcal{D}^* \\
 \searrow & & \downarrow \\
 & & \Delta^*
 \end{array}
 \quad
 \begin{array}{ccc}
 \mathcal{C} & \dashrightarrow^{\alpha} & \mathcal{D} \\
 \searrow & & \downarrow \\
 & & \Delta
 \end{array}
 \quad
 \begin{array}{ccc}
 & & \Delta^* \\
 & \swarrow & \\
 & \Delta &
 \end{array}$$

Proof. We only prove the case of $n = 0$ generically smooth curves. Let $\mathcal{C}' \rightarrow \mathcal{C}, \mathcal{D}' \rightarrow \mathcal{D}$ be the minimal resolutions and let $\Gamma \subset \mathcal{C}' \times_{\Delta} \mathcal{D}'$ be the clpsure of the graph of $\text{id} \times \alpha^* : \mathcal{C}^* \rightarrow \mathcal{C}^* \times_{\Delta^*} \mathcal{D}^*$. Let $\Gamma' \rightarrow \Gamma$ be the minimal resolution. Hence we get birational projective maps $\Gamma' \rightarrow \mathcal{C}'$ and $\Gamma' \rightarrow \mathcal{D}'$. By the same proof of [23] Theorem II.8.19, we get

$$\Gamma(\mathcal{C}', \omega_{\mathcal{C}'/\Delta}^{\otimes k}) \cong \Gamma(\Gamma', \omega_{\Gamma'/\Delta}^{\otimes k}) \cong \Gamma(\mathcal{D}', \omega_{\mathcal{D}'/\Delta}^{\otimes k})$$

for all $k \geq 0$. As the canonical bundle are ample, we get

$$\mathcal{C}' \cong \text{Proj} \bigoplus_k \Gamma(\mathcal{C}', \omega_{\mathcal{C}'/\Delta}^{\otimes k}) \cong \text{Proj} \bigoplus_k \Gamma(\mathcal{D}', \omega_{\mathcal{D}'/\Delta}^{\otimes k}) \cong \mathcal{D}'.$$

Furthermore, we know that \mathcal{C}, \mathcal{D} are stable models of $\mathcal{C}', \mathcal{D}'$, respectively. By the uniqueness of stable models, we get $\alpha : \mathcal{C} \cong \mathcal{D}$ extending α^* . \square

Chapter 11

Gluing and forgetful morphisms

We follow [12].

11.1 Gluing morphisms

Proposition 11.1.1. *There are finite morphisms of algebraic stacks*

$$F : \overline{\mathcal{M}}_{i,n} \times \overline{\mathcal{M}}_{g-i,m} \rightarrow \overline{\mathcal{M}}_{g,n+m-2}$$


$$((C, p_1, \dots, p_n), (C', p'_1, \dots, p'_m)) \mapsto (C \cap C', p_1, \dots, p_{n-1}, p'_1, \dots, p'_m),$$

and

$$G : \overline{\mathcal{M}}_{g-1,n} \rightarrow \overline{\mathcal{M}}_{g,n-2}$$

$$(C, p_1, \dots, p_n) \mapsto (C/(p_{n-1} \sim p_n), p_1, \dots, p_{n-2}).$$

As follows:



Sketch-using pushout. By the stable reduction, these maps are of course representable and proper. As they have the finite fibers, these maps are now finite. Now for F we let $n = m = 1$ and for G we let $n = 2$.

For F : Let $(\pi : \mathcal{C} \rightarrow S, \sigma), (\pi' : \mathcal{C}' \rightarrow S, \sigma')$ are stable curves. As σ, σ' are closed immersions, we get the pushout exists by the theory of Ferrand (St 0ECH) and as we have the finite cover $\mathcal{C} \sqcup \mathcal{C}' \rightarrow \mathcal{C}$, we get this pushout is proper and flat (omitted):

$$\begin{array}{ccccc}
 & \text{Spec}(A) & \xrightarrow{\quad} & \text{Spec}A[y] & \\
 & \swarrow & & \swarrow & \\
 \text{Spec}A[x] & \xrightarrow{\quad} & \text{Spec}A[x, y]/(xy) & \xrightarrow{\quad} & \text{Spec}A[y] \\
 \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\
 \mathcal{C}' & \xrightarrow{\sigma'} & S & \xrightarrow{\sigma} & \mathcal{C} \\
 & \searrow & & \searrow & \\
 & \mathcal{C} & \xrightarrow{\quad} & \mathcal{C} &
 \end{array}$$

where $\text{Spec}A[x]$ is an étale neighborhood of $\sigma(s)$ which is the pulback of étale neighborhood $\text{Spec}(A)$ of any $s \in S$. Since an étale morphism from an affine scheme extend over closed immersions, there is an étale neighborhood $\text{Spec}A[y]$ is an étale neighborhood of $\sigma'(s)$. Then the pushout can be easy to compute as $\text{Spec}(A[x] \times_A A[x]) \cong \text{Spec}A[x, y]/(xy)$. By some results of pushout (in St 0D2G), we get $\text{Spec}A[x, y]/(xy) \rightarrow \mathcal{C}$ is an étale neighborhood of s . Hence $\tilde{\mathcal{C}} \rightarrow S$ is nodal along S . Checking fibers we get $\tilde{\mathcal{C}}_s$ is stable.

For G : Let $(\mathcal{C} \rightarrow S, \sigma_1, \sigma_2)$ are stable curve. Here we consider the pushout:

$$\begin{array}{ccc}
 S \sqcup S & \xrightarrow{\sigma_1 \sqcup \sigma_2} & \mathcal{C} \\
 \downarrow & & \downarrow \\
 S & \longrightarrow & \tilde{\mathcal{C}}
 \end{array}$$

which is étale locally like

$$\begin{array}{ccc}
 \text{Spec}(A \times A) & \xrightarrow{(0,1)} & \text{Spec}A[t] \\
 \downarrow & & \downarrow \\
 \text{Spec}A & \longrightarrow & \text{Spec}A[x, y]/(y^2 - x^2(x+1))
 \end{array}$$

where we find that $x := t^2 - 1, y = t^3 - t$ generate $A \times_{A \times A} A[t]$, then well done. \square

11.2 Boundary divisors of $\overline{\mathcal{M}}_g$

Consider the closed substacks

$$\begin{aligned}
 \delta_0 &= \text{Im}(\overline{\mathcal{M}}_{g-1,2} \rightarrow \overline{\mathcal{M}}_g) \\
 \delta_i &= \text{Im}(\overline{\mathcal{M}}_{i,1} \times \overline{\mathcal{M}}_{g-i,1} \rightarrow \overline{\mathcal{M}}_g)
 \end{aligned}$$

where $i = 1, \dots, \lfloor g/2 \rfloor$.

As these maps are finite, we get $\dim \delta_0 = \dim \overline{\mathcal{M}}_{g-1,2} = 3(g-1) - 3 + 2 = 3g - 4$ and similar $\dim \delta_i = 3g - 4$. Hence these are divisors of $\overline{\mathcal{M}}_g$.


(By analyzing the formal deformation space of a stable curve, one can show that $\delta = \bigcup_{j=0}^{\lfloor g/2 \rfloor} \delta_j$ is a normal crossings divisor.)

11.3 Forgetful morphisms

Proposition 11.3.1. *By Proposition 7.5.1, there is a morphism of algebraic stacks*

$$\overline{\mathcal{M}}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n-1}, (C, p_1, \dots, p_n) \mapsto (C^{st}, p_1, \dots, p_{n-1}).$$

As



11.4 Universal family $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$

This section we follow [12] and [13]. We consider the universal family $\mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n}$ of $\overline{\mathcal{M}}_{g,n}$. Actually the definition of universal family as for any family of stable curves $(\mathcal{C} \rightarrow S, \{\sigma_i\})$, we have the following universal property of cartesian

$$\begin{array}{ccc} \mathcal{C} & \dashrightarrow & \mathcal{U}_{g,n} \\ \downarrow & & \downarrow \\ S & \dashrightarrow^{\exists!} & \overline{\mathcal{M}}_{g,n} \end{array}$$

The existence given by 2-Yoneda's Lemma and some descent theory (omitted). Here we express this family as follows:

Lemma 11.4.1 (See [13]). *$\mathcal{U}_{g,n}(S)$ to be the set of families of curves $(\mathcal{C} \rightarrow S, \sigma_1, \dots, \sigma_n, \sigma)$ where $(\mathcal{C} \rightarrow S, \sigma_1, \dots, \sigma_n) \in \overline{\mathcal{M}}_{g,n}(S)$ and σ is an extra section without smooth condition.*

Proof. Fix $\pi : \mathcal{C} \rightarrow S$. We first let $\Sigma(S) : \mathcal{U}_{g,n}(S) \rightarrow \overline{\mathcal{M}}_{g,n}(S)$ as $(\pi, \sigma_i, \sigma) \mapsto (\pi, \sigma_i)$ be the canonical map and $\Sigma_i(S) : \overline{\mathcal{M}}_{g,n}(S) \rightarrow \mathcal{U}_{g,n}(S)$ as $(\pi, \sigma_1, \dots, \sigma_n) \mapsto (\pi, \sigma_1, \dots, \sigma_n; \sigma_i)$. Finally we need to define $\mathcal{C} \rightarrow \mathcal{U}_{g,n}$ as $(\text{pr}_2 : \mathcal{C} \times_S \mathcal{C} \rightarrow \mathcal{C}, s_i, \Delta)$ where $s_i = (\sigma_i \circ \pi, \text{id}_{\mathcal{C}})$ and $\Delta = (\text{id}_{\mathcal{C}}, \text{id}_{\mathcal{C}})$. Hence we get the following cartesian diagram of fibered categories

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \mathcal{U}_{g,n} \\ \sigma_1, \dots, \sigma_n \uparrow \left(\downarrow \pi \right. & & \left. \downarrow \Sigma \right) \uparrow \Sigma_1, \dots, \Sigma_n \\ S & \longrightarrow & \overline{\mathcal{M}}_{g,n} \end{array}$$

Well done. □

Now we consider

$$\overline{\mathcal{M}}_{g,n+1} \rightarrow \mathcal{U}_{g,n}, (\mathcal{C} \rightarrow S) \mapsto (\mathcal{C}^{st} \rightarrow S, \sigma'_1, \dots, \sigma'_n, \sigma')$$

where this stabilization aiming to make $(\mathcal{C}^{st} \rightarrow S, \sigma'_1, \dots, \sigma'_n)$ in $\overline{\mathcal{M}}_{g,n}(S)$.

$$\begin{array}{ccc} \overline{\mathcal{M}}_{g,n+1} & \longrightarrow & \mathcal{U}_{g,n} \\ & \searrow & \downarrow \\ & & \overline{\mathcal{M}}_{g,n} \end{array}$$

Remark 11.4.2 (More explicit construction). Fix $f : X \rightarrow S$ in $\overline{\mathcal{M}}_{g,n+1}(S)$ and hence we get

$$X = \underline{\text{Proj}}_S \left(\bigoplus_{m \geq 0} f_* \omega_{X/S} \left(\sum_{i=1}^{n+1} \sigma_i \right)^{\otimes m} \right).$$

Now we let

$$c(X) := \underline{\text{Proj}}_S \left(\bigoplus_{m \geq 0} f_* \omega_{X/S} \left(\sum_{i=1}^n \sigma_i \right)^{\otimes m} \right).$$

with $\sigma'_i : S \xrightarrow{\sigma_i} X \rightarrow c(X)$. Hence $(c(X); \sigma'_1, \dots, \sigma'_n)$ be a family of n -pointed stable curves. Hence we get $\overline{\mathcal{M}}_{g,n+1} \rightarrow \mathcal{U}_{g,n}$. Here we follows the proof before chapter 8 in [7].

Proposition 11.4.3. The morphism $\overline{\mathcal{M}}_{g,n+1} \rightarrow \mathcal{U}_{g,n}$ is an isomorphism over $\overline{\mathcal{M}}_{g,n}$.

Sketch. Now we construct an inverse map $\mathcal{U}_{g,n} \rightarrow \overline{\mathcal{M}}_{g,n+1}$.

Step 1. Construct that family of curves. Let $(\mathcal{C} \rightarrow S, \sigma_1, \dots, \sigma_n, \sigma)$ be an element in $\mathcal{U}_{g,n}(S)$. As σ is a closed immersion, it defined by an ideal sheaf $i : \mathcal{I}_\sigma \hookrightarrow \mathcal{O}_\mathcal{C}$. Define the coherent sheaf K by the exact sequence

$$0 \rightarrow \mathcal{O}_\mathcal{C} \xrightarrow{\delta} \mathcal{I}_\sigma^\vee \oplus \mathcal{O}_\mathcal{C}(\sigma_1 + \dots + \sigma_n) \rightarrow K \rightarrow 0$$

where $\delta = (i^\vee, j)$ where j is also an embedding. Now consider $\mathcal{C}' = \underline{\text{Proj}} \text{Sym}(K) \xrightarrow{p} \mathcal{C} \rightarrow S$.

Step 2. Construct the section. In [12], Knudsen introduce a notion called stably reflexive module. Knudsen separate the two cases of σ as this is local on S : (I) σ meets a non-smooth point in the fiber; (II) σ is a divisor meets one of these sections σ_i .

In both cases we find the surjections form as $\sigma^* K \rightarrow \sigma^*(-)$ or $\sigma_i^* K \rightarrow \sigma_i^*(-)$ to getting lifts where showed that all $\sigma^*(-)$ are line bundles (may using stably reflexive module). The picture when $S = \text{Spec}(k)$ as follows:



Hence we omitted all details and get $(\mathcal{C}' \rightarrow S, \sigma'_1, \dots, \sigma'_n, \sigma') \in \overline{\mathcal{M}}_{g,n+1}(S)$. For this detailed proof, we refer the original paper [12] Theorem 2.4 or the new paper [13]. One can also see [2] X.8 for more detailed proof over \mathbb{C} . \square

Chapter 12

Irreducibility

As $\overline{\mathcal{M}}_{g,n}$ is a smooth Deligne-Mumford stack, its irreducibility if and only if connectedness. As $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ be a universal family, it has connected fibers. Hence by induction, we can reduce the case of $\overline{\mathcal{M}}_g$. Moreover, by Keel-Mori theorem we get the coarse moduli space $\overline{\mathcal{M}}_g \rightarrow \overline{M}_g$ which induce the homeomorphism $|\overline{\mathcal{M}}_g| \cong |\overline{M}_g|$. Hence we can reduce the case of \overline{M}_g . Hence we have the following relations:

$$\begin{aligned} \overline{\mathcal{M}}_{g,n} \text{ irreducible} &\Leftrightarrow \overline{\mathcal{M}}_{g,n} \text{ connected} \Leftrightarrow \overline{\mathcal{M}}_g \text{ connected (or irreducible)} \\ &(\Leftrightarrow \mathcal{M}_g \text{ connected and dense in } \overline{\mathcal{M}}_g) \Leftrightarrow \overline{M}_g \text{ connected.} \end{aligned}$$

Here the denseness of \mathcal{M}_g in the proper Deligne-Mumford stack $\overline{\mathcal{M}}_g$ is called **Deligne-Mumford compactification**.

Remark 12.0.1 (Some historical remarks). (i) In 19th century, Clebsch and Hurwitz establishing irreducibility of M_g in characteristic 0 by using the classical topological argument;


(ii) In the appendix of the paper [21] by Fulton in [16] gives a completely algebraic proof for this in characteristic 0 in 1982;

(iii) In the paper [9], Deligne and Mumford give two arguments of irreducibility of $\overline{M}_{g,n}$ in characteristic p (by reduction to characteristic 0) in 1969;

(iv) In paper [15], Fulton established the irreducibility of $\overline{M}_{g,n}$ in characteristic p where $p > g + 1$ in 1969.

12.1 Preliminaries–Branched coverings

Definition 12.1.1. Let C be a connected smooth curve on k . A branched covering of \mathbb{P}_k^1 is a separable finite morphism $f : C \rightarrow \mathbb{P}_k^1$. We say f is simply branched if for any branched point $x \in \mathbb{P}_k^1$, there is at most one ramification point in the fiber $f^{-1}(x)$ and such a point has index 2.



Here (A) is simply branched but (B),(C) are not.

Lemma 12.1.2. *Let C be a smooth, connected and projective curve of genus g over an algebraically closed field k of characteristic 0. If L is a line bundle of degree $d \geq g + 1$ (*I think we may let $d \gg 0$*), then for a subspace $V \subset H^0(C, L)$ of dimension 2 we get $C \rightarrow \mathbb{P}^1$ a simply branched.*

Proof. (*This proof need to re-think.*) As $h^0(C, L) = d + 1 - g$, we get $\dim \text{Gr}(2, H^0(C, L)) = 2(d - g - 1)$. Here $\text{char}(k) = 0$, the map $C \rightarrow \mathbb{P}^1$ is finite separable. So $C \rightarrow \mathbb{P}^1$ is not a simply branched covering if and only if one of the following conditions holds

- (a) V has a base point;
- (b) there exists a ramification point with index > 2 ;
- (c) there exists 2 ramification points in the same fiber.

For (a), then there exists $p \in C$ such that for all $s \in V$ vanishing at p , that is, $s \in H^0(C, L(-p))$. The dimension of $V \in \text{Gr}(2, H^0(C, L))$ have this property is

$$\dim \text{Gr}(2, H^0(C, L(-p))) = 2d - 2g - 4.$$

For (b), then there exists $s \in V$ vanishing to order 3 at a point p , that is, $s \in H^0(C, L(-3p))$. The dimension of $V \in \text{Gr}(2, H^0(C, L))$ have this property is

$$\dim \mathbb{P}H^0(C, L(-3p)) + \dim \mathbb{P}(H^0(C, L)/(s)) = 2d - 2g - 4.$$

Hence varying $p \in C$, the locus of $\text{Gr}(2, H^0(C, L))$ failing (b) has dimension $\dim \text{Gr}(2, H^0(C, L)) - 1$;

For (c), then there exists independent $s_1, s_2 \in V$ vanishing to order 2 at a point p , that is, $s_1, s_2 \in H^0(C, L(-2p))$. The dimension of $V \in \text{Gr}(2, H^0(C, L))$ have this property is

$$\dim \text{Gr}(2, H^0(C, L(-2p))) = 2d - 2g - 6.$$

Hence varying $p \in C$, the locus of $\text{Gr}(2, H^0(C, L))$ failing (b) has dimension $- - -$ □

Lemma 12.1.3. *If $C \rightarrow \mathbb{P}^1$ is a simply branched cover of degree $d > 2$ in characteristic 0, then $\text{Aut}(C/\mathbb{P}^1)$ is trivial.*

Proof. Any $\alpha \in \text{Aut}(C/\mathbb{P}^1)$ must fix the $2g + 2d - 2$ branched points by Riemann-Hurwitz Theorem and simplyness. By Proposition 1.2.7, there are no non-trivial automorphisms of a smooth curve fixing more than $2g + 2$ points. Hence as $d > 2$, $\text{Aut}(C/\mathbb{P}^1)$ is trivial. □

Remark 12.1.4. *Here we give some notes for the proof of Clebsch and Hurwitz in 19th to show that M_g is connected over \mathbb{C} . We define*

$$H_{d,b} = \{C \rightarrow \mathbb{P}^1 \text{ simply branched covering of degree } d \text{ over } b \text{ points}\}$$

where $b = 2g + 2d - 2$. By the previous lemma, $H_{d,b}$ is an algebraic space or a topological space (if $k = \mathbb{C}$, *why?*). Let $\text{Sym}^b \mathbb{P}^1 \setminus \Delta$ as the variety of b unordered distinct points in \mathbb{P}^1 (which can also be written as the complement $\mathbb{P}^b \setminus \Delta$ of the discriminant hypersurface), we have a diagram

$$\begin{array}{ccc} & H_{d,b} & \\ \swarrow & & \searrow \\ \mathcal{M}_g & & \text{Sym}^b \mathbb{P}^1 \setminus \Delta \end{array}$$


with the canonical maps. Then they showed that $H_{d,b} \rightarrow \text{Sym}^b \mathbb{P}^1 \setminus \Delta$ is finite étale (actually this can be showed by using deformation theory pure algebraically, see [1] Lemma 5.7.9. We omitted here). By Lemma 12.1.2, we get $H_{d,b} \rightarrow \mathcal{M}_g$ is surjective. Hence we need to show that $H_{d,b}$ is connected. Combining these and some properties of monodromy theory, they proved this. For more detail, see [1] subsection 5.7.2.

12.2 Irreducibility over characteristic 0 using admissible covers

In this section we will use the method of admissible covers to gives a completely algebraic proof for the irreducibility in characteristic 0, which appears in the appendix of the paper [21] by Fulton in [16].

Proposition 12.2.1. *Let C be a smooth, connected and projective curve of genus g over an algebraically closed field k of characteristic 0. There exists a connected curve T with points $t_1, t_2 \in T$ and a family $\mathcal{C} \rightarrow T$ of stable curves such that $\mathcal{C}_{t_1} \cong C$ and \mathcal{C}_{t_2} is a singular stable curve.*


Proof. By Lemma 12.1.2 we get for $d \gg 0$ there exists a finite cover $C \rightarrow \mathbb{P}^1$ of degree d simply branched over $b = 2g + 2d - 2$ distinct points p_1, \dots, p_b in \mathbb{P}^1 . This gives a b -pointed stable curve $G = [\mathbb{P}^1, \{p_i\}] \in M_{0,b}$. By Remark 12.1.4 ($H_{d,b} \rightarrow \text{Sym}^b \mathbb{P}^1 \setminus \Delta$ is finite étale), we get $G \in M_{0,b}$ in general (WTF?). Then G can degenerates to (D_0, q_1, \dots, q_b) as the following picture



In the other words, there is a DVR R and fraction field K with $\Delta = \text{Spec}(R) \rightarrow \overline{M}_{0,n}$ be a stable curve $(\mathcal{D} \rightarrow \Delta, \sigma_i)$ with generic fiber (\mathbb{P}^1, p_i) and special fiber (D_0, q_1, \dots, q_b) . Hence we have a simply branched covering $\mathcal{C}^* \rightarrow \Delta^*$ and extend to $\mathcal{C} \rightarrow \mathcal{D}$ by taking \mathcal{C} as the integral closure of $\mathcal{O}_{\mathcal{D}}$ in $K(\mathcal{C}^*)$ as

$$\begin{array}{ccccc} \mathcal{C} & \dashrightarrow & \mathcal{D} & \longrightarrow & \Delta \\ \uparrow & & \uparrow & & \uparrow \\ \mathcal{C}^* & \longrightarrow & \mathcal{D}^* & \longrightarrow & \Delta^* \end{array}$$

Hence we get this diagram:



Now we just need to make \mathcal{C} be a singular stable curve. Purity of the branch locus (What's this?) implies the central fiber $\mathcal{C}_0 \rightarrow \mathcal{D}_0$ is ramified at $\sigma_i(0)$. By $\Delta' \rightarrow \Delta, t \mapsto t^m$ we can replace

\mathcal{C} such that $\mathcal{C}_0 \rightarrow \mathcal{D}_0$ is ramified only over $\sigma_i(0)$ and possibly over nodes of \mathcal{D}_0 . By an analysis of possible extensions $\mathcal{C} \rightarrow \mathcal{D}$, one can show that \mathcal{C}_0 is a nodal curve (missing details). Therefore $\mathcal{C} \rightarrow \Delta$ is a family of nodal curves.

Now we take $\mathcal{C} \rightarrow \mathcal{C}^{st}$ and just need to check \mathcal{C}_0^{st} is singular. For any irreducible component $T \subset \mathcal{C}_0^{st}$, apply Riemann-Hurwitz to $T \rightarrow \mathbb{P}^1 \subset \mathcal{D}_0$ we get $2g(T) - 2 = -2d + R$. If \mathbb{P}^1 is the middle one, we get $R \leq 2 + d - 1$; if \mathbb{P}^1 is the boundary one, we get $R \leq 1 + 2d - 2$. Hence $R \leq 2d - 1$ and $g(T) = 0$. Hence T is rational. Hence \mathcal{C}_0^{st} is singular. \square


Proposition 12.2.2. *If $\overline{\mathcal{M}}_{g',n'}$ is irreducible for all $g' < g$, then $\delta = \overline{\mathcal{M}}_{g,2} \setminus \mathcal{M}_{g,2}$ is connected.*

Proof. Let $\delta = \delta_0 \cup \delta_1 \cup \dots \cup \delta_{\lfloor g/2 \rfloor}$ where

$$\delta_0 = \text{Im}(\overline{\mathcal{M}}_{g-1,2} \rightarrow \overline{\mathcal{M}}_g)$$

$$\delta_i = \text{Im}(\overline{\mathcal{M}}_{i,1} \times \overline{\mathcal{M}}_{g-i,1} \rightarrow \overline{\mathcal{M}}_g)$$

where $i = 1, \dots, \lfloor g/2 \rfloor$. Hence δ_0, δ_i are connected by hypotheses. Easy to see that these divisors intersect as the points of $|\overline{\mathcal{M}}_g|$: \square



Theorem 12.2.3. *$\overline{\mathcal{M}}_{g,n}$ is irreducible.*

Proof. By the argument at beginning, we just need to show $\overline{\mathcal{M}}_g$ is connected. By Proposition 12.2.1 every smooth curve degenerates to a stable singular curve in the boundary $\delta = \overline{\mathcal{M}}_g \setminus \mathcal{M}_g$. By induction on g and Proposition 12.2.2 we get δ is connected, so is $\overline{\mathcal{M}}_g$. \square

Remark 12.2.4. *For the irreducibility in positive characteristic, we omitted and we refer the original [9], [15]. For the sketch, we refer subsection 5.7.4 in [1].*

Chapter 13

Projectivity

We will prove the coarse moduli space $\overline{M}_{g,n}$ is projective follows [26] and [35].

Remark 13.0.1. *Some generalizations of the projectivities:*

- (a) In [25] shows the moduli of stable varieties in any dimension is projective;
- (b) In [6] and [36] shows the moduli of K -polystable Fano varieties is projective.

Let the universal family $\pi : \mathcal{U}_g \rightarrow \overline{\mathcal{M}}_g$ and we define k -th pluri-canonical bundle as the vector bundle $\pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k})$. Indeed, $\pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k})$ is a coherent sheaf on the stack $\overline{\mathcal{M}}_g$ by the coherence theorem. We need to check that it is a vector bundle. By definition of the vector bundle over Deligne-Mumford stack, we need to show for any $S \rightarrow \overline{\mathcal{M}}_g$ the sheaf $(\pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k}))|_S$ is a vector bundle over S . As $S \rightarrow \overline{\mathcal{M}}_g$ correspond to $\pi_S : \mathcal{C} \rightarrow S$, we get

$$(\pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k}))|_S \cong \pi_{S,*}(\omega_{\mathcal{C}/S}^{\otimes k}).$$

By some argument with Review 1 we can show that $\pi_{S,*}(\omega_{\mathcal{C}/S}^{\otimes k})$ is a vector bundle.

Moreover, we get use the Riemann-Roch Theorem to deduce that

$$\text{rank}(\pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k})) = \begin{cases} g, & k = 1; \\ (2k-1)(g-1), & k > 1. \end{cases}$$

Now we consider the line bundle over $\overline{\mathcal{M}}_g$

$$\lambda_k := \det \pi_*(\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k}).$$

We will show that for $k \gg 0$, the line bundle λ_k descends to an ample line bundle on \overline{M}_g , then we get \overline{M}_g is a projective scheme.

13.1 Kollár's Criteria

Lemma 13.1.1. *Let \mathcal{X} be a proper Deligne-Mumford stack with coarse moduli space $\mathcal{X} \rightarrow X$. Suppose L line bundle over \mathcal{X} with*

- (a) L is semiample (i.e. L^N is basepoint-free for some $N > 0$);
- (b) for every proper integral curve T and map $f : T \rightarrow \mathcal{X}$ such that $f(T) \subset |\mathcal{X}|$ is not a single point, $\deg L|_T > 0$.

Then for some $N > 0$, $L^{\otimes N}$ descends to an ample line bundle over X .

Proof. This is the stack-version of the Corollary 1.2.15 in [29]. Actually we consider the following diagram which come from (a) and the universal property of coarse moduli space:

$$\begin{array}{ccc} \mathcal{X} & & \\ \downarrow \pi & \searrow f & \\ X & \xrightarrow{g} & \mathbb{P}(H^0(\mathcal{X}, L^{\otimes N})) \end{array}$$

By (b), f doesn't contract curves, so is g . Hence g is quasi-finite and proper, hence finite by Zariski main theorem. Hence $M := g^* \mathcal{O}(1)$ is ample; moreover, $\pi^* M = L^{\otimes N}$, we win. \square

Theorem 13.1.2 (Nakai-Moishezon Criterion). *If X is a proper algebraic space, a line bundle L is ample if and only if for all irreducible closed subvarieties $Z \subset X$,*

$$L^{\dim Z} \cdot Z > 0.$$

Proof. This is the algebraic space-version of the Theorem 1.21 in [8]. By Le Lemme de Gabber (Theorem C.2.1), there exists a finite surjection $f : X' \rightarrow X$ and by the algebraic space version St 0GFB we get L is ample if and only if $f^* L$ is ample. Hence by the scheme-version of Nakai-Moishezon Criterion ([8] Theorem 1.21), we win. \square

Let X be a proper algebraic space over k . Let $W \rightarrow Q$ be a surjection of vector bundles of rank w and q . Suppose that W has structure group $G \rightarrow \mathrm{GL}_w$. There is a classifying map

$$X \rightarrow [\mathrm{Gr}(q, w)/G], x \mapsto [W \otimes \kappa(x) \rightarrow Q \otimes \kappa(x)]$$

which is well defined because these killed by G .

Here we state our main theorem in this section. For simplicity, we only state it in characteristic 0. The criteria first appears in [26] and more general case we refer [25].

Theorem 13.1.3 (Kollár's Criterion). *Let X be a proper algebraic space over a field k of characteristic 0. Let $W \rightarrow Q$ be a surjection of vector bundles of rank w and q , where W has structure group $G \rightarrow \mathrm{GL}_w$. Suppose that*

- (a) *The classifying map $X(k) \rightarrow \mathrm{Gr}(q, w)(k)/G(k)$ has finite fibers;*
- (b) *W is nef.*

Then $\det Q$ is ample.

Proof. By Nakai-Moishezon criterion, for any irreducible subvariety $Z \subset X$ we need to verify $\det(Q)|_Z$ is big. As (a),(b) can restrict to Z , we can let X is an integral scheme and show that $\det Q$ is big.

By Le Lemme de Gabber (Theorem C.2.1), there exists a finite projective surjection $f : Y \rightarrow X$ of schemes. Hence we have $\det(f^* Q)^{\dim Y} = \deg(f) \det(Q)^{\dim X}$ and $\det Q$ is big if and only if $\det(f^* Q)$ is big. By taking the normalization, we can assume Y is normal and integral. So by Lemma 13.1.4 we win. \square

Lemma 13.1.4. *Let Y be a normal projective integral scheme over a field k of characteristic 0. Let $W \rightarrow Q$ be a surjection of vector bundles of rank w and q , where W has structure group $G \rightarrow \mathrm{GL}_w$. Suppose that*

- (a) *The classifying map $Y(k) \rightarrow \mathrm{Gr}(q, w)(k)/G(k)$ generically has finite fibers;*
- (b) *W is nef.*

Then $\det Q$ is big.

Sketch. To add. See Proposition 5.8.9 in [1]. \square

13.2 Nefness of pluri-canonical bundles

Theorem 13.2.1. *Let $\pi : \mathcal{C} \rightarrow T$ be a family of stable curves over a smooth curve T over k , then $\pi_*(\omega_{\mathcal{C}/T}^{\otimes k})$ is nef for $k \geq 2$.*

Proof. We following several steps:

•**Step 1. Reduction to characteristic p .** Now we let k is of characteristic 0. Since \mathcal{C} and T are finite type over k , their defining equations only involve finitely many coefficients of k . Thus there exists a finitely generated \mathbb{Z} -subalgebra $A \subset k$ and a cartesian diagram

$$\begin{array}{ccc} \mathcal{C} & \longrightarrow & \tilde{\mathcal{C}} \\ \downarrow & & \downarrow \\ T & \longrightarrow & \tilde{T} \\ \downarrow & & \downarrow \\ \text{Spec } k & \longrightarrow & \text{Spec } A \end{array}$$

where $\tilde{\mathcal{C}}, \tilde{T}$ are schemes of finite type over A . By possibly enlarging A , we can arrange that $\tilde{T} \rightarrow \text{Spec}(A)$ is smooth and projective family and $\tilde{\mathcal{C}} \rightarrow \tilde{T}$ is a family of stable curves.

(Need to re-think, omitted here.)

•**Step 2. Second reductions.** We reduce to the case that

- (a) \mathcal{C} is a smooth and minimal surface;
- (b) $\mathcal{C} \rightarrow T$ is generically smooth;
- (c) The genus of T is at least 2.

(To add.) These implies \mathcal{C} is of general type.

•**Step 3. Positive characteristic case.** Let $p = \text{char}(k)$. If $\pi_*(\omega_{\mathcal{C}/T}^{\otimes k})$ is not nef, then there exists a quotient line bundle $\pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \twoheadrightarrow M^\vee$ where $d = \deg(M) > 0$. Consider the absolute Frobenius

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\text{Frob}_{\mathcal{C}}} & \mathcal{C} \\ \downarrow & & \downarrow \\ T & \xrightarrow{\text{Frob}_T} & T \end{array}$$

By the property of the dualizing sheaf, we get $\text{Frob}_T^* \pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \cong \pi_*(\omega_{\mathcal{C}/T}^{\otimes k})$. And $\deg \text{Frob}_T^* M = pd$, we can let $d \gg 0$. Hence we can let $M = \omega_T^{\otimes k} \otimes L$ where L is very ample.

The surjection $\pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \twoheadrightarrow (\omega_T^{\otimes k} \otimes L)^\vee$ induce

$$\pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \otimes \omega_T^{\otimes k} \otimes L \twoheadrightarrow \mathcal{O}_T.$$

As $h^1(T, \mathcal{O}_T) \geq 2$, we have $h^1(T, \pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \otimes \omega_T^{\otimes k} \otimes L) \geq 2$. Use the Leray spectral sequence

$$H^1(T, \pi_*(\omega_{\mathcal{C}/T}^{\otimes k}) \otimes \omega_T^{\otimes k} \otimes L) \Rightarrow H^1(\mathcal{C}, \omega_{\mathcal{C}}^{\otimes k} \otimes \pi^* L),$$

hence $h^1(\mathcal{C}, \omega_{\mathcal{C}}^{\otimes k} \otimes \pi^* L) \geq 2$ by some calculation. By Lemma 13.2.2, we win. \square

Lemma 13.2.2 (Bombieri-Ekedahl). *Let S be a smooth projective surface over an algebraically closed field k which is minimal and of general type. Let D be an effective divisor with $D^2 = 0$. If $\text{char}(k) \neq 2$, then $H^1(S, \omega_S^{\otimes n}(D)) = 0$ for all $n \geq 2$. If $\text{char}(k) = 2$, then $h^1(S, \omega_S^{\otimes n}(D)) \leq 1$ for all $n \geq 2$.*

13.3 Positivity via positivity theory

For a morphism $S \rightarrow \overline{\mathcal{M}}_g$ correspond to $\mathcal{C} \rightarrow S$. Consider an integral d , we have

$$\mathrm{Sym}^d \pi_* (\omega_{\mathcal{C}/S}^{\otimes k}) \rightarrow \pi_* (\omega_{\mathcal{C}/S}^{\otimes dk}).$$

When $S = \mathrm{Spec}(K)$, $\mathcal{C} = C$, we get

$$\mathrm{Sym}^d H^0(C, \omega_C^{\otimes k}) \rightarrow H^0(C, \omega_C^{\otimes dk})$$

with kernel consists of degree d equations cutting out the image of $|\omega_C^{\otimes k}| : C \rightarrow \mathbb{P}^{r(k)-1}$. If $k \geq 3$, $\omega_{\mathcal{C}/S}^{\otimes k}$ is very ample and thus $\mathcal{C} \rightarrow S$ can be recovered from the kernel of the multiplication map.

Proposition 13.3.1. *For $k \gg 0$ and N sufficiently divisible, then $\lambda_k = \det \pi_* (\omega_{\mathcal{U}_g/\overline{\mathcal{M}}_g}^{\otimes k})$ descends to an ample line bundle on $\overline{\mathcal{M}}_g$.*

Proof. Consider $\mathcal{C} = \mathcal{U}_g$, $S = \overline{\mathcal{M}}_g$. Choose k, d such that

- (a) $\omega_{\mathcal{C}/S}^{\otimes k}$ is relatively very ample and $R^1 \pi_* \omega_{\mathcal{C}/S}^{\otimes k} = 0$;
- (b) Every curve $|\omega_C^{\otimes k}| : C \hookrightarrow \mathbb{P}^{r(k)-1}$ is cut out by equations degree d ;
- (c) $\pi_* (\omega_{\mathcal{C}/S}^{\otimes k})$ is nef (by Theorem 13.2.1).

These implies surjection

$$W := \mathrm{Sym}^d \pi_* (\omega_{\mathcal{C}/S}^{\otimes k}) \twoheadrightarrow \pi_* (\omega_{\mathcal{C}/S}^{\otimes dk}) =: Q.$$

Let w, q be the rank of W, Q , respectively. Let W has structure group $G \rightarrow \mathrm{GL}_w$. Consider the classifying map

$$\overline{\mathcal{M}}_g \rightarrow [\mathrm{Gr}(q, w)/G], x \mapsto \underbrace{[\mathrm{Sym}^d H^0(C, \omega_C^{\otimes k})]}_{\Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d))} \twoheadrightarrow \underbrace{H^0(C, \omega_C^{\otimes dk})}_{\Gamma(C, \mathcal{O}(d))}$$

is injective as the conditions on d and k imply that the kernel of the multiplication map uniquely determines C .

By Le Lemme de Gabber we get a finite cover $X \rightarrow \overline{\mathcal{M}}_g$. By Kollár's Criterion (Theorem 13.1.3), we get the pullback of λ_k to X is ample for $k \gg 0$. By Proposition C.2.2, we get for N sufficiently divisible, $\lambda_k^{\otimes N}$ descends to a line bundle L on $\overline{\mathcal{M}}_g$. Since the pullback of L under the finite morphism $X \rightarrow \overline{\mathcal{M}}_g \rightarrow \overline{\mathcal{M}}_g$, by St 0GFB we get the conclusion that L is ample. \square

Theorem 13.3.2. *If $2g - 2 + n > 0$, then $\overline{\mathcal{M}}_{g,n}$ is projective.*

Proof. The universal family $\overline{\mathcal{M}}_{g,n+1} \rightarrow \overline{\mathcal{M}}_{g,n}$ is projective by Proposition 7.3.3. Hence we just consider $n = 0$. This is right directly by the previous proposition. \square

Remark 13.3.3. *If we consider $\omega_{\mathcal{U}_{g,n}/\overline{\mathcal{M}}_{g,n}}^{\otimes k} (\Sigma_1 + \dots + \Sigma_n)$ from begining, we can prove the projectivity of $\overline{\mathcal{M}}_{g,n}$ directly.*

13.4 Projectivity via GIT, a sketch

By our old way, we have $\overline{\mathcal{M}}_g \cong [H'/\mathrm{PGL}_{r(k)}]$ for some locally closed $\mathrm{PGL}_{r(k)}$ -invariant subscheme of $\mathrm{Hilb}^P(\mathbb{P}^{r(k)-1})$ where $P(t) = \chi(C, \omega_C^{\otimes kt})$ and $r(k) = (2k - 1)(g - 1)$.

Remark 13.4.1. In fact we have $\overline{\mathcal{M}}_{g,n} \cong [H_{\nu,g,n}/\mathrm{PGL}(N)]$ where $N = (2\nu - 1)(g - 1) + \nu n$ and $H_{\nu,g,n} \subset \underline{\mathrm{Hilb}}^{P_\nu}(\mathbb{P}^{N-1})$ be the Hilbert scheme of ν -log-canonically embedded n -pointed stable curves of genus g where $P_\nu(t) = (2\nu t - 1)(g - 1) + \nu n t$ for $\nu \geq 3$. See [2] Theorem XII.5.6 for the proof.

Let H be the closure of H' in $\underline{\mathrm{Hilb}}^P(\mathbb{P}^{r(k)-1})$, By the proof of the representability of the quotient scheme, we get a closed immersion for $d \gg 0$:

$$\begin{aligned} H &\hookrightarrow \underline{\mathrm{Hilb}}^P(\mathbb{P}^{r(k)-1}) \hookrightarrow \mathrm{Gr}(P(d), \Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d))) \\ [C \hookrightarrow \mathbb{P}^{r(k)-1}] &\longmapsto [\Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d)) \twoheadrightarrow \Gamma(C, \mathcal{O}(d))] \end{aligned}$$

Next consider the Plücker embedding

$$\begin{aligned} \mathrm{Gr}(P(d), \Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d))) &\hookrightarrow \mathbb{P}\left(\bigwedge^{P(d)} \Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d))\right) \\ [\Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d)) \twoheadrightarrow \Gamma(C, \mathcal{O}(d))] &\mapsto \left[\bigwedge^{P(d)} \Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d)) \twoheadrightarrow \bigwedge^{P(d)} \Gamma(C, \mathcal{O}(d))\right] \end{aligned}$$

we can get $L_d := \mathcal{O}_{\mathrm{Gr}(P(d), \Gamma(\mathbb{P}^{r(k)-1}, \mathcal{O}(d)))}(1)|_H$ be the very ample line bundle over H . All these morphisms are $\mathrm{PGL}_{r(k)}$ -equivariant, hence L_d inherits a $\mathrm{PGL}_{r(k)}$ -linearization. Hence L_d can be defined on $[H/\mathrm{PGL}_{r(k)}]$.

Using the theory of Hilbert-Mumford criteria, we can prove the following difficult result.

Theorem 13.4.2. Let $k \geq 5$ and $d \gg 0$. For $h = [C \hookrightarrow \mathbb{P}^{r(k)-1}] \in H$, the curve C is stable if and only if $h \in H$ is GIT semistable with respect to L_d , that is, there exists an equivariant section $s \in \Gamma(H, L_d^{\otimes N})^{\mathrm{PGL}_{r(k)}}$ with $N > 0$ such that $s(h) \neq 0$. Moreover, we have

$$\overline{M}_g \cong \mathrm{Proj}\left(\Gamma(H, L_d^{\otimes N})^{\mathrm{PGL}_{r(k)}}\right),$$

hence projective.

Part III

Some geometry properties of the moduli space of curves

Chapter 14

Preliminaries

We now consider $\mathcal{M}_{g,n}$ and $\overline{\mathcal{M}}_{g,n}$ as the groupoid over the category $(Sch/Spec\mathbb{C})$. Then by the same arguments in the previous part, we can get $\overline{\mathcal{M}}_{g,n}$ is also a proper smooth Deligne-Mumford stack of dimension $3g - 3 + n$ over \mathbb{C} with a coarse moduli space $\overline{M}_{g,n}$ which is a projective variety over \mathbb{C} . Similarly for $\mathcal{M}_{g,n}$ and $M_{g,n}$. We will refer [2].

14.1 Graphs and dual graphs

We can associate a graph to a nodal curve with marked points.

Definition 14.1.1. *A graph Γ is the datum of:*

- (a) *a finite nonempty set $V = V(\Gamma)$ (the set of vertices);*
- (b) *a finite set $L = L(\Gamma)$ (the set of half-edges);*
- (c) *an involution ι of L ;*
- (d) *a partition of L indexed by V , that is, the assignment to each $v \in V$ of a (possibly empty) subset L_v of L such that $L = \bigcup_{v \in V} L_v$ with $L_v \cap L_w = \emptyset$ when $v \neq w$.*


A pair of distinct elements of L interchanged by the involution is called an edge of the graph. A fixed point of the involution is called a leg of the graph. The set of edges of Γ is denoted by $E(\Gamma)$. A dual graph is the datum of a graph together with the assignment of a nonnegative integer weight g_v to each vertex v . The genus of a dual graph Γ is defined to be

$$g_\Gamma = \sum_{v \in V(\Gamma)} g_v + 1 - \chi(\Gamma).$$

A graph (or a dual graph) endowed with a one-to-one correspondence between a finite set P and the set of its legs will be said to be P -marked, or numbered if P is of the form $\{1, \dots, n\}$ for some nonnegative integer n .

Let C be a nodal curve with a finite set D of smooth points of C . As in that case, there is a vertex for each component of the normalization of C , and its weight is the genus of the component. The half-edges issuing from a vertex are the points of the corresponding component which map either to a node of C or to a marked point. Easy to see that the edges of the graph are the pairs of half-edges mapping to the same node of C ; the legs are the half-edges coming from the marked points. This graph we denote it $\text{Graph}(C; D)$.

Easy to see by Theorem 6.2.1, we get the genus of the dual graph associated to $(C; D)$ is equal to the genus of C ! For example:




Remark 14.1.2. (i) Moreover, we also have some kind of localization. For $(C; D)$ we fixed a set S of nodes of C . We let a graph $\text{Graph}^S(C; D)$: The vertices of this graph are the connected components of the partial normalization C^S of C at S , the weight g_v is the arithmetic genus of the corresponding component of C^S , the edges correspond to the nodes in S , and the half-edges are the marked points or the points of C^S mapping to nodes in S .

(ii) We can also define the stable dual graph similarly.

Definition 14.1.3. Let Γ be a P -marked dual graph and let I be a subgraph having no legs and containing all the vertices of Γ . Let Γ_I be the graph that contracting each connected component of I to a point (one can speak this seriously, see [2] page 313). Hence we have a continuous map $c_I : \Gamma \rightarrow \Gamma_I$.

Actually there is a bijection between vertices of Γ_I and the connected components of I , we can let $g_w(\Gamma_I) = g(I_w)$ where w be a vertex of Γ_I and I_w be the corresponding connected component.

A P -marked dual graph Γ' is said to be a specialization of Γ if Γ is isomorphic to Γ'_I for some $I \subset \Gamma'$. We call $c : \Gamma' \rightarrow \Gamma \cong \Gamma'_I$ an I -contraction or simply a contraction. For example:




14.2 More on gluing morphisms

14.2.1 Gluing via graphs

► Gluing of curves.

Fix a P -marked dual graph Γ and for any $v \in V$ we give a L_v -pointed nodal curve C_v of genus g_v . Let $C' = \coprod_{v \in V} C_v$ and let $C = C' / \sim$ where \sim means two points need to gluing together if and only if they are marked points labeled by the two halves of an edge of Γ . Hence $C' \rightarrow C$ is actually a partial normalization. For example:



Here we need to note that this graph here is kind of partial diagram.

► **Gluing of families of curves.**

Fix a P -marked dual graph Γ and for any $v \in V$ we give a family of stable L_v -pointed genus g_v curves $F_v = (f_v : X_v \rightarrow S, \sigma_l \text{ where } l \in L_v)$. Let $X' = \coprod_v X_v$ and we get $F' = (f' : X \rightarrow S, \sigma_i)$ a family of L -pointed nodal curves.

For any $m \in L$, by taking residue along σ_m we get a surjection

$$\omega_{f'}^k(k \sum \sigma_l) \rightarrow \mathcal{O}_{\sigma_m(S)}.$$

Hence we get

$$R_l^{(k)} : f_* \left(\omega_{f'}^k(k \sum \sigma_l) \right) \rightarrow \mathcal{O}_S$$

by some kind of positivity (see [2] Lemma X.6.1(i)) it is surjective for all $k > 1$. Consider

$$R^{(k)} : f_* \left(\omega_{f'}^k(k \sum \sigma_l) \right) \rightarrow \mathcal{O}_S^E$$

indexed by pairs of edges $\{l, l'\}$ with components $R_l^{(k)} + (-1)^{k-1} R_{l'}^{(k)}$. The kernel of its fiber at $s \in S$ is $H^0(X_s, \omega_{f'}^k(k \sum_{p \in P} \sigma_{l_p}(s)))$ where X_s be the gluing of X'_s via Γ , hence its dimension is independent of s . Hence the kernel of $R^{(k)}$, which we denote by \mathcal{S}_k , is locally free. It is locally finitely generated (see [2] Corollary X.6.4). Let

$$X = \text{Proj}_S \bigoplus_{k \geq 0} \mathcal{S}_k$$

and hence the fibers of $X \rightarrow S$ is gluing via Γ . Let $\sigma'_p : S \rightarrow X$ is the composition of σ_{l_p} and $X' \rightarrow X$. Hence we get $F = (X \rightarrow S, \sigma'_p)_{p \in P}$.

14.2.2 Gluing functors

Fix a P -pointed dual genus g dual graph Γ and consider a Deligne-Mumford stack

$$\overline{\mathcal{M}}_\Gamma = \prod_{v \in V} \overline{\mathcal{M}}_{g_v, L_v}.$$

Fixed S and we let $\eta = (\eta_v)_{v \in V} \in \overline{\mathcal{M}}_\Gamma(S)$ where $\eta_v : X_v \rightarrow S$ be a family of stable L_v -pointed curves of genus g_v . The morphisms are isomorphisms between these families. Hence we get a gluing map via Γ to get $\xi_\Gamma(\eta) : X \rightarrow S$, a family of stable P -pointed genus g curves. Hence we get the gluing morphism of stacks


$$\xi_\Gamma : \overline{\mathcal{M}}_\Gamma \rightarrow \overline{\mathcal{M}}_{g, P}.$$

Let $\mathcal{D}_\Gamma \subset \overline{\mathcal{M}}_{g, P}$ be a closed substack as

$$\mathcal{D}_\Gamma(S) = \left\{ \sigma : X \rightarrow S \text{ families of } P\text{-pointed stable curves of genus } g : \begin{array}{l} \text{fibers have dual graphs which are specializations of } \Gamma \end{array} \right\}$$

(as the image of ξ_Γ). It is also a Deligne-Mumford stack with a coarse moduli space $\Delta_\Gamma \subset \overline{M}_{g, P}$ as a closed subvariety. We often refer to the \mathcal{D}_Γ (or the Δ_Γ) as the boundary strata of $\overline{\mathcal{M}}_{g, P}$ (or of $\overline{M}_{g, P}$).

The simplest boundary strata are those of codimension 1 (as a divisor as before), which correspond to the stable graphs with a single edge. If we consider the following graphs:



then for the first we let Γ_{irr} and the second $\Gamma_{a,A}$ (or $\Gamma_{\mathcal{P}}$ if $\mathcal{P} = \{(a, A), (b = g - a, B)\}$ be a stable bipartition). Hence we can also define $\mathcal{D}_{irr} := \mathcal{D}_{\Gamma_{irr}}$ and $\mathcal{D}_{a,A} := \mathcal{D}_{\Gamma_{a,A}}$ (or $\mathcal{D}_{\mathcal{P}}$). The coarse case are the same $\Delta_{irr}, \Delta_{a,A}, \Delta_{\mathcal{P}}$. Moreover, in the case we get the old gluing way:

$$\xi_{irr} : \overline{\mathcal{M}}_{g-1, P \cup \{x, y\}} \rightarrow \overline{\mathcal{M}}_{g, P}, \xi_{a, A} : \overline{\mathcal{M}}_{a, A \cup \{x\}} \times \overline{\mathcal{M}}_{g-a, A^c \cup \{y\}} \rightarrow \overline{\mathcal{M}}_{g, P}.$$

Definition 14.2.1 (Weak Γ -marking). *Consider a family of stable P -pointed genus g curves $(\pi : \mathcal{C} \rightarrow S, \tau_P)$. Let subvariety $\Sigma \subset \text{Sing}(\mathcal{C})$ proper and étale over S , then for any $s \in S$, the fiber Σ_s be a finite set of nodes. Hence we can consider $\text{Graph}^{\Sigma_s}(\mathcal{C}_s)$.*

Fix a P -marked graph Γ of genus g , if $\text{Graph}^{\Sigma_s}(\mathcal{C}_s) \cong \Gamma$ for any s , then we call Σ is a weak Γ -marking. Hence we can define a stack \mathcal{E}_{Γ} as

$$\mathcal{E}_{\Gamma}(S) = \left\{ \begin{array}{l} \pi : \mathcal{C} \rightarrow S \text{ families of } P\text{-pointed stable curves} \\ \text{of genus } g : \text{ endowed with a weak } \Gamma\text{-marking} \end{array} \right\}$$

Definition 14.2.2 (Γ -marking). *If $\mathcal{C} \rightarrow S$ coming from $(X \rightarrow S) \in \overline{\mathcal{M}}_{\Gamma}(S)$ by gluing via Γ with Σ the locus of nodes produced by gluing. As Σ be a union of sections on \mathcal{C} , so is the preimage over X (partial normalization). Hence we can get $\text{Graph}^{\Sigma}(\mathcal{C})$ with a family. Moreover $\Gamma \cong \text{Graph}^{\Sigma}(\mathcal{C})$. If these data exist for $\mathcal{C} \rightarrow S$, we called it endowed a Γ -marking.*

Hence we can see that in this case we can do it conversely, hence we have $\overline{\mathcal{M}}'_{\Gamma}$ as

$$\overline{\mathcal{M}}_{\Gamma}(S) \Leftrightarrow \left\{ \begin{array}{l} \pi : \mathcal{C} \rightarrow S \text{ families of } P\text{-pointed stable curves} \\ \text{of genus } g : \text{ endowed with a } \Gamma\text{-marking} \end{array} \right\} := \overline{\mathcal{M}}'_{\Gamma}(S)$$

Hence we can find that the gluing map can be composited as

$$\begin{array}{ccccc} \overline{\mathcal{M}}_{\Gamma} & & & & \\ \downarrow \cong & \searrow \xi_{\Gamma} & & & \\ \overline{\mathcal{M}}'_{\Gamma} & \xrightarrow{F} \mathcal{E}_{\Gamma} & \xrightarrow{F'} \mathcal{D}_{\Gamma} & \hookrightarrow & \overline{\mathcal{M}}_{g, P} \end{array}$$

where F, F' are forgetful maps.

Proposition 14.2.3. (i) \mathcal{E}_{Γ} be the normalization of substack $\mathcal{D}_{\Gamma} \subset \overline{\mathcal{M}}_{g, P}$;
(ii) The morphism $\overline{\mathcal{M}}_{\Gamma} \rightarrow \mathcal{E}_{\Gamma}$ can be identified with $\overline{\mathcal{M}}_{\Gamma} \rightarrow [\overline{\mathcal{M}}_{\Gamma} / \text{Aut}(\Gamma)]$.

Proof. See [2] Proposition XII.10.11. □

Corollary 14.2.4. We can seen $\text{Im}(\xi_{\Gamma}) = \mathcal{D}_{\Gamma}$ as before.

Proof. Trivial by the Proposition. □

Corollary 14.2.5. Let Γ be a stable P -marked dual graph of genus g . Assume that $\text{Aut}(\Gamma) = \{\text{id}_{\Gamma}\}$. Furthermore, assume that, for every graph Γ' which is a specialization of Γ , all the elements in $\text{Aut}(\Gamma')$ are specializations of id_{Γ} . Then $\xi_{\Gamma} : \overline{\mathcal{M}}_{\Gamma} \rightarrow \overline{\mathcal{M}}_{g, P}$ is a closed immersion.

Proposition 14.2.7. *If we let $\overline{\mathcal{M}}_{\Gamma\Gamma'} := \overline{\mathcal{M}}_{\Gamma} \times_{\overline{\mathcal{M}}_{g,P}} \overline{\mathcal{M}}_{\Gamma'}$, then*

$$\overline{\mathcal{M}}_{\Gamma\Gamma'} = \coprod_{\Lambda \in G_{\Gamma\Gamma'}} \overline{\mathcal{M}}_{\Lambda}.$$

Proof. Fix a scheme T .

First we let $\xi : \mathcal{C} \rightarrow T$ in $\overline{\mathcal{M}}_{\Lambda} = \overline{\mathcal{M}}'_{\Lambda}$, then we are given a subvariety $\Sigma \subset \text{Sing}(C)$, proper and étale over T , whose inverse image in the partial normalization along Σ itself is a union of sections, plus an isomorphism $\gamma : \text{Graph}^{\Sigma}(\mathcal{C}) \cong \Lambda$. Let contractions $c : \Lambda \rightarrow \Gamma$, $c' : \Lambda \rightarrow \Gamma'$ and

$$\Sigma_1 = (c \circ \gamma)^{-1}(E(\Gamma)), \Sigma_2 = (c' \circ \gamma)^{-1}(E(\Gamma'))$$

such that $\Sigma = \Sigma_1 \cup \Sigma_2$ with isomorphisms $\gamma_1 : \text{Graph}^{\Sigma_1}(\mathcal{C}) \cong \Gamma$ and $\gamma_2 : \text{Graph}^{\Sigma_2}(\mathcal{C}) \cong \Gamma'$. Hence ξ is both in $\overline{\mathcal{M}}_{\Gamma}(T)$ and $\overline{\mathcal{M}}_{\Gamma'}(T)$. Hence in $\overline{\mathcal{M}}_{\Gamma\Gamma'}(T)$

Conversely, as we have

$$\overline{\mathcal{M}}_{\Gamma\Gamma'}(T) = \left\{ \begin{array}{l} (\xi, \xi', \phi) : \xi, \xi' \text{ are families of } \Gamma, \Gamma' \text{-marking stable} \\ P\text{-pointed genus } g \text{ curves over } T \text{ with } \phi : \xi \rightarrow \xi' \\ \text{a } T\text{-isomorphism} \end{array} \right\}.$$

Then let $(\xi, \xi', \phi) \in \overline{\mathcal{M}}_{\Gamma\Gamma'}(T)$, hence we have $\gamma : \text{Graph}^{\Sigma_1}(\mathcal{C}) \cong \Gamma$ and $\gamma' : \text{Graph}^{\Sigma_2}(\mathcal{C}) \cong \Gamma'$. Hence we get contractions $c, c' : \text{Graph}^{\Sigma_1 \cup \Sigma_2}(\mathcal{C}) \rightarrow \Gamma, \Gamma'$. Hence we get $(\text{Graph}^{\Sigma_1 \cup \Sigma_2}(\mathcal{C}), c, c') \in G_{\Gamma\Gamma'}$, hence we win. \square

Chapter 15

Picard groups of the moduli space of curves

We will refer [2] chapter XIII.

Chapter 16

The kodaira dimension of moduli space of curves

We will refer [10] and [11] by using limit linear series instead of admissible covers in [21].

Part IV

Analytic construction via Teichmüller space

See [17] and [2] chapter XV.

Part V

Cellular decomposition and cohomology

See [2] chapter XVIII and XIX.

Part VI

Alterations and the moduli space of stable curves

See He Tongmu's Answer for now. And [7] maybe used.

Part VII

Appendix

Appendix A

Some basic result in scheme theory

A.1 Some corollaries of semi-continuity theorem

Review 1 (Cohomology and Base Change, see [23] III.12.11). *Let $f : X \rightarrow Y$ be a proper and finitely presented morphism of schemes with a finitely presented sheaf on X which is flat over Y . Let a point $y \in Y$ and $i \in \mathbb{Z}$, the comparison map $\phi_y^i : R^i f_* F \otimes \kappa(y) \rightarrow H^i(X_y, F_y)$ is surjective. Then*

(i) *There is an open neighborhood $V \subset Y$ of y such that for any morphism $g : Y' \rightarrow V$ of schemes, the comparison map $\phi_{Y'}^i : g^* R^i f_* F \rightarrow R^i f'_*(g')^* F$ is an isomorphism. In particular ϕ_y^i is an isomorphism;*

(ii) *ϕ_y^{i-1} is surjective if and only if $R^i f_* F$ is locally free in some neighborhood of y .*

Review 2 (Grauert's Corollary). (See [1] A.7.16) *Let $f : X \rightarrow Y$ be a flat proper morphism of noetherian schemes such that $h^0(X_y, \mathcal{O}_y) = 1$ for all $y \in Y$ ($\Leftrightarrow \mathcal{O}_Y = f_* \mathcal{O}_X$ and stable under base-change) (resp. the geometric fibers are integral).*

For a line bundle L on X , consider the functor $(\text{Sch}/Y) \rightarrow (\text{Sets})$ by sending $T \rightarrow Y$ to $\{\}$ if L_T is the pullback of a line bundle on T and to \emptyset otherwise. Then this functor is representable by a locally closed (resp. closed) subscheme of Y .*

A.2 Artin approximation and its corollaries

Definition A.2.1. *Let $A \rightarrow B$ of noetherian rings is called geometrically regular if it is flat and for every prime ideal $\mathfrak{p} \subset A$ and any finite field extension $K/\kappa(\mathfrak{p})$, the fiber $B \otimes_A K$ is regular.*

A noetherian local ring R is called a G -ring if $A \rightarrow \hat{A}$ is geometrically regular.

Theorem A.2.2 (Artin approximation, see [1] A.10.9). *Let S be a scheme and $s \in S$ be a point such that $\mathcal{O}_{S,s}$ is G -ring. Let $F : (\text{Sch}/S) \rightarrow (\text{Sets})$ be a colimit preserving contravariant functor (commutes with systems of \mathcal{O}_S -algebras) and $\hat{\xi} \in F(\text{Spec } \hat{\mathcal{O}}_{S,s})$. For any integer $N \geq 0$, there exists an étale morphism $(S', s') \rightarrow (S, s)$ and $\xi' \in F(S')$ with $\kappa(s) = \kappa(s)'$ such that the restrictions of $\hat{\xi}$ and ξ' to $\text{Spec}(\mathcal{O}_{S,s}/\mathfrak{m}_s^{N+1})$ are equal.*

Corollary A.2.3 (See [1] A.10.13). *Let X_1, X_2 be schemes of finite type over S and let $s \in S$ be a point such that $\mathcal{O}_{S,s}$ is a G-ring. If $x_1 \in X_1, x_2 \in X_2$ are points over s such that $\widehat{\mathcal{O}}_{X_1, x_1}$ and $\widehat{\mathcal{O}}_{X_2, x_2}$ are isomorphic as $\mathcal{O}_{S,s}$ -algebras, then there exists a common residually-trivial étale neighborhood as*

$$\begin{array}{ccc} & (X_3, x_3) & \\ \swarrow & & \searrow \\ (X_1, x_1) & & (X_2, x_2) \end{array}$$

A.3 Miscellany

Review 3. *Let k be a field and X be a proper geometrically connected and geometrically reduced k -scheme, then $\Gamma(X, \mathcal{O}_X) = k$.*

Proof. Just need to consider k is algebraically closed. Let $s \in \Gamma(X, \mathcal{O}_X) = \text{Hom}(X, \mathbb{A}_k^1)$ and consider the following diagram:

$$\begin{array}{ccccc} & & Z = s(X) & & \\ & \nearrow & & \searrow & \\ X & & & & \mathbb{A}_k^1 \hookrightarrow \mathbb{P}_k^1 \\ & \xrightarrow{s} & & & \\ & \searrow & & \swarrow & \\ & & \text{Spec}(k) & & \end{array}$$

Well done. □

Review 4 (Openness of ampleness). *Let $X \rightarrow S$ be a proper morphism of schemes and L be a line bundle over X . Let S is noetherian. If for some $s \in S$, the fiber L_s over X_s is ample (resp. very ample), then exists an open neighborhood U of s such that L_U is ample (resp. very ample) over X_U .*

Proposition A.3.1 (St 0C45). *Let X be a locally Noetherian scheme of dimension 1 with normalization $f : \tilde{X} \rightarrow X$. Then*

- (1) f is integral (finite if X is reduced locally finite type over a field), surjective, and induced a bijection on irreducible components;
- (2) there is a factorization $\tilde{X} \rightarrow X_{\text{red}} \rightarrow X$ and the morphism $\tilde{X} \rightarrow X_{\text{red}}$ is the normalization of X_{red} and birational;
- (3) for every closed point $x \in X$, stalk $(f_* \mathcal{O}_{\tilde{X}})_x$ is the integral closure of $\mathcal{O}_{X,x}$ in total ring of fractions of $(\mathcal{O}_{X,x})_{\text{red}} = \mathcal{O}_{X_{\text{red}},x}$;
- (4) \tilde{X} is a disjoint union of integral normal Noetherian schemes.

Proposition A.3.2 (0B5V). *Let R be a Noetherian ring. Let $f : X \rightarrow Y$ be a morphism of schemes proper over R . Let L be an invertible \mathcal{O}_Y -module. Assume f is finite and surjective. Then L is ample if and only if f^*L is ample.*

Appendix B

Some results of resolution of singularities for surfaces

Theorem B.0.1 (Minimal Resolutions). *Let X be a surface. There exists a unique projective birational morphism $\pi : \tilde{X} \rightarrow X$ from a smooth surface such that every other resolution $Y \rightarrow X$ factors as $Y \rightarrow \tilde{X} \rightarrow X$ (or equivalently such that $K_X \cdot E \geq 0$ for every π -exceptional curve E).*

Proof. See [27] Theorem 2.16. □

Theorem B.0.2 (Embedded Resolutions of Curves in Surfaces). *Let X be a surface and $X_0 \subset X$ be a curve. There is a finite sequence of blow-ups at reduced points of X_0 yielding a projective birational morphism $\tilde{X} \rightarrow X$ such that \tilde{X} is smooth and such that the preimage \tilde{X}_0 of X_0 has set-theoretic normal crossings, i.e. $(\tilde{X}_0)_{\text{red}}$ is nodal.*

Proof. See [27] Theorem 1.47. □

Theorem B.0.3 (Castelnuovo's Contraction Theorem). *Let X be a smooth projective surface and E a smooth rational (-1) -curve. Then there is a projective morphism $X \rightarrow Y$ to a smooth surface and a point $y \in Y$ such that $f^{-1}(y) = E$ and $X \setminus E \rightarrow Y \setminus \{y\}$ is an isomorphism.*

Proof. See [27] Theorem 2.14. □

Corollary B.0.4 (Existence of Relative Minimal Models). *A smooth surface X admits a projective birational morphism $X \rightarrow X_{\min}$ to a smooth surface such that every projective birational morphism $X_{\min} \rightarrow Y$ to a smooth surface is an isomorphism. In particular X_{\min} has no smooth rational (-1) -curves.*

Appendix C

Basic theory of algebraic spaces and stacks

C.1 Some basic facts

Theorem C.1.1. (See [32] 8.3.3) Let \mathcal{X}/S be an algebraic stack, then the following statement

- (a) \mathcal{X} is a Deligne-Mumford stack;
- (b) the diagonal $\Delta : \mathcal{X} \rightarrow \mathcal{X} \times_S \mathcal{X}$ is formally unramified;
- (c) for any algebraic closed field k and any point $x \in \mathcal{X}(k)$, the group scheme $\underline{\text{Aut}}_x$ is reduced finite k -scheme.

Then (a) \Leftrightarrow (b), and if \mathcal{X} noetherian, then (a) \Leftrightarrow (b) \Leftrightarrow (c).

Theorem C.1.2. Let \mathcal{X} be a smooth noetherian algebraic stack over k and $x \in \mathcal{X}(k)$ be a point with smooth stabilizer. Then

$$\dim_x \mathcal{X} = \dim T_{\mathcal{X},x} - \dim G_x.$$

Theorem C.1.3 (Valuative Criteria). Let $f : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of noetherian algebraic stacks. Assume f is of finite type and with separated diagonals. Then consider any DVR and its fraction field K with a 2-commutative diagram

$$\begin{array}{ccc} \text{Spec} K & \longrightarrow & \mathcal{X} \\ \downarrow & \nearrow \text{dashed} & \downarrow \\ \text{Spec} R & \longrightarrow & \mathcal{Y} \end{array}$$

Then

- (1) f is proper if and only if there exists an extension of DVRs $R \rightarrow R'$ and $K \rightarrow K'$ of fraction fields having finite transcendence degree and a lifting unique up to unique isomorphism

$$\begin{array}{ccccc} \text{Spec} K' & \longrightarrow & \text{Spec} K & \longrightarrow & \mathcal{X} \\ \downarrow & & \downarrow & \nearrow \text{dashed} & \downarrow \\ \text{Spec} R' & \longrightarrow & \text{Spec} R & \longrightarrow & \mathcal{Y} \end{array}$$

- (2) f is separated if and only if every two liftings of the first diagram are uniquely isomorphic.

(3) f is universally closed if for the first diagram, there exists an extension of DVRs $R \rightarrow R'$ and $K \rightarrow K'$ of fraction fields having finite transcendence degree and a lifting as in the second diagram.

Proof. See [1] Theorem 3.8.5 or St 0CLY. □

C.2 Miscellany

Theorem C.2.1 (Le Lemme de Gabber). *Let \mathcal{X} be a Deligne-Mumford stack separated and of finite type over a noetherian scheme S . Then there exists a finite, generically étale and surjective morphism $Z \rightarrow \mathcal{X}$ where Z be a scheme.*

Proposition C.2.2. *Let \mathcal{X} be a Deligne-Mumford stack separated and of finite type over a noetherian algebraic space S . Let $\pi : \mathcal{X} \rightarrow X$ be the coarse moduli space. If \mathcal{L} is a line bundle on \mathcal{X} , then for N sufficiently divisible $\mathcal{L}^{\otimes N}$ descends to X .*

Proof. See [1] Proposition 4.3.37. □

Bibliography

- [1] Jarod Alper. *Notes on Stacks and Moduli*. <https://sites.math.washington.edu/~jarod/moduli.pdf>, 2022-9-28.
- [2] Enrico Arbarello, Maurizio Cornalba, and Phillip A. Griffiths. *Geometry of Algebraic Curves, Volume II*, volume 268. Springer, 2011.
- [3] Enrico Arbarello, Maurizio Cornalba, Phillip A. Griffiths, and Joe Harris. *Geometry of Algebraic Curves, Volume I*, volume 267. Springer, 1985.
- [4] Michael Artin and Gayn Winters. Degenerate fibres and stable reduction of curves. *Topology*, 10(4):373–383, 1971.
- [5] The Stacks Project Authors. *The Stacks Project*. <https://stacks.math.columbia.edu/>, 2022.
- [6] G Codogni and Z Patakfalvi. Positivity of the cm line bundle for families of k-stable klt fano varieties. *Invent. math.*, 223:811–894, 2020.
- [7] Brian Conrad. Math 249b notes: Alterations. *Online Notes*, 2017.
- [8] Olivier Debarre. *Higher-Dimensional Algebraic Geometry*. Springer New York, NY, 2001.
- [9] Pierre Deligne and David Mumford. The irreducibility of the space of curves of given genus. *Publications Mathématiques de l’IHES*, 36:75–109, 1969.
- [10] David Eisenbud and Joe Harris. Limit linear series: Basic theory. *Invent. math.*, 85:337–371, 1986.
- [11] David Eisenbud and Joe Harris. The kodaira dimension of the moduli space of curves of genus ≥ 23 . *Invent. math.*, 90:359–387, 1987.
- [12] Finn F. Knudsen. The projectivity of the moduli space of stable curves, ii: The stacks $m_{g,n}$. *Mathematica Scandinavica*, 52(2):161–199, 1983.
- [13] Finn F. Knudsen. A closer look at the stacks of stable pointed curves. *Journal of Pure and Applied Algebra*, 216:2377–2385, 2012.
- [14] Lei Fu. *Algebraic Geometry*. Tsinghua University Press, 2006.
- [15] William Fulton. Hurwitz schemes and irreducibility of moduli of algebraic curves. *Ann. of Math.*, 90:542, 1969.

- [16] William Fulton. On the irreducibility of the moduli space of curves. *Invent. math.*, 67:87–88, 1982.
- [17] John H. Hubbard and Sarah Koch. An analytic construction of the Deligne-Mumford compactification of the moduli space of curves. *Journal of Differential Geometry*, 98:261 – 313, 2014.
- [18] Jack Hall. Moduli of singular curves. *Unknown*, 2010.
- [19] Jack Hall. The moduli stack of (all) curves. *Unknown*, 2013.
- [20] Joe Harris and Ian Morrison. *Moduli of curves*, volume 187. Springer Science & Business Media, 2006.
- [21] Joe Harris and David Mumford. On the kodaira dimension of the moduli space of curves. *Invent. math.*, 67:23–86, 1982.
- [22] Robin Hartshorne. *Residues and Duality*, volume 20. Springer Berlin, Heidelberg, 1966.
- [23] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer Science & Business Media, 1977.
- [24] Daniel Huybrechts. *Complex Geometry, An Introduction*. Springer Berlin, Heidelberg, 2005.
- [25] Sándor J. Kovács and Zsolt Patakfalvi. Projectivity of the moduli space of stable log-varieties and subadditivity of log-kodaira dimension. *J. Amer. Math. Soc.*, 30:959–1021, 2017.
- [26] János Kollár. Projectivity of complete moduli. *Journal of Differential Geometry*, 32:235–268, 1990.
- [27] János Kollár. *Lectures on Resolution of Singularities*, volume 166. PRINCETON UNIVERSITY PRESS, 2007.
- [28] Steven L. Kleiman. The picard scheme. *Arxiv: 0504020*, 2005.
- [29] Robert Lazarsfeld. *Positivity in Algebraic Geometry I*, volume 48. Springer Berlin, Heidelberg, 2004.
- [30] Qing Liu. *Algebraic Geometry and Arithmetic Curves*. Oxford University Press, USA, 2006.
- [31] Hideyuki Matsumura. *Commutative ring theory*. Cambridge university press, 1989.
- [32] Martin Olsson. *Algebraic spaces and stacks*, volume 62. American Mathematical Soc., 2016.
- [33] Edoardo Sernesi. *Deformations of Algebraic Schemes*. Springer-Verlag Berlin Heidelberg, 2006.
- [34] Ravi Vakil. *THE RISING SEA: Foundations of Algebraic Geometry*. Working Draft, August 29, 2022.
- [35] Eckart Viehweg. *Quasi-projective moduli for polarized manifolds*, volume 30. Springer, 1995.
- [36] Chenyang Xu and Ziquan Zhuang. On positivity of the cm line bundle on k-moduli spaces. *Ann. of Math*, 192:1005 – 1068, 2020.