

Title: Opioid Prescription Control CDS System

Author: Divya Verma

Variables:

[End_of_Session]

Confidence Variable

Prompt: END OF SESSION

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Message_1_The_recommendations_do_not_apply_]

Confidence Variable

Prompt: THE RECOMMENDATIONS ONLY APPLY TO OUTPATIENTS AGED 18 YEARS AND OLDER.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: True

Never Ask User: False

Display with results: True

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

HTML Link: <https://www.oregonpainguidance.org/opioidmedcalculator/>

Ask with: Edit Box

Arrange: One item per line

[Message_2_The_recommendations_exclude_the_conditions]

Confidence Variable

Prompt: THE RECOMMENDATIONS DO NOT APPLY TO PAIN RELATED TO SICKLE CELL DISEASE
OR CANCER OR TO PATIENTS RECEIVING PALLIATIVE OR END-OF-LIFE CARE.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Message_3_All_recommendations_are_met]

Confidence Variable

Prompt: ALL CDC RECOMMENDATIONS HAVE BEEN MET. PLEASE PROCEED TO PRESCRIBING
OPIOID THERAPY :)

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Q_A_Outpatient_Status]

Static List Variable

Prompt: Is the patient Outpatient?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q_B_Age]

Static List Variable

Prompt: Is the patient aged 18 or over?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q_C_Type_of_pain]

Static List Variable

Prompt: Which type of pain are you prescribing for?

Static List Values:

Acute_Pain_1_month

Acute Pain (<1 month)

Subacute_Pain_1_to_3_months_or_Chronic_Pain_3_months

Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q_D_Excluded_Conditions]

Static List Variable

Prompt: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q01_Opioid_Naive_acute_pain]

Static List Variable

Prompt: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event)

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q02_Opioid_Naive_subacute_or_chronic_pain]

Static List Variable

Prompt: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present]

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q03_Type_of_Opioid_analgesic]

Static List Variable

Prompt: Which type of Opioid analgesic you want to prescribe?

Static List Values:

1_Immediate-release_opioidseg_Suboxone_Sublingual_Film_Morphine_Sulfate

1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

2_Extended-release_and_longacting_ERLA_opioidseg_12_HR_Oxycodone_Hydrochloride_72_HR_Fentan

2) Extended-release and long-acting (ER/LA) opioids(e.g. 12 HR Oxycodone Hydrochloride, 72 HR Fentanyl)

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q0405_MME_dose]

Numeric Variable

Prompt: Enter MME dose Click here to Calculate Morphine Equivalent Dose

[Q06_Prescription_duration_in_days_for_Acute_pain]

Numeric Variable

Prompt: How many days are you prescribing for Acute pain?

[Q07_Opioid_risk_benefit_assessment]

Static List Variable

Prompt: Has Opioid risk benefit assessment been performed in the past 90 days?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q08A_Naloxone_Medications]

Static List Variable

Prompt: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q08B_Risks_of_opioid_overdose]

Static List Variable

Prompt: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day ≥ 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q09_PDMP_data_review_and_PDMP_review_procedure]

Static List Variable

Prompt: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q10A_Urine_drug_screening]

Static List Variable

Prompt: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q10B_Positive_drug_result_subacute_or_chronic_pain]

Static List Variable

Prompt: Is there positive result for either of following in the patient with subacute or chronic pain?- Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone.

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q11_Concurrent_medications]

Static List Variable

Prompt: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Q12_Evidence_of_Opioid_Use_disorder]

Static List Variable

Prompt: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days?

Static List Values:

Yes

Yes

No

No

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

[Rec#1_Consider_Nonpharmacologic_and_Nonopioid_Pharmacologic_Therapy]

Confidence Variable

Prompt: Recommendation #1 - Recommend use of nonpharmacologic therapy and nonopioid pharmacologic therapy as alternative. Nonopioid therapies are at least as effective as opioids for many common types of acute pain.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: True

Never Ask User: False

Display with results: True

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

HTML Link: <https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm?>

Ask with: Edit Box

Arrange: One item per line

[Rec#10A_Consider_Urine_Drug_Testing]

Confidence Variable

Prompt: Recommendation #10 - Urine Drug Screening is recommended. ORDER URINE DRUG TESTING or GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#10B_Positive_Cocaine_or_PCP_in_Urine_Screening]

Confidence Variable

Prompt: Recommendation #10 -Positive Cocaine or PCP in Urine Screening- this may be a false positive or indicate recreational or addictive use. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#11_Concurrent_Use_of_Opioids_and_Benzodiazepines]

Confidence Variable

Prompt: Recommendation #11 - Avoid prescribing opioid pain medication and benzodiazepine concurrently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#12_Evidencebased_Treatment_for_Patients_with_Opioid_Use_Disorder]

Confidence Variable

Prompt: Recommendation #12 - Recommend opioid agonist or partial agonist treatment with methadone maintenance therapy or buprenorphine and/or behavioral therapy. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#2_Opioid_Therapy_Goals_Discussion]

Confidence Variable

Prompt: Recommendation #2 - Maximize Nonopioid Therapies as Appropriate. RESOLVE, THEN GO
BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#3_Consider_Opioid_Immediate_Release_Form]

Confidence Variable

Prompt: Recommendation #3: Recommend use of immediate-release opioids instead of extended release/long acting opioids when starting patient on opioids. Record reason for opioid therapy OR Attempt alternative therapy. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#4#5_Lowest_Effective_Dose]

Confidence Variable

Prompt: Recommendations #4 and #5 -High risk for opioid overdose - taper now. Access MME conversation Tables at https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm#T1_down.
RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

HTML Link: https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm#T1_down

Ask with: Edit Box

Arrange: One item per line

[Rec#6_Prescribe_Lowest_Effective_Dose_and_Duration]

Confidence Variable

Prompt: Recommendation #6 - Limit therapy to ≤ 3 days to maximum of ≤ 7 days and to limit to immediate release opioids. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#7_Opioid_Therapy_Risk_Assessment]

Confidence Variable

Prompt: Recommendation #7A - Perform assessment of risk for opioid use procedure: Patients on opioid therapy should be evaluated for benefits and harms within 1 to 4 weeks of starting opioid therapy and every 3 months or more subsequently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#8_Naloxone_Consideration]

Confidence Variable

Prompt: Recommendation #8 - Incorporate into the management plan strategies to mitigate risk; including considering offering naloxone when factors that increase risk for opioid overdose are present.

Consider offering naloxone given the risk factor(s). RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec#9_Consider_Patients_History_of_Controlled_Substance_Prescriptions]

Confidence Variable

Prompt: Recommendation #9 - PDMP Data Review: Consider Patient's History of Controlled Substance Prescriptions. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.

Calculation Mode: Sum

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Display:

Ask with: Edit Box

Arrange: One item per line

[Rec_12_actions]

Static List Variable

Prompt: Recommendation #12 - Offer evidence-based medications to treat patients with opioid use disorder: Clinicians should offer or arrange treatment with evidence-based medications to treat patients with opioid use disorder. Detoxification on its own, without medications for opioid use disorder, is not recommended for opioid use disorder because of increased risks for resuming drug use, overdose, and overdose death. -----CHOOSE THE NEXT ACTION:

Static List Values:

Document__Will_order_medication_to_treat_opioid_use_disorder

Document - Will order medication to treat opioid use disorder

Refer__Will_refer_patient_to_SUD_specialist

Refer - Will refer patient to SUD specialist

Flags:

Always obtain a value: False

Display with results: False

Never Ask User: False

Display with results: False

Initialize: False

Check for PARAM data passed in Applet call: False

In backward chaining, stop after first value is set: False

In backward chaining, skip redundant rules: False

Use backward chaining to derive value: True

Use External Source to get value: False

Any number of values can be assigned

Display:

Ask with: Radio Buttons

Arrange: One item per line

Logic Block: Try to initiate opioid or not

```
1 | Q_A_Outpatient_Status = No
2 |   --> [Message_1_The_recommendations_do_not_apply_] = 10
3 | Q_A_Outpatient_Status = Yes
4 |   Q_B_Age = Yes
5 |     Q_C_Type_of_pain = Acute_Pain_1_month
6 |       Q_D_Excluded_Conditions = No
7 |         Q01_Opioid_Naive_acute_pain = Yes
8 |           --> [Rec#1_Consider_Nonpharmacologic_and_Nonopioid_Pharmacologic_Therapy]
9 |             = 10
10 |         Q01_Opioid_Naive_acute_pain = No
11 |           Q03_Type_of_Opioid_analgesic =
12 |             1_Immediate-release_opioidseg_Suboxone_Sublingual_Film_Morphine_Sulfate
13 |               [Q0405_MME_dose] <= 50
14 |                 [Q06_Prescription_duration_in_days_for_Acute_pain] <= 7
15 |                   Q08A_Naloxone_Medications = Yes
16 |                     Q09_PDMP_data_review_and_PDMP_review_procedure = No
17 |                       --> [Rec#9_Consider_Patients_History_of_Controlled_Substance_Pre
18 |                         scriptions] = 10
19 |                     Q09_PDMP_data_review_and_PDMP_review_procedure = Yes
20 |                       Q11_Concurrent_medications = No
21 |                         Q12_Evidence_of_Opioid_Use_disorder = Yes
22 |                           --> ASK [Rec_12_actions]
23 |                           --> [End_of_Session] = 10
24 |                         Q12_Evidence_of_Opioid_Use_disorder = No
25 |                           --> [Message_3_All_recommendations_are_met] = 10
26 |                       Q11_Concurrent_medications = Yes
27 |                         --> [Rec#11_Concurrent_Use_of_Opioids_and_Benzodiazepines] =
28 |                           10
29 |                     Q08A_Naloxone_Medications = No
30 |                       Q08B_Risks_of_opioid_overdose = Yes
31 |                         --> [Rec#8_Naloxone_Consideration] = 10
32 |                       Q08B_Risks_of_opioid_overdose = No
33 |                         Q09_PDMP_data_review_and_PDMP_review_procedure = Yes
34 |                           --> [Rec#9_Consider_Patients_History_of_Controlled_Substance_P
35 |                             rescriptions] = 10
36 |                         Q09_PDMP_data_review_and_PDMP_review_procedure = No
37 |                           Q11_Concurrent_medications = No
38 |                             Q12_Evidence_of_Opioid_Use_disorder = Yes
39 |                               --> ASK [Rec_12_actions]
40 |                               --> [End_of_Session] = 10
41 |                             Q12_Evidence_of_Opioid_Use_disorder = No
42 |                               --> [Message_3_All_recommendations_are_met] = 10
43 |                           Q11_Concurrent_medications = Yes
44 |                             --> [Rec#11_Concurrent_Use_of_Opioids_and_Benzodiazepines]
45 |                               = 10
46 |                         [Q06_Prescription_duration_in_days_for_Acute_pain] > 7
47 |                           --> [Rec#6_Prescribe_Lowest_Effective_Dose_and_Duration] = 10
48 |                       [Q0405_MME_dose] > 50
49 |                         --> [Rec#4#5_Lowest_Effective_Dose] = 10
50 |                   Q03_Type_of_Opioid_analgesic =
51 |                     2_Extended-release_and_longacting_ERLA_opioidseg_12_HR_Oxycodone_Hydroc
52 |                       hloride_72_HR_Fentanyl_
```

```

45      --> [Rec#3_Consider_Opioid_Immediate_Release_Form] = 10
46      Q_D_Excluded_Conditions = Yes
47      --> [Message_2_The_recommendations_exclude_the_conditions] = 10
48      Q_C_Type_of_pain = Subacute_Pain_1_to_3_months_or_Chronic_Pain_3_months
49      Q_D_Excluded_Conditions = No
50      Q02_Opioid_Naive_subacute_or_chronic_pain = Yes
51      --> [Rec#2_Opioid_Therapy_Goals_Discussion] = 10
52      Q02_Opioid_Naive_subacute_or_chronic_pain = No
53      Q03_Type_of_Opioid_analgesic =
54          1_ImmediateRelease_opioidseg_Suboxone_Sublingual_Film_Morphine_Sulfate
55          [Q0405_MME_dose] <=50
56          Q07_Opioid_risk_benefit_assessment = No
57          --> [Rec#7_Opioid_Therapy_Risk_Assessment] = 10
58          Q07_Opioid_risk_benefit_assessment = Yes
59          Q08A_Naloxone_Medications = Yes
60          Q09_PDMP_data_review_and_PDMP_review_procedure = No
61          --> [Rec#9_Consider_Patients_History_of_Controlled_Substance_Pre
62              scriptions] = 10
63          Q09_PDMP_data_review_and_PDMP_review_procedure = Yes
64          Q10A_Urine_drug_screening = Yes
65          Q10B_Positive_drug_result_subacute_or_chronic_pain = Yes
66          --> [Rec#10B_Positive_Cocaine_or_PCP_in_Urine_Screening] =
67              10
68          Q10B_Positive_drug_result_subacute_or_chronic_pain = No
69          Q11_Concurrent_medications = Yes
70          --> [Rec#11_Concurrent_Use_of_Opioids_and_Benzodiazepin
71              es] = 10
72          Q11_Concurrent_medications = No
73          Q12_Evidence_of_Opioid_Use_disorder = Yes
74          --> ASK [Rec_12_actions]
75          --> [End_of_Session] = 10
76          Q12_Evidence_of_Opioid_Use_disorder = No
77          --> [Message_3_All_recommendations_are_met] = 10
78          Q10A_Urine_drug_screening = No
79          --> [Rec#10A_Consider_Urine_Drug_Testing] = 10
80          Q08A_Naloxone_Medications = No
81          Q08B_Risks_of_opioid_overdose = No
82          Q09_PDMP_data_review_and_PDMP_review_procedure = No
83          --> [Rec#9_Consider_Patients_History_of_Controlled_Substance_P
84              rescriptions] = 10
85          Q09_PDMP_data_review_and_PDMP_review_procedure = Yes
86          Q10A_Urine_drug_screening = Yes
87          Q10B_Positive_drug_result_subacute_or_chronic_pain = Yes
88          --> [Rec#10A_Consider_Urine_Drug_Testing] = 10
89          Q10B_Positive_drug_result_subacute_or_chronic_pain = No
90          Q11_Concurrent_medications = Yes
91          --> [Rec#11_Concurrent_Use_of_Opioids_and_Benzodiaze
              pines] = 10
          Q11_Concurrent_medications = No
          Q12_Evidence_of_Opioid_Use_disorder = Yes
          --> ASK [Rec_12_actions]
          --> [End_of_Session] = 10
          Q12_Evidence_of_Opioid_Use_disorder = No
    
```

```

92      --> [Message_3_All_recommendations_are_met] = 10
93      Q10A_Urine_drug_screening = No
94      --> [Rec#10A_Consider_Urine_Drug_Testing] = 10
95      Q08B_Risks_of_opioid_overdose = Yes
96      --> [Rec#8_Naloxone_Consideration] = 10
97      [Q0405_MME_dose] >50
98      --> [Rec#4#5_Lowest_Effective_Dose] = 10
99      Q03_Type_of_Opioid_analgesic =
      2_Extendedrelease_and_longacting_ERLA_opioidseg_12_HR_Oxycodone_Hydroc
      hloride_72_HR_Fentanyl_
      --> [Rec#3_Consider_Opioid_Immediate_Release_Form] = 10
100      Q_D_Excluded_Conditions = Yes
101      --> [Message_2_The_recommendations_exclude_the_conditions] = 10
102      Q_B_Age = No
103      --> [Message_1_The_recommendations_do_not_apply_] = 10
104

```


Command Block: Default Command Block

```
1  DERIVE CONF  // Derive the value of all
2  RESULTS
```


Rules:

Block: Try to initiate opioid or not Row:2

IF:

Is the patient Outpatient? No

THEN:

THE RECOMMENDATIONS ONLY APPLY TO OUTPATIENTS AGED 18 YEARS AND OLDER.: Confidence =10

Block: Try to initiate opioid or not Row:8

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) Yes

THEN:

Recommendation #1 - Recommend use of nonpharmacologic therapy and nonopioid pharmacologic therapy as alternative. Nonopioid therapies are at least as effective as opioids for many common types of acute pain.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No

THEN:

Recommendation #9 - PDMP Data Review: Consider Patient's History of Controlled Substance Prescriptions. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No
- AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? Yes

THEN:

ASK [Rec_12_actions]
END OF SESSION: Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No
- AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? No

THEN:

ALL CDC RECOMMENDATIONS HAVE BEEN MET. PLEASE PROCEED TO PRESCRIBING OPIOID THERAPY :): Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? Yes

THEN:

Recommendation #11 - Avoid prescribing opioid pain medication and benzodiazepine concurrently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition Yes

THEN:

Recommendation #8 - Incorporate into the management plan strategies to mitigate risk; including considering offering naloxone when factors that increase risk for opioid overdose are present. Consider offering naloxone given the risk factor(s). RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes
AND: Is the patient aged 18 or over? Yes
AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
AND: [Q0405_MME_dose] <=50
AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No
AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No
AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

THEN:

Recommendation #9 - PDMP Data Review: Consider Patient's History of Controlled Substance Prescriptions. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No
- AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No
- AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? Yes

THEN:

ASK [Rec_12_actions]
END OF SESSION: Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No
- AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No
- AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? No

THEN:

ALL CDC RECOMMENDATIONS HAVE BEEN MET. PLEASE PROCEED TO PRESCRIBING OPIOID THERAPY :); Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Acute Pain (<1 month)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: [Q06_Prescription_duration_in_days_for_Acute_pain] <=7
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No
- AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? Yes

THEN:

Recommendation #11 - Avoid prescribing opioid pain medication and benzodiazepine concurrently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: [Q06_Prescription_duration_in_days_for_Acute_pain] >7

THEN:

Recommendation #6 - Limit therapy to <=3 days to maximum of <= 7 days and to limit to immediate release opioids. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] >50

THEN:

Recommendations #4 and #5 -High risk for opioid overdose - taper now. Access MME conversation Tables at https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm#T1_down. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.:
Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with acute pain opioid naive? (i.e. No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event) No

AND: Which type of Opioid analgesic you want to prescribe? 2) Extended-release and long-acting (ER/LA) opioids(e.g. 12 HR Oxycodone Hydrochloride, 72 HR Fentanyl)

THEN:

Recommendation #3: Recommend use of immediate-release opioids instead of extended release/long acting opioids when starting patient on opioids. Record reason for opioid therapy OR Attempt alternative therapy. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Acute Pain (<1 month)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? Yes

THEN:

THE RECOMMENDATIONS DO NOT APPLY TO PAIN RELATED TO SICKLE CELL DISEASE OR CANCER OR TO PATIENTS RECEIVING PALLIATIVE OR END-OF-LIFE CARE.: Confidence = 10

Block: Try to initiate opioid or not Row:51

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] Yes

THEN:

Recommendation #2 - Maximize Nonopioid Therapies as Appropriate. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? No

THEN:

Recommendation #7A - Perform assessment of risk for opioid use procedure: Patients on opioid therapy should be evaluated for benefits and harms within 1 to 4 weeks of starting opioid therapy and every 3 months or more subsequently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

Block: Try to initiate opioid or not Row:60

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No

THEN:

Recommendation #9 - PDMP Data Review: Consider Patient's History of Controlled Substance Prescriptions. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?-
Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid
OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate
OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. Yes

THEN:

Recommendation #10 -Positive Cocaine or PCP in Urine Screening- this may be a false positive or indicate recreational or addictive use. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?-
Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid
OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate
OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? Yes

THEN:

Recommendation #11 - Avoid prescribing opioid pain medication and benzodiazepine concurrently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?-
Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid
OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate
OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No

AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? Yes

THEN:

ASK [Rec_12_actions]

END OF SESSION: Confidence = 10

Block: Try to initiate opioid or not Row:73

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?- Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No

AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? No

THEN:

ALL CDC RECOMMENDATIONS HAVE BEEN MET. PLEASE PROCEED TO PRESCRIBING OPIOID THERAPY :): Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? Yes

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? No

THEN:

Recommendation #10 - Urine Drug Screening is recommended. ORDER URINE DRUG TESTING or GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.:
Confidence = 10

Block: Try to initiate opioid or not Row:79

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs No

THEN:

Recommendation #9 - PDMP Data Review: Consider Patient's History of Controlled Substance Prescriptions. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?-
Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid
OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate
OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. Yes

THEN:

Recommendation #10 - Urine Drug Screening is recommended. ORDER URINE DRUG TESTING or GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.:
Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?- Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? Yes

THEN:

Recommendation #11 - Avoid prescribing opioid pain medication and benzodiazepine concurrently. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes

AND: Is there positive result for either of following in the patient with subacute or chronic pain?- Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No

AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No

AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? Yes

THEN:

ASK [Rec_12_actions]

END OF SESSION: Confidence = 10

IF:

- Is the patient Outpatient? Yes
- AND: Is the patient aged 18 or over? Yes
- AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)
- AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No
- AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No
- AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)
- AND: [Q0405_MME_dose] <=50
- AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes
- AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No
- AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No
- AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes
- AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? Yes
- AND: Is there positive result for either of following in the patient with subacute or chronic pain?- Positive result for synthetic opioid in the past year and not currently prescribed synthetic opioid OR-Positive result for phencyclidine(PCP) in the past year OR-Positive result for fentanyl in the past year and not currently prescribed fentanyl OR-Positive result for cocaine in the past year OR-Positive result for amphetamine in the past year and not currently prescribed amphetamine OR-Positive result for tetrahydrocannabinol (THC) in the past year and not currently prescribed tetrahydrocannabinol (THC) OR-Positive result for opiate in the past year and not currently prescribed natural opiate OR-Positive result for alcohol in the past year OR-Positive result for methadone in the past year and not currently prescribed methadone. No
- AND: Is there any concurrent medications usage of a benzodiazepine OR other central nervous system depressant medication? No
- AND: Any current Opioid Use disorder OR order for Buprenorphine or Methadone medication in the past 90 days? No

THEN:

ALL CDC RECOMMENDATIONS HAVE BEEN MET. PLEASE PROCEED TO PRESCRIBING OPIOID THERAPY :): Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition No

AND: Has Prescription Drug Monitoring Program(PDMP) data been reviewed in the past 90 days and has PDMP review procedure been performed in the past 90 days? Click here to access PDMPs Yes

AND: Has urine drug screening (Nonopioid or Opioid drug) been performed in last 12 months? No

THEN:

Recommendation #10 - Urine Drug Screening is recommended. ORDER URINE DRUG TESTING or GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.:
Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] <=50

AND: Has Opioid risk benefit assessment been performed in the past 90 days? Yes

AND: Has patient been prescribed Naloxone medications to reduce or reverse opioid overdose? No

AND: Are any of the following factors that increase risk for opioid overdose present? - 1) Currently receiving benzodiazepine or other central nervous system depressant medications, or 2) High opioid dosages (MME/day >= 50), or 3) History of substance use disorder, or 4) Diagnosed with sleep-disordered breathing condition Yes

THEN:

Recommendation #8 - Incorporate into the management plan strategies to mitigate risk; including considering offering naloxone when factors that increase risk for opioid overdose are present. Consider offering naloxone given the risk factor(s). RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 1) Immediate-release opioids(e.g. Suboxone Sublingual Film, Morphine Sulfate)

AND: [Q0405_MME_dose] >50

THEN:

Recommendations #4 and #5 -High risk for opioid overdose - taper now. Access MME conversation Tables at https://www.cdc.gov/mmwr/volumes/71/rr/rr7103a1.htm#T1_down. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.:
Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? No

AND: Is the patient with Subacute or Chronic opioid naive?-- [i.e. 1) No indication of opioid use in the past 90 days from a prescription, self report, or a dispensing event ; 2) Pain management procedure in past 90 days is not present; 3) Pain treatment plan created, edited, or reviewed by the prescribing physician in past 90 days is not present] No

AND: Which type of Opioid analgesic you want to prescribe? 2) Extended-release and long-acting (ER/LA) opioids(e.g. 12 HR Oxycodone Hydrochloride, 72 HR Fentanyl)

THEN:

Recommendation #3: Recommend use of immediate-release opioids instead of extended release/long acting opioids when starting patient on opioids. Record reason for opioid therapy OR Attempt alternative therapy. RESOLVE, THEN GO BACK AND CHOOSE THE CORRECT CHOICE TO CONTINUE THE SESSION.: Confidence = 10

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? Yes

AND: Which type of pain are you prescribing for? Subacute Pain (1 to 3 months) or Chronic Pain (>3 months)

AND: Is there an evidence of sickle cell disease or limited life expectancy or palliative or end-of-life care or active cancer treatment? Yes

THEN:

THE RECOMMENDATIONS DO NOT APPLY TO PAIN RELATED TO SICKLE CELL DISEASE OR CANCER OR TO PATIENTS RECEIVING PALLIATIVE OR END-OF-LIFE CARE.: Confidence = 10

Block: Try to initiate opioid or not Row:104

IF:

Is the patient Outpatient? Yes

AND: Is the patient aged 18 or over? No

THEN:

THE RECOMMENDATIONS ONLY APPLY TO OUTPATIENTS AGED 18 YEARS AND
OLDER.: Confidence = 10