

MATH50003


Numerical Analysis

IV.2 Differential Equations via Finite Differences

Dr Sheehan Olver

Part IV

Applications of Linear Algebra

1. Polynomial Interpolation and Regression for approximating data
 2. Differential Equations via finite difference
- 

IV.2.1 Indefinite Integration

Solve the simplest ODE replacing derivatives w/ divided differences

Indefinite integration can be thought of as an ODE:


$$\begin{aligned} u(a) &= c, \\ u'(x) &= f(x) \quad \text{for} \quad a \leq x \leq b. \end{aligned}$$

Of course $u(x) = \int_a^x f(x) dx + c$, but let's ignore that since we want to generalise.

Idea: replace derivatives with divided differences.

Do so in 4 steps.

Step 1: ODE on interval \rightarrow ODE on grid



$$\begin{array}{l} u(a) = c, \\ \hline u'(x) = f(x) \end{array}$$

inf. dim
problem



$$\begin{bmatrix} u(x_0) \\ u'(x_0) \\ u'(x_1) \\ \vdots \\ u'(x_{n-1}) \end{bmatrix} = \begin{bmatrix} c \\ f(x_0) \\ f(x_1) \\ \vdots \\ f(x_{n-1}) \end{bmatrix}$$

$\underbrace{\hspace{10em}}_{\mathbf{b}}$

Stop at
 x_{n-1} .

finite dim,

Step 2: ODE on grid \rightarrow Divided differences on grid

\nearrow
righted-sided

$$u'(x) \approx \frac{u(x+h) - u(x)}{h} \Rightarrow$$

$$u'(x_j) \approx \frac{u(x_{j+1}) - u(x_j)}{h} = \frac{u(x_{j+1}) - u(x_j)}{h}$$

$$\begin{bmatrix} u(x_0) \\ u'(x_0) \\ u'(x_1) \\ \vdots \\ u'(x_{n-1}) \end{bmatrix} = \begin{bmatrix} c \\ f(x_0) \\ f(x_1) \\ \vdots \\ f(x_{n-1}) \end{bmatrix}$$

$\underbrace{\hspace{10em}}_{\mathbf{b}}$



$$\begin{bmatrix} u(x_0) \\ (u(x_1) - u(x_0))/h \\ (u(x_2) - u(x_1))/h \\ \vdots \\ (u(x_n) - u(x_{n-1}))/h \end{bmatrix} \approx \mathbf{b}$$

Step 3: Divided differences on grid \rightarrow Discrete system

New unknowns $\begin{bmatrix} u_0 \\ u_1 \\ \vdots \\ u_n \end{bmatrix}$ so that (hopefully)
 $u_j \approx u(x_j)$

$$\begin{bmatrix} u(x_0) \\ (u(x_1) - u(x_0))/h \\ (u(x_2) - u(x_1))/h \\ \vdots \\ (u(x_n) - u(x_{n-1}))/h \end{bmatrix} \approx \mathbf{b} \quad \rightarrow \quad \begin{bmatrix} u_0 \\ (u_1 - u_0)/h \\ (u_2 - u_1)/h \\ \vdots \\ (u_n - u_{n-1})/h \end{bmatrix} = \mathbf{b}$$

Step 4: Discrete system \rightarrow Linear system


$$\begin{bmatrix} u_0 \\ (u_1 - u_0)/h \\ (u_2 - u_1)/h \\ \vdots \\ (u_n - u_{n-1})/h \end{bmatrix} = \mathbf{b}$$



$$\underbrace{\begin{bmatrix} 1 & & & & \\ -1/h & 1/h & & & \\ & \ddots & \ddots & & \\ & & & -1/h & 1/h \end{bmatrix}}_L \underbrace{\begin{bmatrix} u_0 \\ u_1 \\ \vdots \\ u_n \end{bmatrix}}_{\mathbf{u}} = \mathbf{b}$$

$\left[\begin{matrix} c \\ f_0 \\ \vdots \\ f_{n-1} \end{matrix} \right]$

$$u(0) = 0, \quad u' = \cos x$$



Solve it using forward substitution in $O(n)$ ops

thereby find u ;

IV.2.1 Forward Euler


Generalise to first order linear ODEs

Consider general first order linear ODEs:

$$\begin{aligned} u(a) &= c \\ u'(x) - \omega(x)u(x) &= f(x) \end{aligned}$$

Repeat 4 steps as before.

Step 1: ODE on interval \rightarrow ODE on grid



$$\begin{aligned} &u(a) = c \\ &\underline{u'(x) - \omega(x)u(x) = f(x)} \end{aligned}$$

∞ -dim



$$\begin{bmatrix} \underline{u(x_0)} \\ u'(x_0) + \omega(x_0)u(x_0) \\ u'(x_1) + \omega(x_1)u(x_1) \\ \vdots \\ u'(x_{n-1}) + \omega(x_{n-1})u(x_{n-1}) \end{bmatrix} = \begin{bmatrix} \underline{c} \\ f(x_0) \\ f(x_1) \\ \vdots \\ \underbrace{f(x_{n-1})}_b \end{bmatrix}$$

finite dim

Step 2: ODE on grid \rightarrow Divided differences on grid

$$u'(x'_j) \approx \frac{u(x'_{j+1}) - u(x'_j)}{h}$$

$$\begin{bmatrix} u(x_0) \\ u'(x_0) + \omega(x_0)u(x_0) \\ u'(x_1) + \omega(x_1)u(x_1) \\ \vdots \\ u'(x_{n-1}) + \omega(x_{n-1})u(x_{n-1}) \end{bmatrix} = \underbrace{\begin{bmatrix} c \\ f(x_0) \\ f(x_1) \\ \vdots \\ f(x_{n-1}) \end{bmatrix}}_{\mathbf{b}} \quad \rightarrow \quad \begin{bmatrix} u(x_0) \\ (u(x_1) - u(x_0))/h + \omega(x_0)u(x_0) \\ (u(x_2) - u(x_1))/h + \omega(x_1)u(x_1) \\ \vdots \\ (u(x_n) - u(x_{n-1}))/h + \omega(x_{n-1})u(x_{n-1}) \end{bmatrix} \approx \mathbf{b}$$

right-sided
divided diff

Step 3: Divided differences on grid \rightarrow Discrete system

Now we write $u(x_j) \approx u_j$

$$\begin{bmatrix} u(x_0) \\ (u(x_1) - u(x_0))/h + \omega(x_0)u(x_0) \\ (u(x_2) - u(x_1))/h + \omega(x_1)u(x_1) \\ \vdots \\ (u(x_n) - u(x_{n-1}))/h + \omega(x_{n-1})u(x_{n-1}) \end{bmatrix} \approx \mathbf{b} \quad \rightarrow \quad \begin{bmatrix} u_0 \\ (u_1 - u_0)/h + \omega(x_0)u_0 \\ (u_2 - u_1)/h + \omega(x_1)u_1 \\ \vdots \\ (u_n - u_{n-1})/h + \omega(x_{n-1})u_{n-1} \end{bmatrix} = \mathbf{b}$$


Step 4: Discrete system \rightarrow Linear system

$$\begin{bmatrix} u_0 \\ (u_1 - u_0)/h + \omega(x_0)u_0 \\ (u_2 - u_1)/h + \omega(x_1)u_1 \\ \vdots \\ (u_n - u_{n-1})/h + \omega(x_{n-1})u_{n-1} \end{bmatrix} = \mathbf{b}$$



$$\underbrace{\begin{bmatrix} 1 & & & \\ \omega(x_0) - 1/h & 1/h & & \\ & \ddots & \ddots & \\ & & \omega(x_{n-1}) - 1/h & 1/h \end{bmatrix}}_L \underbrace{\begin{bmatrix} u_0 \\ u_1 \\ \vdots \\ u_n \end{bmatrix}}_{\mathbf{u}} = \mathbf{b}$$

$$u(0) = 1, u' + xu = e^x$$




Then solve via forward sub.

IV.2.3 Poisson

Use second order divided differences


Consider the simplest second order ODE with boundary conditions:



$$\begin{array}{l} u(0) = c, \\ \hline u''(x) = f(x), \\ \hline u(1) = d \end{array}$$

Repeat 4 steps as before but with second-order divided differences and one more boundary condition.

Step 1: ODE on interval \rightarrow ODE on grid



$$\begin{array}{c} u(0) = c, \\ \hline u''(x) = f(x), \\ \hline u(1) = d \end{array}$$



$$\begin{bmatrix} u(x_0) \\ u''(x_1) \\ u''(x_2) \\ \vdots \\ u''(x_{n-1}) \\ u(x_n) \end{bmatrix} = \underbrace{\begin{bmatrix} c \\ f(x_1) \\ f(x_2) \\ \vdots \\ f(x_{n-1}) \\ d \end{bmatrix}}_b$$

just use
interior
 x ;

Step 2: ODE on grid \rightarrow Divided differences on grid

2nd order divided diff:

$$u''(x_i) \approx \frac{u(x_{i-1}) - 2u(x_i) + u(x_{i+1}))}{h^2}$$

$$\begin{bmatrix} u(x_0) \\ u''(x_1) \\ u''(x_2) \\ \vdots \\ u''(x_{n-1}) \\ u(x_n) \end{bmatrix} = \underbrace{\begin{bmatrix} c \\ f(x_1) \\ f(x_2) \\ \vdots \\ f(x_{n-1}) \\ d \end{bmatrix}}_{\mathbf{b}}$$



$$\begin{bmatrix} u(x_0) \\ \frac{u(x_0) - 2u(x_1) + u(x_2)}{h^2} \\ \frac{u(x_1) - 2u(x_2) + u(x_3)}{h^2} \\ \vdots \\ \frac{u(x_{n-2}) - 2u(x_{n-1}) + u(x_n)}{h^2} \\ u(x_n) \end{bmatrix} \approx \mathbf{b}$$

Step 3: Divided differences on grid \rightarrow Discrete system

Again $u; \approx u(x;)$

$$\begin{bmatrix} u(x_0) \\ \frac{u(x_0) - 2u(x_1) + u(x_2)}{h^2} \\ \frac{u(x_1) - 2u(x_2) + u(x_3)}{h^2} \\ \vdots \\ \frac{u(x_{n-2}) - 2u(x_{n-1}) + u(x_n)}{h^2} \\ u(x_n) \end{bmatrix} \approx b$$



$$\begin{bmatrix} u_0 \\ \frac{u_0 - 2u_1 + u_2}{h^2} \\ \frac{u_1 - 2u_2 + u_3}{h^2} \\ \vdots \\ \frac{u_{n-2} - 2u_{n-1} + u_n}{h^2} \\ u_n \end{bmatrix} = b$$


Step 4: Discrete system \rightarrow Linear system

$$\begin{bmatrix} u_0 \\ \frac{u_0 - 2u_1 + u_2}{h^2} \\ \frac{u_1 - 2u_2 + u_3}{h^2} \\ \vdots \\ \frac{u_{n-2} - 2u_{n-1} + u_n}{h^2} \\ u_n \end{bmatrix} = \mathbf{b}$$



$$\underbrace{\begin{bmatrix} 1 & & & & \\ 1/h^2 & -2/h^2 & 1/h & & \\ & \ddots & \ddots & \ddots & \\ & & 1/h^2 & -2/h^2 & 1/h \\ & & & & 1 \end{bmatrix}}_A \underbrace{\begin{bmatrix} u_0 \\ u_1 \\ \vdots \\ u_n \end{bmatrix}}_{\mathbf{u}} = \mathbf{b}$$

$$\begin{aligned} u(0) &= \underbrace{1}_c \\ u''(x) &= \underbrace{-4x^2 \cos(x^2) - 2 \sin(x^2)}_{f(x)} \\ u(1) &= \underbrace{\cos 1}_d \end{aligned}$$



A is tridiagonal, can compute LU, QR

and solve in $O(n)$ operations,

~~Complex~~ Examinable Material! Week 1 - today