Week 4. Static games with complete information III: Nash equilibria

Exercise 1: Best responses

Consider the stag hunt game:

Suppose player 1 uses the mixed strategy (x, 1 - x), where x is player 1's probability to Stag. Similarly, player 2's strategy is (y, 1 - y).

- (i) For given x, y compute the players' payoffs $\pi^{(1)}(x, y), \pi^{(2)}(x, y)$ (see Remarks 2.6, 2.7).
- (ii) For a given y compute player 1's best response (BR(y)). In particular, show that there is some y^* such that all $x \in [0, 1]$ are a best response.
- (iii) Draw the two best response correspondences BR(x), BR(y) into a x-y plane. How often do they intersect? What does it mean if they intersect?

Exercise 2: Cournot Duopoly

The Cournot duopoly game is defined by:

- Players: $N = \{ \text{Firm } 1, \text{Firm } 2 \}$
- Actions: Amount of good produced, $x^{(i)} \in [0, \infty)$ for $i \in \{1, 2\}$
- Payoffs: $\pi^{(i)}(x^{(1)}, x^{(2)}) = [a b(x^{(1)} + x^{(2)})]x^{(i)} cx^{(i)}$

Show that there is a Nash equilibrium in pure strategies. For simplicity assume a=10,b=1,c=1.

[Hint: For each $x^{(i)}$ computer $BR(x^{(-i)})$. Then solve simultaneously:

$$x^{(1)} = BR(x^{(2)})$$

 $x^{(2)} = BR(x^{(1)})$

1

Exercise 3: Matching Pennies

Compute the Nash equilibria for the following two games, and interpret the result.

Bonus Exercise 1: Finding games with a non-generic number of equilibria

Find an example of a symmetric 2 player game, with 2 actions per player, with:

- Exactly 2 Nash equilibria
- infinitely many Nash equilibria

[Note: These should include all Nash equilibria. Not just pure Nash equilibria.]

Bonus Exercise 2: Verifying NE in games with finitely many players & actions

Show that to verify whether a strategy profile $\hat{\sigma} = (\hat{\sigma}^{(1)}, \dots, \hat{\sigma}^{(n)})$ is a Nash equilibrium, it is sufficient to check all deviations towards pure strategies.

Specifically show that $\hat{\sigma}$ is a Nash equilibrium if and only if for all players i the following two conditions hold:

- (i) All actions that player i uses give the same payoff: if $\sigma_j^{(i)} > 0$ and $\sigma_k^{(i)} > 0$ then $\pi^{(i)}(s_j^{(i)}, \hat{\sigma}^{(-i)}) = \pi^{(i)}(s_k^{(i)}, \hat{\sigma}^{(-i)})$.
- (ii) Actions that are not played are not profitable: if $\sigma_j^{(i)} = 0$ then $\pi^{(i)}(s_j^{(i)}, \hat{\sigma}^{(-i)}) \leq \pi^{(i)}(\hat{\sigma}^{(i)}, \hat{\sigma}^{(-i)})$.

[Hint: One way to prove the above is once again by contradiction.]