

Mathematical modeling

Jean-Louis Bouquard



Beijing Institute of Technology

Optimization Methods



- 1 Mathematical modeling
- 2 Solving linear problems

1 Mathematical modeling

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the constraints
- and the objective function
- Restriction! constraints and objective function must be LINEAR
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be LINEAR
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be LINEAR
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be **LINEAR**
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be **LINEAR**
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be **LINEAR**
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be **LINEAR**
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

Mathematical modeling

- Why a mathematical modeling?
- Define clearly the problem, the data
- Choose the **variables**
- Write the **constraints**
- and the **objective function**
- Restriction! constraints and objective function must be **LINEAR**
- Do not confuse variables (unknown, decisions) and data (known)
- Do not confuse math-model-variables and algorithm-variables

A first model for the F2// $\sum T_j$

- For every job j , $C_{1,j}$ and $C_{2,j}$ are the completion times of j
- T_j is the tardiness of j , $T_j \geq 0$

Succession of the operations and tardinesses:

$$\begin{aligned}\forall j, 1 \leq j \leq n, \quad & C_{1,j} \geq p_{1,j} \\ & C_{2,j} \geq C_{1,j} + p_{2,j} \\ & T_j \geq C_{2,j} - d_j\end{aligned}$$

Objective function:

$$\text{Minimize } \sum_{j=1}^n T_j$$

A first model for the F2// $\sum T_j$

- For every job j , $C_{1,j}$ and $C_{2,j}$ are the completion times of j
- T_j is the tardiness of j , $T_j \geq 0$

Succession of the operations and tardinesses:

$$\begin{aligned}\forall j, 1 \leq j \leq n, \quad & C_{1,j} \geq p_{1,j} \\ & C_{2,j} \geq C_{1,j} + p_{2,j} \\ & T_j \geq C_{2,j} - d_j\end{aligned}$$

Objective function:

$$\text{Minimize } \sum_{j=1}^n T_j$$

A first model for the F2// $\sum T_j$

- For every job j , $C_{1,j}$ and $C_{2,j}$ are the completion times of j
- T_j is the tardiness of j , $T_j \geq 0$

Succession of the operations and tardinesses:

$$\begin{aligned}\forall j, 1 \leq j \leq n, \quad & C_{1,j} \geq p_{1,j} \\ & C_{2,j} \geq C_{1,j} + p_{2,j} \\ & T_j \geq C_{2,j} - d_j\end{aligned}$$

Objective function:

$$\text{Minimize } \sum_{j=1}^n T_j$$

A first model for the F2// $\sum T_j$

- For every job j , $C_{1,j}$ and $C_{2,j}$ are the completion times of j
- T_j is the tardiness of j , $T_j \geq 0$

Succession of the operations and tardinesses:

$$\begin{aligned}\forall j, 1 \leq j \leq n, \quad & C_{1,j} \geq p_{1,j} \\ & C_{2,j} \geq C_{1,j} + p_{2,j} \\ & T_j \geq C_{2,j} - d_j\end{aligned}$$

Objective function:

$$\text{Minimize } \sum_{j=1}^n T_j$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0,1\}$)
 indicating whether j_1 is before j_2
- $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
 - Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
 - Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
 - Idea: link this choice to a binary variable
 - For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0,1\}$)
 indicating whether j_1 is before j_2
 - $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0,1\}$)
 indicating whether j_1 is before j_2
- $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0, 1\}$)
 indicating whether j_1 is before j_2
- $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0, 1\}$)
 indicating whether j_1 is before j_2
- $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0, 1\}$)
 indicating whether j_1 is before j_2

$$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

- How to express the *non overlap* constraints?
- For every 2 jobs (j_1, j_2) and every machine i
- Either j_1 is before j_2 and $C_{i,j_1} \leq C_{i,j_2} - p_{i,j_2}$
- Or j_1 is after j_2 and $C_{i,j_2} \leq C_{i,j_1} - p_{i,j_1}$
- Idea: link this choice to a binary variable
- For every 2 jobs j_1 and j_2 ($j_1 < j_2$)
 Z_{j_1,j_2} is a binary variable ($\{0, 1\}$)
 indicating whether j_1 is before j_2
- $$Z_{j_1,j_2} = \begin{cases} 0 & \text{if } j_1 \text{ before } j_2 \\ 1 & \text{if } j_1 \text{ after } j_2 \end{cases}$$

A first model for the F2// ΣT_j

Non overlap constraints:

$$\forall (j_1, j_2), 1 \leq j_1 < j_2 \leq n, \quad C_{1,j_1} \leq C_{1,j_2} - p_{1,j_2} \quad \text{if } Z_{j_1,j_2} = 0$$

$$C_{1,j_2} \leq C_{1,j_1} - p_{1,j_1} \quad \text{if } Z_{j_1,j_2} = 1$$

They can be turned into:

$$\forall i \in \{1, 2\}, \forall (j_1, j_2), \quad C_{i,j_1} + p_{i,j_2} \leq C_{i,j_2} + HV \times Z_{j_1,j_2}$$

$$1 \leq j_1 < j_2 \leq n, \quad C_{i,j_1} + p_{i,j_2} \leq C_{i,j_1} + HV \times (1 - Z_{j_1,j_2})$$

Where HV is a *high value* bigger than any $C_{i,j}$

A first model for the F2// ΣT_j

Non overlap constraints:

$$\begin{aligned} \forall (j_1, j_2), 1 \leq j_1 < j_2 \leq n, \quad & C_{1,j_1} \leq C_{1,j_2} - p_{1,j_2} \text{ if } Z_{j_1,j_2} = 0 \\ & C_{1,j_2} \leq C_{1,j_1} - p_{1,j_1} \text{ if } Z_{j_1,j_2} = 1 \end{aligned}$$

They can be turned into:

$$\begin{aligned} \forall i \in \{1, 2\}, \forall (j_1, j_2), \quad & C_{i,j_1} + p_{i,j_2} \leq C_{i,j_2} + HV \times Z_{j_1,j_2} \\ 1 \leq j_1 < j_2 \leq n, \quad & C_{i,j_1} + p_{i,j_2} \leq C_{i,j_1} + HV \times (1 - Z_{j_1,j_2}) \end{aligned}$$

Where HV is a *high value* bigger than any $C_{i,j}$

Example of a permutation

The permutation (253614) is represented by the (triangular) matrix (Z_{j_1, j_2}) :

	1	2	3	4	5	6
1		1	1	0	1	1
2			0	0	0	0
3				0	1	0
4					1	1
5						0
6						

Example of a permutation

The permutation (253614) is represented by the (triangular) matrix (Z_{j_1, j_2}) :

	1	2	3	4	5	6
1		1	1	0	1	1
2			0	0	0	0
3				0	1	0
4					1	1
5						0
6						

A first model for the F2// ΣT_j

Exclusion on each machine:

$$\forall i \in \{1, 2\}, \forall (j_1, j_2), 1 \leq j_1 < j_2 \leq n, \quad (1)$$

$$p_{i,j_2} \leq C_{i,j_2} - C_{i,j_1} + HV \times Z_{j_1,j_2} \leq HV - p_{i,j_1}$$

Succession of the operations and tardinesses:

$$\forall j, 1 \leq j \leq n, \quad C_{1,j} \geq p_{1,j} \quad (2)$$

$$C_{2,j} \geq C_{1,j} + p_{2,j} \quad (3)$$

$$T_j \geq C_{2,j} - d_j \quad (4)$$

Objective function

$$\text{Minimize} \quad \sum_{j=1}^n T_j \quad (5)$$

A first model for the F2// ΣT_j

Exclusion on each machine:

$$\forall i \in \{1, 2\}, \forall (j_1, j_2), 1 \leq j_1 < j_2 \leq n, \quad (1)$$

$$p_{i,j_2} \leq C_{i,j_2} - C_{i,j_1} + HV \times Z_{j_1,j_2} \leq HV - p_{i,j_1}$$

Succession of the operations and tardinesses:

$$\forall j, 1 \leq j \leq n, \quad C_{1,j} \geq p_{1,j} \quad (2)$$

$$C_{2,j} \geq C_{1,j} + p_{2,j} \quad (3)$$

$$T_j \geq C_{2,j} - d_j \quad (4)$$

Objective function

$$\text{Minimize} \quad \sum_{j=1}^n T_j \quad (5)$$

A first model for the F2// ΣT_j

Exclusion on each machine:

$$\begin{aligned} \forall i \in \{1, 2\}, \forall (j_1, j_2), 1 \leq j_1 < j_2 \leq n, \\ p_{i,j_2} \leq C_{i,j_2} - C_{i,j_1} + HV \times Z_{j_1,j_2} \leq HV - p_{i,j_1} \end{aligned} \quad (1)$$

Succession of the operations and tardinesses:

$$\forall j, 1 \leq j \leq n, \quad C_{1,j} \geq p_{1,j} \quad (2)$$

$$C_{2,j} \geq C_{1,j} + p_{2,j} \quad (3)$$

$$T_j \geq C_{2,j} - d_j \quad (4)$$

Objective function

$$\text{Minimize} \quad \sum_{j=1}^n T_j \quad (5)$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not a linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: "Is $j = S_k$?"
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: "Is $j = S_k$?"
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
 - It is **not** a **linear** constraint
 - p_{1,S_k} is depending on S_k but not linearly
 - Again, we use binary variables:
 - To the yes-no question: "Is $j = S_k$?"
 - we associate the binary variable $X_{k,j}$
 - $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2//\sum T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
 - p_{1,S_k} is depending on S_k but not linearly
 - Again, we use binary variables:
 - To the yes-no question: "Is $j = S_k$?"
 - we associate the binary variable $X_{k,j}$
 - $X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: "Is $j = S_k$?"
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
 - To the yes-no question: "Is $j = S_k$?"
 - we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: “Is $j = S_k$?”
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2//\sum T_j$

- The main decision is the choice of the permutation
- Can we choose $S_k =$ number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: “Is $j = S_k$?”
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model for the $F2/\Sigma T_j$

- The main decision is the choice of the permutation
- Can we choose S_k = number of the job in position k as a variable ?
- We would have: $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$
- It is **not** a **linear** constraint
- p_{1,S_k} is depending on S_k but not linearly
- Again, we use binary variables:
- To the yes-no question: “Is $j = S_k$?”
- we associate the binary variable $X_{k,j}$
- $$X_{k,j} = \begin{cases} 1 & \text{if } j \text{ has rank } k \\ 0 & \text{otherwise} \end{cases}$$

A second model

Constraints

- Every rank is assigned to some job:

$$\forall k = 1..n \quad \sum_{j=1}^n X_{k,j} = 1 \quad (6)$$

- Every job has some rank:

$$\forall j = 1..n \quad \sum_{k=1}^n X_{k,j} = 1 \quad (7)$$

A second model

Constraints

- Every rank is assigned to some job:

$$\forall k = 1..n \quad \sum_{j=1}^n X_{k,j} = 1 \quad (6)$$

- Every job has some rank:

$$\forall j = 1..n \quad \sum_{k=1}^n X_{k,j} = 1 \quad (7)$$

Example of a permutation

The permutation (253614) is represented by the matrix $(X_{k,j})$:

	1	2	3	4	5	6
1	0	1	0	0	0	0
2	0	0	0	0	1	0
3	0	0	1	0	0	0
4	0	0	0	0	0	1
5	1	0	0	0	0	0
6	0	0	0	1	0	0

Example of a permutation

The permutation (253614) is represented by the matrix $(X_{k,j})$:

	1	2	3	4	5	6
1	0	1	0	0	0	0
2	0	0	0	0	1	0
3	0	0	1	0	0	0
4	0	0	0	0	0	1
5	1	0	0	0	0	0
6	0	0	0	1	0	0

A second model

- In $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$
- we will replace C_{1,s_k} with $F_{1,k}$
- For every rank k , $F_{1,k}$ and $F_{2,k}$ are the completion times of the k^{th} job
- Similarly T_{s_k} is renamed G_k
- G_k is the tardiness of the k^{th} job, $G_k \geq 0$

A second model

- In $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$
- we will replace C_{1,s_k} with $F_{1,k}$
- For every rank k , $F_{1,k}$ and $F_{2,k}$ are the completion times of the k^{th} job
- Similarly T_{S_k} is renamed G_k
- G_k is the tardiness of the k^{th} job, $G_k \geq 0$

A second model

- In $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$
- we will replace C_{1,s_k} with $F_{1,k}$
- For every rank k , $F_{1,k}$ and $F_{2,k}$ are the completion times of the k^{th} job
- Similarly T_{S_k} is renamed G_k
- G_k is the tardiness of the k^{th} job, $G_k \geq 0$

A second model

- In $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$
- we will replace C_{1,s_k} with $F_{1,k}$
- For every rank k , $F_{1,k}$ and $F_{2,k}$ are the completion times of the k^{th} job
- Similarly T_{S_k} is renamed G_k
- G_k is the tardiness of the k^{th} job, $G_k \geq 0$

A second model

- In $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$
- we will replace C_{1,s_k} with $F_{1,k}$
- For every rank k , $F_{1,k}$ and $F_{2,k}$ are the completion times of the k^{th} job
- Similarly T_{S_k} is renamed G_k
- G_k is the tardiness of the k^{th} job, $G_k \geq 0$

A second model

- $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,S_k}$
- We need p_{1,S_k} (and p_{2,S_k} and d_{S_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,S_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,S_k}$
- We need p_{1,S_k} (and p_{2,S_k} and d_{S_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,S_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,s_k}$
- We need p_{1,s_k} (and p_{2,s_k} and d_{s_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,s_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,s_k}$
- We need p_{1,s_k} (and p_{2,s_k} and d_{s_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,s_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,s_k}$
- We need p_{1,s_k} (and p_{2,s_k} and d_{s_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,s_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,S_k} \geq C_{1,S_{k-1}} + p_{1,S_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,S_k}$
- We need p_{1,S_k} (and p_{2,S_k} and d_{S_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,S_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,s_k}$
- We need p_{1,s_k} (and p_{2,s_k} and d_{s_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,s_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

- $C_{1,s_k} \geq C_{1,s_{k-1}} + p_{1,s_k}$ is written
- $F_{1,k} \geq F_{1,k-1} + p_{1,s_k}$
- We need p_{1,s_k} (and p_{2,s_k} and d_{s_k})
- S_k is replaced by the row $(X_{k,1}, X_{k,2}, \dots, X_{k,n})$
- $(X_{k,1}, X_{k,2}, \dots, X_{k,n}) = (0, \dots, 0, 1, 0, \dots, 0)$
- The 1 has the S_k^{th} position
- Thus p_{1,s_k} is $\sum_{j=1}^n p_{1,j} X_{k,j}$
- $\sum_{j=1}^n p_{1,j} X_{k,j}$ is linear

A second model

Constraints

The first job:

$$F_{1,1} = \sum_{j=1}^n p_{1,j} X_{1,j} \quad (8)$$

$$F_{2,1} = \sum_{j=1}^n (p_{1,j} + p_{2,j}) X_{1,j} \quad (9)$$

The other jobs on machine 1:

$$\forall k = 2..n \quad F_{1,k} = F_{1,k-1} + \sum_{j=1}^n p_{1,j} X_{k,j} \quad (10)$$

The other jobs on machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{2,k-1} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (11)$$

A second model

Constraints

The first job:

$$F_{1,1} = \sum_{j=1}^n p_{1,j} X_{1,j} \quad (8)$$

$$F_{2,1} = \sum_{j=1}^n (p_{1,j} + p_{2,j}) X_{1,j} \quad (9)$$

The other jobs on machine 1:

$$\forall k = 2..n \quad F_{1,k} = F_{1,k-1} + \sum_{j=1}^n p_{1,j} X_{k,j} \quad (10)$$

The other jobs on machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{2,k-1} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (11)$$

A second model

Constraints

The first job:

$$F_{1,1} = \sum_{j=1}^n p_{1,j} X_{1,j} \quad (8)$$

$$F_{2,1} = \sum_{j=1}^n (p_{1,j} + p_{2,j}) X_{1,j} \quad (9)$$

The other jobs on machine 1:

$$\forall k = 2..n \quad F_{1,k} = F_{1,k-1} + \sum_{j=1}^n p_{1,j} X_{k,j} \quad (10)$$

The other jobs on machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{2,k-1} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (11)$$

A second model

The other jobs from machine 1 to machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{1,k} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (12)$$

The tardinesses:

$$\forall k = 1..n \quad G_k \geq F_{2,k} - \sum_{j=1}^n d_j X_{k,j} \quad (13)$$

The criterion:

$$\text{Minimize} \quad \sum_{k=1}^n G_k \quad (14)$$

A second model

The other jobs from machine 1 to machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{1,k} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (12)$$

The tardinesses:

$$\forall k = 1..n \quad G_k \geq F_{2,k} - \sum_{j=1}^n d_j X_{k,j} \quad (13)$$

The criterion:

$$\text{Minimize} \quad \sum_{k=1}^n G_k \quad (14)$$

A second model

The other jobs from machine 1 to machine 2:

$$\forall k = 2..n \quad F_{2,k} \geq F_{1,k} + \sum_{j=1}^n p_{2,j} X_{k,j} \quad (12)$$

The tardinesses:

$$\forall k = 1..n \quad G_k \geq F_{2,k} - \sum_{j=1}^n d_j X_{k,j} \quad (13)$$

The criterion:

$$\text{Minimize} \quad \sum_{k=1}^n G_k \quad (14)$$

2 Solving linear problems

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{i,j} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_j$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
- is a linear problem if all the X_j 's are in \mathbb{R}
- It is said a mixed integer linear problem if some X_j 's are in \mathbb{Z}

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{ij} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_j$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
- is a linear problem if all the X_j 's are in \mathbb{R}
- It is said a mixed integer linear problem if some X_j 's are in \mathbb{Z}

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{i,j} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_i$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
- is a **linear problem** if all the X_j 's are in \mathbb{R}
- It is said a mixed integer linear problem if some X_j 's are in \mathbb{Z}

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{i,j} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_i$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
 - is a **linear problem** if all the X_j 's are in \mathbb{R}
 - It is said a **mixed integer linear problem** if some X_j 's are in \mathbb{Z}

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{i,j} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_i$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
- is a **linear problem** if all the X_j 's are in \mathbb{R}
- It is said a **mixed integer linear problem** if some X_j 's are in \mathbb{Z}

Linear problems

- The problem defined by:
- Variables are $(X_1, \dots, X_n) = X$
- Constraints are $\sum_{j=1}^n a_{i,j} X_j \begin{cases} \leq \\ = \\ \geq \end{cases} b_i$
- Minimize (or Maximize) $f(X) = \sum_{j=1}^n c_j X_j$
- is a **linear problem** if all the X_j 's are in \mathbb{R}
- It is said a **mixed integer linear problem** if some X_j 's are in \mathbb{Z}

Solving linear problems

- For the LP problems (continuous variables)
 - proved polynomial (Khachiyan 1979)
 - very efficient methods:
 - Simplex method (Dantzig, 1947)
 - Interior-point method (Karmarcar, 1984)

Solving linear problems

- For the LP problems (continuous variables)
- proved polynomial (Khachiyan 1979)
- very efficient methods:
- Simplex method (Dantzig, 1947)
- Interior-point method (Karmarcar, 1984)

Solving linear problems

- For the LP problems (continuous variables)
- proved polynomial (Khachiyan 1979)
- very efficient methods:
 - Simplex method (Dantzig, 1947)
 - Interior-point method (Karmarcar, 1984)

Solving linear problems

- For the LP problems (continuous variables)
- proved polynomial (Khachiyan 1979)
- very efficient methods:
- Simplex method (Dantzig, 1947)
- Interior-point method (Karmarcar, 1984)

Solving linear problems

- For the LP problems (continuous variables)
- proved polynomial (Khachiyan 1979)
- very efficient methods:
- Simplex method (Dantzig, 1947)
- Interior-point method (Karmarcar, 1984)

Solving linear problems

- For the ILP problems (integer variables)
- NP-hard
- Branch & bound, branch & cut, branch & price
- ...

Solving linear problems

- For the ILP problems (integer variables)
- NP-hard
- Branch & bound, branch & cut, branch & price
- . . .

Solving linear problems

- For the ILP problems (integer variables)
- NP-hard
- Branch & bound, branch & cut, branch & price
- . . .

Solving linear problems

- For the ILP problems (integer variables)
- NP-hard
- Branch & bound, branch & cut, branch & price
- . . .

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- ...
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- ...
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- ...
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- ...
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- . . .
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- Solvers
- GLPK
- CPLEX
- MATLAB
- ...
- Spreadsheets: OpenOffice, Excel

Solving linear problems

- This provides an exact method
- therefore only for small instances
- otherwise provides an approximation method with a time limit

Solving linear problems

- This provides an exact method
- therefore only for small instances
- otherwise provides an approximation method with a time limit

Solving linear problems

- This provides an exact method
- therefore only for small instances
- otherwise provides an approximation method with a time limit

The solver GLPK

- Models are **written** in the **language** GNU MathProg
- **Data** as well
- Then run GLPSOL (model1 or model2) to get an **optimal solution**
- otherwise, with a time limit, get an suboptimal solution
- or with the **relaxed** LP problem, get a lower bound

The solver GLPK

- Models are **written** in the **language** GNU MathProg
- **Data** as well
- Then run GLPSOL (model1 or model2) to get an **optimal solution**
- otherwise, with a time limit, get an **suboptimal solution**
- or with the **relaxed LP** problem, get a **lower bound**

The solver GLPK

- Models are **written** in the **language** GNU MathProg
- **Data** as well
- Then run GLPSOL (**model1** or **model2**) to get an **optimal solution**
- otherwise, with a time limit, get an **suboptimal solution**
- or with the **relaxed LP** problem, get a **lower bound**

The solver GLPK

- Models are **written** in the **language** GNU MathProg
- **Data** as well
- Then run GLPSOL (**model1** or **model2**) to get an **optimal solution**
- otherwise, with a time limit, get an **suboptimal solution**
- or with the relaxed LP problem, get a **lower bound**

The solver GLPK

- Models are **written** in the **language** GNU MathProg
- **Data** as well
- Then run GLPSOL (**model1** or **model2**) to get an **optimal solution**
- otherwise, with a time limit, get an **suboptimal solution**
- or with the **relaxed** LP problem, get a **lower bound**

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to Python and its PULP module
- Similarly to C++ and its CPLEX library
- Beside, to OpenOffice

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to **Python** and its **PULP** module
- Similarly to C++ and its CPLEX library
- Beside, to OpenOffice

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to Python and its PULP module
- Similarly to C++ and its CPLEX library
- Beside, to OpenOffice

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to **Python** and its **PULP** module
 - Similarly to C++ and its CPLEX library
 - Beside, to OpenOffice

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to **Python** and its **PULP** module
- Similarly to **C++** and its CPLEX library
- Beside, to OpenOffice

Solving linear problems

Using GLPK

- Solvers can also be handled with a programming language
- thanks to a library
- C++, C, Python, Java, ...
- A glance to **Python** and its **PULP** module
- Similarly to **C++** and its CPLEX library
- Beside, to **OpenOffice**