



School of Mathematical Sciences
Dalian University of Technology

Prof. Xian Liao

Lingong Road 2
116024 Dalian, P.R.China

E-Mail: liao@dlut.edu.cn

Website: <https://faculty.dlut.edu.cn/liao>

Lecture Notes for “Fourier Analysis and Differential Equations”^{1 2}

Time³ & Place:

Tuesday, 13:30-15:05, 研教楼 306

(Sep. 8, Sep. 15, Sep. 22, Sep. 29, Oct. 13, Oct. 20, Oct. 27, Nov. 3)

Wednesday, 10:05-11:40, 综合教学 1 号楼 356

(Sep. 9, Sep. 16, Sep. 23, Sep. 30, Oct. 14, Oct. 21, Oct. 28, Nov. 4)

There are NO lectures at Sep. 20 & Oct. 10!

Teaching assistant:

Dingli Jiang ([dinglijiang1018\(at\)foxmail.com](mailto:dinglijiang1018@foxmail.com))

¹The lecture notes are based on a lecture given at the Dalian University of Technology during the fall term 2026.

²Comments are welcome to be sent to me by email.

³It takes place from Sep. 6 to Nov. 6, 2026 (2.-10. semester week).

Contents

1	Fourier series	3
1.1	Examples of boundary value problems	3
1.1.1	An ODE with Dirichlet boundary condition	3
1.1.2	Fourier's idea to solve the heat equation	4
1.2	Spectral theorem and Fourier series	6
1.2.1	Resolvent computation	7
1.2.2	Spectral theorem and Fourier sine series	9
1.2.3	Eigenvalue problem with Neumann boundary condition	10
1.2.4	Full Fourier series	11
1.3	Pointwise convergence of Fourier series	13
1.3.1	Dirichlet kernel	13
1.3.2	Dini's criterion and du Bois-Reymond's counterexample	15
1.3.3	Dirichlet-Jordan's criterion	17
1.3.4	Fejér's kernel	19
1.4	Convergence in norm of Fourier series	20
1.4.1	Conjugate function and Riesz projections	22
1.4.2	Harmonic extension and Poisson kernel	24
1.4.3	Harmonic conjugate and conjugate Poisson kernel . . .	26
1.4.4	Other related topics	28

This two-month lecture provides an introduction to the sophisticated theory of Fourier analysis. The one-dimensional case is particularly emphasized, serving as motivation and providing examples for the illustration of the general theory. It also helps to connect the various concepts encountered in the theories of ordinary differential equations (ODEs), functional analysis, complex analysis, harmonic analysis and many other fields. Due to time constraints and the ambitious goal, we are unable to define all the related mathematical concepts rigorously, and instead aim to draw the audience's interest to the relevant theory. (The footnotes are left as exercises to the audience.) Finally the application to partial differential equations (PDEs) is demonstrated.

1 Fourier series

1.1 Examples of boundary value problems

1.1.1 An ODE with Dirichlet boundary condition

Consider the boundary value problem for a second-order ordinary differential equation on the interval $[0, \pi]$,

$$u''(x) + \lambda u(x) = 0 \text{ on } (0, \pi), \quad u(0) = u(\pi) = 0. \quad (1.1)$$

We consider three different cases:

- Case $\lambda < 0$.

The following two functions are the fundamental solutions for the ODE $u''(x) = -\lambda u(x)$

$$\begin{aligned} \sinh(\sqrt{-\lambda}x) &= \frac{e^{\sqrt{-\lambda}x} - e^{-\sqrt{-\lambda}x}}{2}, \\ \cosh(\sqrt{-\lambda}x) &= \frac{e^{\sqrt{-\lambda}x} + e^{-\sqrt{-\lambda}x}}{2}, \end{aligned}$$

The boundary conditions imply that the solution should be zero. ⁴

- Case $\lambda = 0$.

A regular solution of $u''(x) = 0$ is a linear function $u(x) = ax + b$, which is 0 due to the boundary conditions.

⁴One can also argue from another perspective: We take the $L^2(0, \pi)$ -inner product

- Case $\lambda > 0$.

The following two functions are fundamental solutions for the ODE $u''(x) = -\lambda u(x)$

$$\begin{aligned}\sin(\sqrt{\lambda}x) &= \frac{e^{i\sqrt{\lambda}x} - e^{-i\sqrt{\lambda}x}}{2i}, \\ \cos(\sqrt{\lambda}x) &= \frac{e^{i\sqrt{\lambda}x} + e^{-i\sqrt{\lambda}x}}{2},\end{aligned}$$

The boundary conditions imply that the solution should be a linear combination of

$$\sin(nx), \quad n = 1, 2, \dots$$

To conclude, there is an infinite set of discrete values of λ :

$$\lambda = n^2, \quad n = 1, 2, \dots$$

such that the boundary value problem (1.1) has a nontrivial solution. These values n^2 , $n \in \mathbb{N}$ are called the eigenvalues of the (eigenvalue) problem (1.1) and the functions

$$u_n(x) = b_n \sin(nx), \quad n \in \mathbb{N}, \quad b_n \in \mathbb{R}(\text{or } \mathbb{C})$$

are the corresponding eigenfunctions. The concept of eigenvalues and eigenfunctions becomes clearer in Section 1.2 below.

1.1.2 Fourier's idea to solve the heat equation

Consider the one-dimensional initial-boundary value problem for the (real-valued) heat equation

$$\begin{cases} \partial_t u - \partial_{xx} u = 0 & \text{on } (0, \infty) \times (0, \pi), \\ u(0, x) = f(x) & x \in (0, \pi), \\ u(t, 0) = u(t, \pi) = 0 & t \in (0, \infty), \end{cases} \quad (1.3)$$

between the equation and the solution u itself, to obtain by integration by parts

$$-\int_0^\pi |u'|^2 dx + \lambda \int_0^\pi |u|^2 dx = 0. \quad (1.2)$$

This implies, if u is a regular solution (e.g. in $H^1(0, \pi)$), then $u = 0$. This argument can be easily generalized to the case when $\lambda \in \mathbb{C}$ with $\text{Im } \lambda \neq 0$, where $u = 0$ (in $H^1(0, \pi)$) follows.

where the compatibility condition is satisfied

$$f(0) = f(\pi) = 0,$$

and we assume that f is not identically zero: $f \neq 0$.

J. Fourier⁵ made the following Ansatz (separation of variables)

$$u(t, x) = T(t)X(x).$$

This leads to the following:

- $T'(t)X(x) - T(t)X''(x) = 0$, which implies (at least formally),

$$\frac{T'(t)}{T(t)} = \frac{X''(x)}{X(x)} = -\lambda \text{ for some constant } \lambda.$$

- $T(t)X(0) = T(t)X(\pi) = 0$ for all time t , which implies

$$X(0) = X(\pi) = 0.$$

- $T(0)X(x) = f(x)$.

From the ODE results above, if $f \neq 0$ and one searches for nontrivial (real-valued) solution, λ can “only” take the value n^2 for some $n \in \mathbb{N}$, and $X(x)$ is a multiple of $\sin(nx)$, and in this case, the time function $T(t) = T(0)e^{-n^2t}$. That is, if the initial data is

$$f(x) = b_n \sin(nx),$$

for some $n \in \mathbb{N}$ and some constant b_n , then the problem (1.3) has a (unique) solution

$$u(t, x) = T(t)X(x) = T(0)e^{-n^2t} \frac{b_n \sin(nx)}{T(0)} = b_n e^{-n^2t} \sin(nx).$$

If the initial data is a finite sum of sine functions

$$f(x) = \sum_{n=1}^N b_n \sin(nx),$$

then by the superposition principle the solution is

$$u(t, x) = \sum_{n=1}^N b_n e^{-n^2t} \sin(nx).$$

⁵Joseph Fourier, *Théorie Analytique de la Chaleur*. Chez F. Didot, 1822 (Firstly presented to *Institute de France* in 1807).

Thus, formally, if we can expand the initial data $f(x)$ as a sine series

$$f(x) = \sum_{n=1}^{\infty} b_n \sin(nx), \quad (1.4)$$

then the following function (when it is well-defined) is a solution of (1.1)

$$u(x) = \sum_{n=1}^{\infty} b_n e^{-n^2 t} \sin(nx).$$

A function with such an expansion (1.4) (nowadays called Fourier sine series) “seems” very special. For half a century around 1800 there was a great debate whether such a series expansion holds only for specific functions.

1.2 Spectral theorem and Fourier series

We claim that any function in the functional space

$$\{f \in C^2([0, \pi]) \mid f(0) = f(\pi) = 0\} \quad (1.5)$$

can be expanded as Fourier sine series. Historically one can make use of the Weierstrass approximation theorem and the density of trigonometric functions in the space of polynomial functions to prove this statement. Here we illustrate this in modern mathematical language.

We can reformulate (1.1) as the following eigenvalue problem

$$\mathcal{A}u = \lambda u, \quad (1.6)$$

where the operator $\mathcal{A}$ is defined by⁶

$$\mathcal{A} : D(\mathcal{A}) \rightarrow X, \quad \mathcal{A}(u) = -u'' \text{ for } u \in D(\mathcal{A}). \quad (1.7)$$

Here we take the underlying inner product space and scalar product below

$$X = C([0, \pi]; \mathbb{C}), \quad \langle u, v \rangle = \int_0^\pi u \bar{v} \, dx,$$

and the domain of the operator $\mathcal{A}$ is a subspace of X :

$$D(\mathcal{A}) = \{u \in C^2([0, \pi]) \mid u(0) = u(\pi) = 0\} \subset X.$$

⁶There is a certain conceptual mismatch between the classical setting of continuous functions and the spectral-theoretic framework, which is more naturally aligned with spaces such as $L^2(0, \pi)$. For example, one may define $\mathcal{A} : H_0^2(0, \pi) = D(\mathcal{A}) \rightarrow L^2(0, \pi)$.

The induced norm on X is

$$\|u\| = \langle u, u \rangle^{\frac{1}{2}},$$

and the distance between u and v is $\|u - v\|$. Note that $D(\mathcal{A})$ and X are not complete with respect to the defined metric.

Recall that the Lebesgue space

$$L^2(0, \pi) = \{f \text{ is Lebesgue measurable on } (0, \pi) \mid \|f\|_{L^2(0, \pi)} = \left(\int_0^\pi |f(x)|^2 dx \right)^{\frac{1}{2}} < \infty\} / \sim,$$

where

- The integral is understood in the sense of Riemann-Lebesgue:

$$\|f\|_{L^2(0, \pi)} = \left(2 \int_0^\infty \alpha \, m(\{x \in (0, \pi) \mid |f(x)| > \alpha\}) d\alpha \right)^{\frac{1}{2}}$$

- The equivalence $\sim$ means identity almost everywhere (a.e.):

$$f \sim g \Leftrightarrow m(\{x \in (0, \pi) \mid f(x) \neq g(x)\}) = 0.$$

The Lebesgue space $L^2(0, \pi)$ is endowed with the scalar product $\langle \cdot, \cdot \rangle$, and is a Hilbert space. By the density of the space⁷ of test functions $C_c^\infty(0, \pi)$ in $L^2(0, \pi)$, the closure of $C_c^\infty(0, \pi) \subset D(A) \subset X$ in the Lebesgue space $(L^2(0, \pi), \langle \cdot, \cdot \rangle)$ is the Lebesgue space itself

$$\overline{X}^{L^2} = \overline{D(\mathcal{A})}^{L^2} = \overline{C_c^\infty(0, \pi)}^{L^2} = L^2(0, \pi).$$

1.2.1 Resolvent computation

From the ODE results above (recalling (1.2)), the eigenvalues and normalized eigenfunctions of the operator $\mathcal{A}$ are

$$\lambda_n = n^2, \quad u_n = \sqrt{\frac{2}{\pi}} \sin(nx), \quad n \in \mathbb{N}, \quad (1.8)$$

with $\|u_n\|_{L^2(0, \pi)} = 1$.

[Sep 8, 2026]
[Sep 9, 2026]

⁷The density result also holds on the whole line or in higher dimensions.

We now compute the resolvent of $\mathcal{A}$ at $\lambda = -1 \notin \mathbb{N}$, that is, we compute $R_{\mathcal{A}}(-1) = (\mathcal{A} + 1)^{-1} : X \rightarrow D(\mathcal{A})$. That is, for a given function $g \in X$, we seek the solution $u \in D(\mathcal{A})$ of the following boundary value problem for the inhomogeneous ODE:

$$-u'' + u = g, \quad u(0) = u(\pi) = 0. \quad (1.9)$$

It suffices to compute the Green function of this boundary value problem. The homogeneous equation $-u'' + u = 0$ has fundamental solutions

$$e^{-x}, \quad e^x.$$

The solution satisfying the boundary condition $u(0) = 0$ is a multiple of

$$\phi(x) = e^x - e^{-x},$$

while the solution satisfying the boundary condition $u(\pi) = 0$ is a multiple of

$$\psi(x) = e^{-(\pi-x)} - e^{\pi-x}.$$

Their Wronskian $W(\phi, \psi) = \phi\psi' - \phi'\psi = \det \begin{pmatrix} \phi & \psi \\ \phi' & \psi' \end{pmatrix}$ is independent of x ⁸ and is given by

$$W(\phi, \psi) = 2(e^\pi - e^{-\pi}).$$

The Green function⁹ is

$$\begin{aligned} G(x, y) &= \begin{cases} -\frac{\phi(x)\psi(y)}{W(\phi, \psi)} & \text{for } x < y \\ -\frac{\phi(y)\psi(x)}{W(\phi, \psi)} & \text{for } x > y \end{cases} \\ &= -\frac{e^{-\pi+x+y} + e^{\pi-(x+y)} - e^{\pi-|x-y|} - e^{-\pi+|x-y|}}{W(\phi, \psi)}. \end{aligned}$$

Thus the unique solution of (1.9) reads¹⁰

$$u(x) = \int_0^\pi G(x, y)g(y) \, dy, \quad (1.10)$$

⁸It follows from Abel's identity (also called Liouville's formula).

⁹The $-$ sign before the quantity $\frac{\phi(x)\psi(y)}{W(\phi, \psi)}$ or $\frac{\phi(y)\psi(x)}{W(\phi, \psi)}$ in the definition of the Green function here is due to the $-$ sign before the highest order derivative ∂_{xx} in the equation (1.9), which (usually) is assumed to be $+$ in the standard formulation of ODE.

¹⁰The unique solvability of the inhomogeneous problem (1.9) follows from the fact that the homogeneous problem (1.1) with $\lambda = -1$ has only trivial zero solution, by the Fredholm

that is,

$$R_{\mathcal{A}}(-1)g(x) = (\mathcal{A} + 1)^{-1}g(x) = \int_0^\pi G(x, y)g(y) dy.$$

One may verify that

- $R_{\mathcal{A}}(-1) : X \rightarrow D(\mathcal{A})$.

This follows immediately from (1.9).

- $R_{\mathcal{A}}(-1)$ is a compact operator in X . That is, for any bounded sequence $(g_k)_k$ in X , the function sequence $(R_{\mathcal{A}}(-1)g_k)_k$ has a subsequence which converges in X .

This follows from the properties of G and a compactness argument, e.g. Arzela-Ascoli theorem.

- $R_{\mathcal{A}}(-1)$ is symmetric. That is,

$$\langle R_{\mathcal{A}}(-1)g, h \rangle = \langle g, R_{\mathcal{A}}(-1)h \rangle, \quad \forall g, h \in X.$$

This follows from the symmetry property of $G(x, y) = G(y, x) \in \mathbb{R}$.

1.2.2 Spectral theorem and Fourier sine series

Spectral theorem¹¹ states that given an infinite-dimensional inner product space X and a compact symmetric operator $\mathcal{R} : X \rightarrow X$, there exists a sequence of real eigenvalues $(z_n)_n \subset \mathbb{R}$ converging to 0. The corresponding normalized eigenvectors $(v_n)_n$ form an orthonormal set and every $f \in \text{Ran}(\mathcal{R})$ can be written as

$$f = \sum_{n=1}^{\infty} \langle f, v_n \rangle v_n.$$

If $\text{Ran}(\mathcal{R})$ is dense, then the eigenvectors form an orthonormal basis.

alternative theorem.

The unique solution of (1.9) can be represented by Green function (1.10): $u(x) = \int_0^x G(x, y)g(y) dy + \int_x^\pi G(x, y)g(y) dy$, where the Green function has the following properties

- $G(x, y) \in C([0, \pi]^2)$, $G(x, y) \in C^2(\Delta_\pm)$ with $\Delta_\pm = \{(x, y) \in [0, \pi]^2 \mid \pm(x - y) \leq 0\}$, and $\partial_x G$ has a jump at the diagonal line: $\partial_x G(x, x - 0) - \partial_x G(x, x + 0) = -1$;
- $G(\cdot, y)$ satisfies the homogeneous ODE (with respect to x variable) in $\Delta_\pm$ and the boundary conditions.

¹¹We don't give the proof here.

Thus for the compact symmetric operator $R_{\mathcal{A}}(-1)$ in X , there exists a sequence of real eigenvalues $(z_n) \rightarrow 0$ and the corresponding normalized eigenfunctions

$$(v_n) \subset D(\mathcal{A}), \quad \|v_n\| = \langle v_n, v_n \rangle = 1,$$

which form an orthonormal basis in the closure of its range $\overline{\text{Ran}(R_{\mathcal{A}}(-1))}^{L^2} = \overline{D(\mathcal{A})}^{L^2} = L^2(0, \pi)$. That is, for any $f \in L^2(0, \pi)$, we can expand it as follows:

$$f = \sum_{n=1}^{\infty} \langle f, v_n \rangle v_n.$$

Now we return to the operator $\mathcal{A}$. The fact that $R_{\mathcal{A}}(-1)v_n = z_nv_n$ is equivalent to

$$\mathcal{A}v_n = (-1 + \frac{1}{z_n})v_n,$$

that is, v_n is a normalized eigenfunction of $\mathcal{A}$ associated with the eigenvalue $-1 + \frac{1}{z_n}$ which tends to infinity. Recalling (1.8), we find

$$\lambda_n = n^2 = -1 + \frac{1}{z_n}, \quad u_n = \sqrt{\frac{2}{\pi}} \sin(nx) = v_n.$$

To conclude, the expansion $f = \sum_{n=1}^{\infty} \langle f, v_n \rangle v_n$ for any function $f \in L^2(0, \pi)$ is indeed (1.4):

$$f = \sum_{n=1}^{\infty} b_n \sin(nx), \quad b_n = \sqrt{\frac{2}{\pi}} \langle f, v_n \rangle = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(nx) dx.$$

This is the Fourier sine series. Let us emphasize here that the series converges with respect to our scalar product, i.e. in the sense of $L^2(0, \pi)$.

We remark that if the function $f \in D(\mathcal{A})$, then the sine series converges uniformly to f . The convergence of Fourier series is discussed in more detail in Sections 1.3 and 1.4 later.

[Sep 9, 2026]

[Sep 15, 2026]

1.2.3 Eigenvalue problem with Neumann boundary condition

Instead of Dirichlet boundary condition studied in Section 1.2, we consider the Laplace eigenvalue problem associated to the Neumann boundary condition. More precisely, we consider the eigenvalue problem

$$\mathcal{B}u = \lambda u, \quad \mathcal{B} : D(\mathcal{B}) \rightarrow X, \quad \mathcal{B}(u) = -u'' \text{ for } u \in D(\mathcal{B}), \quad (1.11)$$

where the domain of the operator $\mathcal{B}$ is

$$D(\mathcal{B}) = \{u \in C^2([0, \pi]) \mid u'(0) = u'(\pi) = 0\} \subset X,$$

with $X = C([0, \pi]; \mathbb{C})$ equipped with the inner product $\langle \cdot, \cdot \rangle$.

The same analysis as in Subsection 1.1.1 yields the following eigenvalues and normalized eigenfunctions

$$\lambda_0 = 0, \quad \lambda_n = n^2, \quad u_0 = \sqrt{\frac{1}{\pi}}, \quad u_n = \sqrt{\frac{2}{\pi}} \cos(nx), \quad n \in \mathbb{N}.$$

We can then compute the resolvent $R_{\mathcal{B}}(-1) : X \rightarrow D(\mathcal{B})$ in terms of Green function

$$R_{\mathcal{B}}(-1)h(x) = \int_0^\pi G_{\mathcal{B}}(x, y)h(y) \, dy, \quad G_{\mathcal{B}}(x, y) = \begin{cases} -\frac{\phi_{\mathcal{B}}(x)\psi_{\mathcal{B}}(y)}{W(\phi_{\mathcal{B}}, \psi_{\mathcal{B}})} & \text{for } x \leq y \\ -\frac{\phi_{\mathcal{B}}(y)\psi_{\mathcal{B}}(x)}{W(\phi_{\mathcal{B}}, \psi_{\mathcal{B}})} & \text{for } x \geq y \end{cases}$$

and verify that it is compact and symmetric, as in Subsection 1.2.1¹². Finally the spectral theorem ensures that any function $f \in L^2(0, \pi)$ can be expanded as Fourier cosine series

$$f = a_0 + \sum_{n=1}^{\infty} a_n \cos(nx),$$

$$a_0 = \frac{1}{\pi} \int_0^\pi f(x) \, dx, \quad a_n = \frac{2}{\pi} \int_0^\pi f(x) \cos(nx) \, dx, \quad n \in \mathbb{N},$$

in the sense of $L^2(0, \pi)$.

1.2.4 Full Fourier series

We now consider the eigenvalue problem

$$\mathcal{C}u = \lambda u, \quad \mathcal{C} : D(\mathcal{C}) \rightarrow Y, \quad \mathcal{C}(u) = -u'' \text{ for } u \in D(\mathcal{C}), \quad (1.12)$$

where

- $Y = C([-\pi, \pi]; \mathbb{C})$ is equipped with the inner product $\langle \cdot, \cdot \rangle_{L^2(-\pi, \pi)}$;
- we assume periodic boundary conditions on the interval $[-\pi, \pi]$

$$D(\mathcal{C}) = \{u \in C^2([-\pi, \pi]) \mid u(-\pi) = u(\pi), \quad u'(-\pi) = u'(\pi)\} \subset Y.$$

¹²We omit the details here.

There are eigenvalues

$$\lambda_0 = 0, \quad \lambda_n = n^2, \quad n \in \mathbb{N},$$

and the corresponding normalized eigenfunctions

$$u_0 = \sqrt{\frac{1}{2\pi}}, \quad \{u_n, v_n\} = \left\{ \sqrt{\frac{1}{\pi}} \cos(nx), \sqrt{\frac{1}{\pi}} \sin(nx) \right\}, \quad n \in \mathbb{N}.$$

The spectral theorem implies that any function $f \in L^2(-\pi, \pi)$ can be expanded as Fourier series in the sense of $L^2(-\pi, \pi)$:

$$f = a_0 + \sum_{n=1}^{\infty} a_n \cos(nx) + \sum_{n=1}^{\infty} b_n \sin(nx), \quad a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx,$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx, \quad b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx, \quad n \in \mathbb{N}.$$

Observe that

- If $f \in C([-\pi, \pi])$ is an odd function, then $a_0 = a_n = 0$, and hence the full Fourier series of any odd function on $(-\pi, \pi)$ coincides with Fourier sine series on $(0, \pi)$ in the sense of $L^2(0, \pi)$.

Conversely, if $f \in C([0, \pi])$ with Dirichlet boundary condition $f(0) = f(\pi) = 0$, then one can extend it to an odd function in $C([-\pi, \pi])$ satisfying $f(-\pi) = f(0) = f(\pi) = 0$, and the Fourier sine series coincides with the full Fourier series on $(0, \pi)$ in the sense of $L^2(0, \pi)$.

This is the physical intuition why the Fourier sine series is “complete” on $[0, \pi]$.

- If $f \in C([-\pi, \pi])$ is an even function, then $b_n = 0$, and hence the full Fourier series of any even function on $(-\pi, \pi)$ coincides with Fourier cosine series on $(0, \pi)$ in the sense of $L^2(0, \pi)$.

Conversely, if $f \in C([0, \pi])$ with Neumann boundary condition $f'(0) = f'(\pi) = 0$, then one can extend it to an even function in $C([-\pi, \pi])$ satisfying $f'(-\pi) = f'(0) = f'(\pi) = 0$, and the Fourier cosine series coincides with full Fourier series on $(0, \pi)$ in the sense of $L^2(0, \pi)$.

- Replacing

$$\cos(nx) = \frac{1}{2}(e^{inx} + e^{-inx}), \quad \sin(nx) = \frac{1}{2i}(e^{inx} - e^{-inx}),$$

we can rewrite the full Fourier series in the elegant complex form

$$\begin{aligned} f &= \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} c_n e^{inx}, \quad \frac{1}{\sqrt{2\pi}} c_0 = a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-i0x} dx, \\ \frac{1}{\sqrt{2\pi}} c_n &= \frac{1}{2} a_n + \frac{1}{2i} b_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-inx} dx, \quad n \in \mathbb{N}, \\ \frac{1}{\sqrt{2\pi}} c_{-n} &= \frac{1}{2} a_n - \frac{1}{2i} b_n = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-i(-n)x} dx, \quad n \in \mathbb{N}. \end{aligned}$$

1.3 Pointwise convergence of Fourier series

The spectral theorem guarantees the convergence of Fourier series

$$f(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} c_n e^{inx}, \quad c_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} f(x) e^{-inx} dx \quad (1.13)$$

in the sense of $L^2(-\pi, \pi)$. There is a long history of mathematicians searching for the convergence of the partial sum of the Fourier series

$$S_N f(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-N}^N c_n e^{inx}, \quad c_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} f(x) e^{-inx} dx \quad (1.14)$$

as $N \rightarrow \infty$, at first in the sense of pointwise convergence.

1.3.1 Dirichlet kernel

The first positive results is due to P.G.L. Dirichlet (1829). He rewrote the partial sum of a 2π -periodic function as

$$\begin{aligned} S_N f(x) &= \frac{1}{2\pi} \sum_{n=-N}^N \int_{-\pi}^{\pi} f(t) e^{-int} dt e^{inx} \\ &= \frac{1}{2\pi} \sum_{n=-N}^N \int_{-\pi}^{\pi} f(t) e^{in(x-t)} dt \\ &= \int_{-\pi}^{\pi} f(t) D_N(x-t) dt = \int_{-\pi}^{\pi} f(x-t) D_N(t) dt = (f * D_N)(x), \end{aligned}$$

where D_N is the so-called Dirichlet kernel

$$D_N(y) = \frac{1}{2\pi} \sum_{n=-N}^N e^{iny}$$

$$\begin{aligned}
&= \frac{1}{2\pi} (1 + 2\cos y + 2\cos(2y) + \cdots + 2\cos(Ny)) \\
&= \frac{1}{2\pi} \frac{1}{\sin(\frac{1}{2}y)} \left(\sin(\frac{1}{2}y) + (\sin(\frac{3}{2}y) - \sin(\frac{1}{2}y)) + (\sin(\frac{5}{2}y) - \sin(\frac{3}{2}y)) + \right. \\
&\quad \left. + \cdots + (\sin((N + \frac{1}{2})y) - \sin((N - \frac{1}{2})y)) \right) \\
&= \frac{\sin((N + \frac{1}{2})y)}{2\pi \sin(\frac{1}{2}y)}.
\end{aligned}$$

Observe that $D_N(y)$ satisfies

$$\int_{-\pi}^{\pi} D_N(y) dy = 1 \text{ and } |D_N(y)| \leq \frac{1}{2\pi \sin(\frac{1}{2}\delta)} \text{ if } \delta \leq |y| \leq \pi. \quad (1.15)$$

Notice that $\int_{-\pi}^{\pi} |D_N(y)| dy$ is growing in terms of N of the rate $\ln N$ ¹³:

$$\begin{aligned}
\int_{-\pi}^{\pi} |D_N(y)| dy &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left| \frac{\sin((N + \frac{1}{2})y)}{\sin(\frac{1}{2}y)} \right| dy = \frac{1}{\pi} \int_0^{\pi} \frac{|\sin((N + \frac{1}{2})y)|}{\frac{1}{2}y} dy + O(1) \\
&= \frac{2}{\pi} \int_0^{(N + \frac{1}{2})\pi} \frac{|\sin t|}{t} dt + O(1) = \frac{2}{\pi} \sum_{k=0}^{N-1} \int_{k\pi}^{(k+1)\pi} \frac{|\sin t|}{t} dt + O(1) \\
&= \frac{2}{\pi} \sum_{k=0}^{N-1} \int_0^{\pi} \frac{|\sin t|}{t + k\pi} dt + O(1) \\
&= \frac{2}{\pi} \int_0^{\pi} \sin t \left(\sum_{k=0}^{N-1} \frac{1}{t + k\pi} \right) dt + O(1) = O(\ln N). \quad (1.16)
\end{aligned}$$

We claim that the pointwise convergence of a Fourier series at some point x depends on the regularity/smoothness of the function at x , and hence is (surprisingly) a local property (cf. Riemann's "second theorem" in his Habilitation thesis (1854)¹⁴), although the definition of Fourier series depends on the values of this function on the whole interval $[-\pi, \pi]$. To see this, consider first the special case $f \in L^2(-\pi, \pi)$ which satisfies $f = 0$ on a neighborhood of x : $(x - \delta, x + \delta)$, and we compute

$$S_N f(x) = \int_{\delta \leq |y| \leq \pi} f(x - y) \frac{\sin((N + \frac{1}{2})y)}{2\pi \sin(\frac{1}{2}y)} dy$$

¹³This growth is the reason for the failure of the pointwise convergence of Fourier series for continuous functions, see du Bois-Reymond's negative result later.

¹⁴Bernhard Riemann, Ueber die Darstellbarkeit einer Function durch eine trigonometrische Reihe. Abhandlungen der Königlichen Gesellschaft der Wissenschaften zu Göttingen 13, 87-132, 1867. (Habilitationsschrift, Göttingen, 1854)

$$\begin{aligned}
&= \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x-y) \frac{1}{\sin(\frac{1}{2}y)} 1_{\{\delta \leq |y| \leq \pi\}} \left(\frac{1}{2i} (e^{(N+\frac{1}{2})iy} - e^{-(N+\frac{1}{2})iy}) \right) dy \\
&= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(f(x-y) \frac{1}{\sin(\frac{1}{2}y)} 1_{\{\delta \leq |y| \leq \pi\}} \frac{1}{2i} e^{\frac{1}{2}iy} \right) e^{iNy} dy \\
&\quad - \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(f(x-y) \frac{1}{\sin(\frac{1}{2}y)} 1_{\{\delta \leq |y| \leq \pi\}} \frac{1}{2i} e^{-\frac{1}{2}iy} \right) e^{-iNy} dy.
\end{aligned}$$

This is the difference of the Fourier coefficients of two integrable functions at $-N$ and N , both of which tends to 0 as $N \rightarrow \infty$: At this point x , by the Riemann-Lebesgue lemma^{15 16},

$$\lim_{N \rightarrow \infty} S_N f(x) = 0 = f(x).$$

1.3.2 Dini's criterion and du Bois-Reymond's counterexample

Now we consider the function f with general behavior on $(x - \delta, x + \delta)$ and compute (due to the unit size of the integral of D_N)

$$S_N f(x) - f(x) = \int_{-\pi}^{\pi} (f(x-y) - f(x)) \frac{\sin((N + \frac{1}{2})y)}{2\pi \sin(\frac{1}{2}y)} dy = \int_{|y| < \delta} + \int_{\delta \leq |y| \leq \pi}.$$

Dini's criterion If we assume the following continuity condition for the function f at the point x :

$$\int_{|y| < \delta} \left| \frac{f(x-y) - f(x)}{y} \right| dy < \infty, \quad (1.17)$$

¹⁵For any integrable function g which is periodic on $\mathbb{R}$ with period 2π , we compute

$$\begin{aligned}
\hat{g}(N) &:= \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} g(y) e^{-iNy} dy = -\frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} g(y) e^{-iN(y+\frac{\pi}{N})} dy \\
&= -\frac{1}{\sqrt{2\pi}} \int_{-\pi+\frac{\pi}{N}}^{\pi+\frac{\pi}{N}} g(t - \frac{\pi}{N}) e^{-iNt} dt = -\frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} g(y - \frac{\pi}{N}) e^{-iNy} dy,
\end{aligned}$$

and thus

$$\hat{g}(N) = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} g(y) e^{-iNy} dy = \frac{1}{2} \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} (g(y) - g(y - \frac{\pi}{N})) e^{-iNy} dy.$$

If g is continuous, then its Fourier coefficient $\hat{g}(N)$ tends to 0 as N tends to infinity. If $g \in L^1(-\pi, \pi)$, then we can approximate it with some continuous function to derive the convergence result.

¹⁶Henri Lebesgue, *Leçons sur les séries trigonométriques*, Gauthier-Villars, Paris, 1906; the integral itself is from *Intégrale, longueur, aire*, *Annali di Matematica Pura ed Applicata* (3) 7, 231–359, 1902.

for some $\delta > 0$, then all the functions

$$\frac{f(x-y) - f(x)}{\sin(\frac{1}{2}y)} 1_{\{|y| < \delta\}} \frac{1}{2i} e^{\pm \frac{1}{2}iy}, \quad \frac{f(x-y) - f(x)}{\sin(\frac{1}{2}y)} 1_{\{\delta \leq |y| \leq \pi\}} \frac{1}{2i} e^{\pm \frac{1}{2}iy}$$

are integrable on $[-\pi, \pi]$, and hence Riemann-Lebesgue lemma implies

$$S_N f(x) \rightarrow f(x) \text{ as } N \rightarrow \infty.$$

The continuity property (1.17) is called the Dini's criterion (1880)¹⁷ for the pointwise convergence of Fourier series.

Any Hölder continuous function $f \in C^\alpha$ such that $|f(x-y) - f(x)| \leq C|y|^\alpha$ for $|y| < \delta$ for some $\delta > 0$ satisfies the Dini's criterion (1.17). In particular, Fourier sine series in (1.4) converges uniformly on $[0, \pi]$ to the function f in (1.5).

[Sep 15, 2026]

[Sep 16, 2026]

Du Bois-Reymond's counterexample However, continuous functions need not satisfy Dini's condition, and we have the negative result by P. du Bois-Reymond (1873)¹⁸: *There exists a continuous function whose Fourier series diverges at a point.*

We don't follow his original proof, while apply the Banach-Steinhaus theorem (so-called uniform boundedness principle) instead to show this negative result. To this end, let¹⁹

$$Z = \{f \in C([-\pi, \pi]; \mathbb{C}) \mid \|f\|_\infty = \sup_{x \in [-\pi, \pi]} |f(x)| < \infty\} = C_b([-\pi, \pi]; \mathbb{C})$$

denote the usual continuous function space on $[-\pi, \pi]$, which is a Banach space. Let $\{T_N\}_N$ be a family of linear bounded operators from Z to $\mathbb{C}$, which is a normed vector space, given by

$$T_N : Z \rightarrow \mathbb{C}, \text{ via } T_N f = (S_N f)(0) = \int_{-\pi}^{\pi} f(t) D_N(t) dt,$$

where $D_N(x) = \frac{\sin((N+\frac{1}{2})x)}{2\pi \sin(\frac{1}{2}x)}$. By Banach-Steinhaus theorem²⁰, either

$$\sup_N \|T_N\|_{Z \rightarrow \mathbb{C}} < \infty,$$

¹⁷Ulisse Dini, *Serie di Fourier e altre rappresentazioni analitiche delle funzioni di una variabile reale*. Pisa: Tipografia T. Nistri, 1880.

¹⁸Paul du Bois-Reymond, Über die Fourier'schen Reihen, *Nachrichten der Königlichen Gesellschaft der Wissenschaften und der Georg-August-Universität zu Göttingen*, 1873.

¹⁹The space Z here is different from the space Y given in (1.12).

²⁰Stefan Banach and Hugo Dyonizy Steinhaus, *Sur le principe de la condensation de singularités*, *Fundamenta Mathematicae* 9, 50–61, 1927.

or there exists an element $f \in Z$ such that

$$\sup_N |T_N f| = \infty.$$

For any fixed $N \in \mathbb{N}$, D_N has only a finite number of zeros, and hence $\operatorname{sgn}(D_N)$ has only a finite number of jumps. Thus for any $\varepsilon > 0$, there exists a continuous function f such that

$$\|f\|_\infty = 1 \text{ and } T_N f \geq d_N - \varepsilon, \quad d_N := \int_{-\pi}^{\pi} |D_N(t)| dt.$$

Obviously the operator norm $\|T_N\|_{Z \rightarrow \mathbb{C}}$ is bounded by d_N , and therefore

$$\|T_N\|_{Z \rightarrow \mathbb{C}} = d_N. \tag{1.18}$$

Recall (1.16): $d_N = O(\ln N)$ as $N \rightarrow \infty$, and hence

$$\|T_N\|_{Z \rightarrow \mathbb{C}} = O(\ln N) \text{ as } N \rightarrow \infty.$$

Banach-Steinhaus theorem then implies that there exists a continuous function f such that

$$\limsup_{N \rightarrow \infty} |(S_N f)(0)| = \infty.$$

To conclude, we find a continuous function $f \in C([-\pi, \pi])$, which is uniformly bounded on $[-\pi, \pi]$, while $|(S_N f)(0)| \rightarrow \infty$, and hence $(S_N f)(0) \not\rightarrow f(0)$ as $N \rightarrow \infty$.

1.3.3 Dirichlet-Jordan's criterion

The Jordan's criterion (1881)²¹ investigates this pointwise convergence property from another perspective. Namely, if *the function f has a bounded variation in the neighborhood of the point x* , then the pointwise convergence holds in the following sense:

$$\lim_{N \rightarrow \infty} S_N f(x) = \frac{1}{2}(f(x-0) + f(x+0)). \tag{1.19}$$

We give a short proof here. Since every function of bounded variation is the difference of two monotonic functions, we assume without loss of generality

²¹Camille Jordan, Sur la série de Fourier, *Comptes Rendus de l'Académie des Sciences de Paris*, 1881.

that f is monotonic in some neighborhood of the point x . Since $D_N(t) = D_N(-t)$, we rewrite

$$\begin{aligned} S_N f(x) &= \int_0^\pi f(x-t) D_N(t) dt + \int_{-\pi}^0 f(x-t) D_N(t) dt \\ &= \int_0^\pi (f(x-t) + f(x+t)) D_N(t) dt, \end{aligned}$$

and it suffices to show for any monotonic function g , it holds

$$\lim_{N \rightarrow \infty} \int_0^\pi g(t) D_N(t) dt = \frac{1}{2} g(0_+). \quad (1.20)$$

Indeed, one may assume further $g(0_+) = 0$, then for any $\varepsilon > 0$ there exists $\delta > 0$ such that $|g|_{(0,\delta)} < \varepsilon$, and there exists $\xi \in [0, \delta]$ such that

$$\begin{aligned} \int_0^\pi g(t) D_N(t) dt &= \int_0^\delta g(t) D_N(t) dt + \int_\delta^\pi g(t) D_N(t) dt \\ &= g(\delta_-) \int_\xi^\delta D_N(t) dt + \int_\delta^\pi g(t) D_N(t) dt, \end{aligned}$$

where the second equality follows from the second mean value theorem for integrals²². The first integral above is bounded by

$$\varepsilon \left| \int_\xi^\delta \frac{\sin((N + \frac{1}{2})t)}{2\pi \sin(\frac{1}{2}t)} dt \right| \leq \varepsilon \left(\left| \int_\xi^\delta \frac{\sin((N + \frac{1}{2})t)}{\pi t} dt \right| + C \right)$$

²²Given any monotonically increasing function g and continuous function f on $[a, b]$, we compute further the following Lebesgue-Stieltjes integral as

$$\int_a^b f(x) dg(x) = (g(b) - g(a)) \int_a^b f(x) d \frac{g(x) - g(a)}{g(b) - g(a)} \in (g(b) - g(a)) [\min_{[a,b]} f, \max_{[a,b]} f].$$

Thus there exists $\xi \in [a, b]$ such that $\int_a^b f(x) dg(x) = f(\xi)(g(b) - g(a))$, which is the second mean value theorem for Stieltjes integrals. The same result holds for monotonically decreasing function g .

Now let $F(x) = \int_a^x f(t) dt$, then we use integration by parts and the second mean value theorem above to derive

$$\begin{aligned} \int_a^b f(x) g(x) dx &= \int_a^b F'(x) g(x) dx = [F(x) g(x)]_a^b - \int_a^b F(x) dg(x) \\ &= F(b) g(b) - F(\xi)(g(b) - g(a)) = g(b) \int_\xi^b f(x) dx + g(a) \int_a^\xi f(x) dx. \end{aligned}$$

$$\leq \varepsilon \left(\left| \int_{(N+\frac{1}{2})\xi}^{(N+\frac{1}{2})\delta} \frac{\sin t}{\pi t} dt \right| + C \right) \leq \tilde{C}\varepsilon,$$

while the second integral tends to 0 as $N \rightarrow \infty$ by Riemann-Lebesgue lemma. Hence (1.20) holds and correspondingly (1.19) follows. In particular, if f is continuous and has a bounded variation in a neighborhood of x , then

$$\lim_{N \rightarrow \infty} S_N f(x) = f(x).$$

The classical Dirichlet's pointwise convergence criterion (1829)²³:

- f is bounded and piecewise continuous;
- f has a finite number of maxima and minima,

is a special case of Jordan's criterion.

1.3.4 Fejér's kernel

The pointwise convergece theory for Fourier series of continuous functions was “rescued” by L. Fejér in 1900²⁴. Instead of the natural partial sum of Fourier series where Dirichlet kernel D_N is involved, L. Fejér made use of the arithmetic means of partial sums (Cesàro's summation)

$$\sigma_N = \frac{1}{N+1} \sum_{k=0}^N S_k,$$

and the associated Fejér kernel

$$\begin{aligned} F_N(y) &= \frac{1}{N+1} \sum_{k=0}^N D_k(y) \\ &= \frac{1}{N+1} \frac{1}{2\pi} \frac{\sin(\frac{1}{2}y) + \sin(\frac{3}{2}y) + \sin(\frac{5}{2}y) + \cdots + \sin((N+\frac{1}{2})y)}{\sin(\frac{1}{2}y)} \\ &= \frac{1}{N+1} \frac{1}{2\pi} \frac{\cos 0 - \cos((N+1)y)}{2(\sin(\frac{1}{2}y))^2} \\ &= \frac{1}{N+1} \frac{1}{2\pi} \left(\frac{\sin(\frac{1}{2}(N+1)y)}{\sin(\frac{1}{2}y)} \right)^2 \geq 0. \end{aligned}$$

²³Peter Gustav Lejeune Dirichlet, Sur la convergence des séries trigonométriques qui servent à représenter une fonction arbitraire entre des limites données, *Journal für die reine und angewandte Mathematik* (Crelle's journal), 1829.

²⁴Lipót Fejér, Sur les fonctions bornées et intégrables, *Comptes rendus hebdomadaires des séances de l'Académie des sciences*, 1900.

Note that unlike the sign-changing Dirichlet kernel D_N , $F_N \geq 0$ is nonnegative. F_N satisfies also (1.15):

$$\int_{-\pi}^{\pi} F_N(y) dy = 1 \text{ and } F_N(y) \leq \frac{1}{N+1} \frac{1}{2\pi} \frac{1}{(\sin(\frac{1}{2}\delta))^2} \text{ if } \delta \leq |y| \leq \pi,$$

and is an approximate identity. Then we have immediately that for any continuous function $f \in C([-\pi, \pi])$,

$$\lim_{N \rightarrow \infty} \sup_{x \in [-\pi, \pi]} |(\sigma_N f)(x) - f(x)| = 0.$$

Indeed,

$$\begin{aligned} |(\sigma_N f)(x) - f(x)| &= \left| \int_{-\pi}^{\pi} (f(x-t) - f(x)) F_N(t) dt \right| \\ &\leq \int_{|t| < \delta} |f(x-t) - f(x)| F_N(t) dt + 2\|f\|_{\infty} \int_{\delta \leq |t| \leq \pi} F_N(t) dt, \end{aligned}$$

where δ can be chosen such that the first integral be arbitrarily small, while for any fixed δ , the second integral tends to 0 as N tends to ∞ .

The above argument can be used to show the $L^p(-\pi, \pi)$ convergence of f by $\sigma_N f$ for $p \in [1, \infty)$, as long as one notes $\lim_{t \rightarrow 0} \|f(x-t) - f(x)\|_{L^p_x} = 0$ for $p \in [1, \infty)$. This implies the following facts:

- The trigonometric polynomials are dense in $L^p(-\pi, \pi)$, $p \in [1, \infty)$.
- Weierstrass approximation theorem²⁵: Every continuous function on the torus is a uniform limit of trigonometric polynomials.
- If f is integrable with $c_n = 0$ for all $n \in \mathbb{Z}$, then f is identically zero (a.e.).

[Sep 16, 2026]

[Sep 22, 2026]

1.4 Convergence in norm of Fourier series

We have already achieved the satisfactory convergence result of Fourier series in the sense of $L^2(-\pi, \pi)$ by the spectral theorem:

$$\lim_{N \rightarrow \infty} \|S_N f - f\|_{L^2(-\pi, \pi)} = 0, \quad \forall f \in L^2(-\pi, \pi). \quad (1.21)$$

²⁵Karl Theodor Wilhelm Weierstrass, *Über die analytische Darstellbarkeit sogenannter willkürlicher Functionen einer reellen Veränderlichen*, Sitzungsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin (1885), 633–639, 789–805.

where the partial sum $S_N f$ is defined in (1.14). Furthermore, the mapping $f \mapsto (c_n)$ is an isometry from $L^2(-\pi, \pi)$ to $\ell^2(\mathbb{Z})$ by Parseval's identity²⁶ Δ ,

$$\|f\|_{L^2(-\pi, \pi)}^2 = \sum_{n=-\infty}^{\infty} |c_n|^2. \quad (1.22)$$

We may also ask whether the convergence result (1.21) holds in the sense of $L^p(-\pi, \pi)$.

- Case $p = \infty$.

Du Bois-Reymond's counterexample in Subsection 1.3.2 gives the negative answer for the case $p = \infty$.

- If $p \in [1, \infty)$, we have the following equivalence relation for $p \in [1, \infty)$:

$$\|S_N f - f\|_{L^p(-\pi, \pi)} \rightarrow 0 \text{ as } N \rightarrow \infty, \quad \forall f \in L^p(-\pi, \pi) \quad (1.23)$$

$$\Leftrightarrow \exists C = C_p \text{ such that uniformly in } N, \quad \|S_N\|_{L^p(-\pi, \pi) \rightarrow L^p(-\pi, \pi)} \leq C. \quad (1.24)$$

The necessity of (1.24) follows from the uniform boundedness principle. The other direction follows from the following facts:

- Density property: $C_0^\infty(-\pi, \pi)$ is dense in $L^p(-\pi, \pi)$;
- Uniform convergence and hence convergence in L^p -norm for a dense subset $C_0^\infty(-\pi, \pi)$: For any $\varphi \in C_0^\infty(-\pi, \pi)$, $\|S_N \varphi - \varphi\|_{L^p(-\pi, \pi)} \leq (2\pi)^{\frac{1}{p}} \|S_N \varphi - \varphi\|_{L^\infty(-\pi, \pi)} \rightarrow 0$ as $N \rightarrow \infty$,

and the decomposition $S_N f - f = S_N(f - \varphi) + (S_N \varphi - \varphi) + (\varphi - f)$. Therefore

- Case $p = 1$.

Similar as the analysis for the norm of T_N in (1.18)²⁷, we obtain

$$\|S_N\|_{L^1 \rightarrow L^1} = \|D_N\|_{L^1(-\pi, \pi)} = O(\ln N) \text{ as } N \rightarrow \infty,$$

and thus the convergence statement (1.23) fails for $p = 1$ ²⁸.

²⁶Marc-Antoine Parseval des Chênes, *Mémoire sur les séries et sur l'intégration complète d'une équation aux différences partielles linéaires du second ordre à coefficients constants*, Mémoires de l'Académie des Sciences de l'Institut de France 2, 593–611, 1806.

²⁷This is due to Young's inequality: $\|S_N f\|_{L^1} \leq \|D_N\|_{L^1} \|f\|_{L^1}$ and the result (1.18): $\|S_N\|_{L^1 \rightarrow L^1} = \|S_N^*\|_{L^\infty \rightarrow L^\infty} \geq \|D_N\|_{L^1}$.

²⁸Andrey Nikolaevich Kolmogorov, Une série de Fourier-Lebesgue divergente presque partout. *Fundamenta Mathematicae* 4, 324–328, 1923.

– Case $p \in (1, \infty)$.

We claim that the convergence result (1.23) for $p \in (1, \infty)$ follows from the L^p -boundedness of the conjugate function discussed below.

1.4.1 Conjugate function and Riesz projections

In the functional setting of $L^2(-\pi, \pi)$, we have shown (1.13):

$$f(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{inx}, \quad \hat{f}(n) := c_n = \frac{1}{\sqrt{2\pi}} \int_{-\pi}^{\pi} f(x) e^{-inx} dx. \quad (1.25)$$

We define the conjugate function of a function f as²⁹³⁰

$$\mathcal{H}(f)(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \left(\frac{1}{i} \operatorname{sgn}(n)\right) \hat{f}(n) e^{inx}, \quad (1.26)$$

where $\operatorname{sgn}(0) = 0$, as well as Riesz projections $P_{\pm}$ by³¹

$$P_+(f)(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\infty} \hat{f}(n) e^{inx}, \quad (1.27)$$

$$P_-(f)(x) = -\frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{-1} \hat{f}(n) e^{inx}. \quad (1.28)$$

²⁹Historically it was the concept of harmonic conjugate (see (1.29)-(1.32) below) appeared first, and the pair (u, v) satisfies the Cauchy-Riemann equations. The conjugate function $\mathcal{H}(f)$ happens to be the limit of the harmonic conjugate v on the unit circle (see (1.34) below).

Augustin-Louis Cauchy, *Mémoire sur les intégrales définies prises entre des limites imaginaires*, de Bure Frères, Paris, 1825.

Bernhard Riemann, *Grundlagen für eine allgemeine Theorie der Functionen einer veränderlichen complexen Grösse*, Inauguraldissertation, Göttingen, 1851; reprinted in *Gesammelte mathematische Werke und wissenschaftlicher Nachlass*, B. G. Teubner, Leipzig, 3–48, 1876.

³⁰We will discuss the counter part, namely the Hilbert transform, on the whole line in Subsection 3.3.2 later.

³¹This is after Marcel Riesz (1924, 1927), to recognize his contribution to the L^p -boundedness of the conjugate function for $p \in (1, \infty)$.

The Riesz projection $P = \frac{1}{2\pi i} \oint_{\Gamma} (\lambda I - T)^{-1} d\lambda$, which is after his brother Frigyes Riesz, is related to the spectral projection which we discussed earlier in Section 1.2.

Hardy space (1915) L^2_+ consists of L^2 functions with Fourier transform supported on the domain $\{n \geq 0\}$, that is $L^2_+ = \{P_+ f \mid f \in L^2\}$. It identifies holomorphic functions F on the unit disc such that $\sup_{r < 1} \int_{-\pi}^{\pi} |F(re^{ix})|^2 dx < \infty$.

Then the following relations hold

$$f = P_+(f) - P_-(f), \quad P_+(f) + P_-(f) = i\mathcal{H}(f) + \frac{1}{\sqrt{2\pi}}\hat{f}(0),$$

or equivalently,

$$P_{\pm}(f) = \frac{1}{2}\left(i\mathcal{H}(f) \pm f + \frac{1}{\sqrt{2\pi}}\hat{f}(0)\right).$$

We now reformulate the partial sum S_N in terms of Riesz operator P_+ and multiplication operators e^{iNx} , e^{i2Nx} . We first rewrite

$$\begin{aligned} S_N f &= \frac{1}{\sqrt{2\pi}} \sum_{n=-N}^N \hat{f}(n) e^{inx} \\ &= \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{2N} \hat{f}(n-N) e^{i(n-N)x} \\ &= e^{-iNx} \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{2N} \widehat{f(\cdot)e^{iN\cdot}}(n) e^{inx}, \end{aligned}$$

where we compute for any $g \in L^2(-\pi, \pi)$ that

$$\begin{aligned} \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{2N} \hat{g}(n) e^{inx} &= \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\infty} \hat{g}(n) e^{inx} - \frac{1}{\sqrt{2\pi}} \sum_{n=2N+1}^{\infty} \hat{g}(n) e^{inx} \\ &= P_+(g) - e^{i(2N+1)x} \frac{1}{\sqrt{2\pi}} \sum_{n=0}^{\infty} \widehat{g(\cdot)e^{-i(2N+1)\cdot}}(n) e^{inx} \\ &= P_+(g) - e^{i(2N+1)x} P_+(ge^{-i(2N+1)\cdot}). \end{aligned}$$

To conclude, we can represent the partial sum operator S_N as

$$S_N f = e^{-iNx} P_+(fe^{iN\cdot}) - e^{i2Nx} P_+(fe^{-i2N\cdot}).$$

Since the multiplication operator e^{iax} , $a \in \mathbb{R}$ is a bounded operator on L^p -space with operator norm 1, we derive for all $N \in \mathbb{N}$,

$$\|S_N\|_{L^p(-\pi, \pi) \rightarrow L^p(-\pi, \pi)} \leq 2\|P_+\|_{L^p(-\pi, \pi) \rightarrow L^p(-\pi, \pi)},$$

which is bounded by the norm $\|\mathcal{H}\|_{L^p(-\pi, \pi) \rightarrow L^p(-\pi, \pi)}$, up to some constants. The fact³² that $\mathcal{H}$ is a bounded operator in $L^p(-\pi, \pi)$, $p \in (1, \infty)$ implies the uniform L^p -boundedness of S_N in (1.24), and hence the convergence result (1.23) holds for $p \in (1, \infty)$.

³²We don't give the proof here, which can be found e.g. in [3, Subsection 4.1.2].

Historically, the $L^p(-\pi, \pi)$, $p \in (1, \infty)$ -norm convergence of Fourier series was established by M. Riesz (1927)³³, who proved the L^p boundedness of the conjugate function for $p \in (1, \infty)$. A modern proof rests on the Marcinkiewicz interpolation theorem³⁴ interpolating the L^2 isometry (1.22) against Kolmogorov's weak $(1, 1)$ bound³⁵ yields the strong (p, p) inequality for $p \in (1, 2)$, while the range $p \in (2, \infty)$ follows by duality, the adjoint of the conjugation operator being its own negative.

The real-variable form of the argument is due to Calderón and Zygmund³⁶, whose decomposition replaced the complex-analytic proofs and supplied the weak $(1, 1)$ estimate for general singular integrals in $\mathbb{R}^n$; in this formulation the decisive object is the maximal function³⁷, $\sup_{r < 1} |P_r f|$ for the Poisson kernel and $\sup_{t > 0} |Q_t f|$ for the conjugate function, whose weak $(1, 1)$ and strong (p, p) bounds govern pointwise and norm behaviour alike, and whose partial-sum analogue $\sup_N |S_N f|$ is precisely what later carried the problem from norm convergence to almost-everywhere convergence.

1.4.2 Harmonic extension and Poisson kernel

We turn to view Fourier series from another perspective, by considering the following function defined in the unit disk on the complex plane:

$$u(z) = \frac{1}{\sqrt{2\pi}} \left(\sum_{n=0}^{\infty} \hat{f}(n) z^n + \sum_{n=-\infty}^{-1} \hat{f}(n) \bar{z}^{|n|} \right), \quad |z| < 1, \quad (1.29)$$

that is,

$$u(re^{ix}) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \hat{f}(n) r^{|n|} e^{inx}, \quad r \in [0, 1), \quad x \in [-\pi, \pi].$$

Thus the Fourier series can be viewed as the formal limit of the harmonic function u on the unit circle,

$$\frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \hat{f}(n) e^{inx} = \lim_{r \rightarrow 1^-} u(re^{ix}), \quad x \in [-\pi, \pi]. \quad (1.30)$$

³³Marcel Riesz, Sur les fonctions conjuguées, *Math. Z.* 27, 218–244, 1927 (announced in Les fonctions conjuguées et les séries de Fourier, *C. R. Acad. Sci. Paris* 178, 1464–1467, 1924).

³⁴Józef Marcinkiewicz, Sur l'interpolation d'opérateurs, *C. R. Acad. Sci. Paris* 208, 1272–1273, 1939.

³⁵Andrey Nikolaevich Kolmogorov, Sur les fonctions harmoniques conjuguées et les séries de Fourier, *Fund. Math.* 7, 23–28, 1925.

³⁶Alberto Pedro Calderón and Antoni Zygmund, On the existence of certain singular integrals, *Acta Math.* 88, 85–139, 1952.

³⁷Godfrey Harold Hardy and John Edensor Littlewood, A maximal theorem with function-theoretic applications, *Acta Math.* 54, 81–116, 1930.

When f is real-valued, $\widehat{f}(-n) = \overline{\widehat{f}(n)}$, u is a real-valued harmonic function. We now analyze the limit on the right hand side. Rewrite, for $r < 1$,

$$u(re^{ix}) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \widehat{f}(n) r^{|n|} e^{inx} = \int_{-\pi}^{\pi} f(x) P_r(x-t) dt,$$

where $P_r(t)$ denotes the Poisson kernel^{38 3940}

$$P_r(t) = \frac{1}{2\pi} \sum_{n=-\infty}^{\infty} r^{|n|} e^{int}.$$

It can be computed explicitly by use of complex variables for $r \in [0, 1)$: Observe that for $\zeta = re^{it}$ with $|\zeta| < 1$,

$$\begin{aligned} P_r(t) &= \frac{1}{2\pi} \operatorname{Re} \left[1 + 2 \sum_{n=1}^{\infty} \zeta^n \right] = \frac{1}{2\pi} \operatorname{Re} \left[\frac{1+\zeta}{1-\zeta} \right] \\ &= \frac{1}{2\pi} \operatorname{Re} \left[\frac{(1+\zeta)(1-\bar{\zeta})}{|1-\zeta|^2} \right] = \frac{1}{2\pi} \frac{1-|\zeta|^2}{|1-\zeta|^2} \\ &= \frac{1}{2\pi} \frac{1-r^2}{1-2r \cos t + r^2} > 0. \end{aligned}$$

Note that similar as the Fejér kernel, $\{P_r\}_{r \in (0,1)}$ is an approximate identity, satisfying

$$P_r(t) \geq 0, \quad \int_{-\pi}^{\pi} P_r(t) dt = 1, \quad \lim_{r \rightarrow 1-} \int_{\delta \leq |t| \leq \pi} P_r(t) dt = 0 \text{ if } \delta > 0.$$

Therefore the argument in Subsection 1.3.4 yields the limit on the right hand side of (1.30) in the following sense:

- Case $p \in [1, \infty)$.

If $f \in L^p(-\pi, \pi)$, then

$$\lim_{r \rightarrow 1-} \|u(re^{ix}) - f(x)\|_{L^p} = \lim_{r \rightarrow 1-} \|P_r * f - f\|_{L^p} = 0;$$

³⁸Siméon Denis Poisson, Mémoire sur l'intégration de quelques équations linéaires aux différences partielles, et particulièrement de l'équation générale du mouvement des fluides élastiques, *Mémoires de l'Académie Royale des Sciences de l'Institut de France* **3** (1820/1823), 121–176.

³⁹Karl Hermann Amandus Schwarz, Über die Integration der partiellen Differentialgleichung $\partial^2 u / \partial x^2 + \partial^2 u / \partial y^2 = 0$ unter vorgeschriebenen Grenz- und Unstetigkeitsbedingungen, *Monatsberichte der Königlich Preussischen Akademie der Wissenschaften zu Berlin*, 767–795, 1870.

⁴⁰Formally, “ $\lim_{r \rightarrow 1-} P_r = \lim_{N \rightarrow \infty} D_N$ ”.

- Case $p = \infty$.

If $f \in C([-\pi, \pi])$ is continuous, then

$$\lim_{r \rightarrow 1-} \|u(re^{ix}) - f(x)\|_{L^\infty} = \lim_{r \rightarrow 1-} \|P_r * f - f\|_{L^\infty} = 0.$$

Note that if f is real-valued, then the function u defined in terms of f in (1.29) is harmonic function inside the unit disk. The above convergence result implies that so-defined function $u(re^{ix}) = (P_r * f)(x)$ is the (unique) solution of the Dirichlet problem for the Laplace equation on the unit disk:

$$\begin{cases} \Delta u = 0 & \text{if } |z| < 1, \\ u = f & \text{if } |z| = 1, \end{cases} \quad (1.31)$$

where the boundary condition is satisfied in the above sense. The function u is thus called the harmonic extension of f inside the unit disk.

1.4.3 Harmonic conjugate and conjugate Poisson kernel

We continue the study of the harmonic function $u(z) = u(re^{ix})$ defined by (1.29) inside the unit disk. Let

$$v(z) = \frac{1}{\sqrt{2\pi}} \left(\sum_{n=1}^{\infty} \frac{1}{i} \hat{f}(n) z^n + \sum_{n=-\infty}^{-1} i \hat{f}(n) \bar{z}^{|n|} \right), \quad |z| < 1, \quad (1.32)$$

that is,

$$v(re^{ix}) = \frac{1}{\sqrt{2\pi}} \sum_{n \in \mathbb{Z}} \left(\frac{1}{i} \operatorname{sgn}(n) \right) \hat{f}(n) r^{|n|} e^{inx}, \quad r \in [0, 1), \quad x \in [-\pi, \pi].$$

Observe that

- If $\{\hat{f}(n)\}$ is a bounded sequence, then v is the harmonic conjugate of u , in the sense that

$$G(z) = u(z) + iv(z) = \frac{1}{\sqrt{2\pi}} \left(\hat{f}(0) + 2 \sum_{n=1}^{\infty} \hat{f}(n) z^n \right), \quad (1.33)$$

is holomorphic in the unit disk.

- Formally, the conjugate function defined in (1.26) can be viewed as the limit of $v(z)$ on the unit circle:

$$\mathcal{H}(f)(x) = \frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{\infty} \left(\frac{1}{i} \operatorname{sgn}(n) \right) \hat{f}(n) e^{inx} = \lim_{r \rightarrow 1-} v(re^{ix}). \quad (1.34)$$

Similarly as for u , we rewrite as a convolution

$$v(re^{ix}) = \int_{-\pi}^{\pi} f(t)Q_r(x-t) dt = (Q_r * f)(x)$$

with the conjugate Poisson kernel

$$Q_r(t) = \frac{1}{2\pi} \sum_{n \in \mathbb{Z}} \left(\frac{1}{i} \operatorname{sgn}(n) \right) r^{|n|} e^{int}, \quad r \in [0, 1), \quad t \in [-\pi, \pi].$$

It is real-valued and we compute by use of complex variable $\zeta = re^{it}$ with $|\zeta| = r < 1$, similarly as for Poisson kernel:

$$\begin{aligned} Q_r(t) &= \frac{1}{2\pi} \operatorname{Im} \left[\sum_{n=1}^{\infty} \zeta^n - \sum_{m=1}^{\infty} \bar{\zeta}^m \right] = \frac{1}{2\pi} \operatorname{Im} \left[1 + 2 \sum_{n=1}^{\infty} \zeta^n \right] \\ &= \frac{1}{2\pi} \operatorname{Im} \left[\frac{1+\zeta}{1-\zeta} \right] = \frac{1}{2\pi} \operatorname{Im} \left[\frac{(1+\zeta)(1-\bar{\zeta})}{|1-\zeta|^2} \right] \\ &= \frac{1}{2\pi} \frac{2r \sin t}{1 - 2r \cos t + r^2}. \end{aligned}$$

Note that

- $P_r(t) + iQ_r(t)$ is holomorphic function in ζ in the unit disk:

$$P_r(t) + iQ_r(t) = \frac{1}{2\pi} \frac{1+\zeta}{1-\zeta} = \frac{1}{2\pi} \left(-1 + \frac{2}{1-\zeta} \right), \quad \text{with } \zeta = re^{it}, \quad r \in [0, 1),$$

such that the holomorphic function $G(z)$ can be represented by it as

$$\begin{aligned} G(z) &= u(re^{ix}) + iv(re^{ix}) = ((P_r + iQ_r) * f)(x) \\ &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(-1 + \frac{2}{1 - re^{i(x-t)}} \right) f(t) dt \\ &\stackrel{w=e^{it}}{=} -\hat{f}(0) + 2 \cdot \frac{1}{2\pi i} \int_{|w|=1} \frac{1}{w-z} f(t) dw. \end{aligned}$$

Here the integral is indeed the Cauchy integral of the function f on the unit circle:

$$F(z) = \frac{1}{2\pi i} \int_{|w|=1} \frac{f(t)}{w-z} dw \stackrel{w=e^{it}}{=} \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{f(t)e^{it}}{e^{it}-z} dt. \quad (1.35)$$

- Unlike P_r , the kernel Q_r is not an approximate identity:

- It has zero mean $\int_{-\pi}^{\pi} Q_r(t) dt = 0$;
- Its L^1 -norm diverges as r tends to 1_- :

$$\begin{aligned}\|Q_r\|_{L^1(-\pi,\pi)} &= \frac{2}{\pi} \int_0^\pi \frac{r \sin t}{1 - 2r \cos t + r^2} dt \stackrel{y=2r \cos t}{=} \frac{1}{\pi} \int_{-2r}^{2r} \frac{1}{1 + r^2 - y} dy \\ &= \frac{1}{\pi} \ln(1 + r^2 - y)|_{-2r}^{2r} = \frac{1}{\pi} \ln \frac{(1+r)^2}{(1-r)^2} \\ &\sim \frac{2}{\pi} \log \frac{1}{1-r} \rightarrow \infty \text{ as } r \rightarrow 1_-.\end{aligned}$$

Hence the argument used for P_r breaks down, and this failure is exactly where the difficulty of the convergence problem lives. Indeed, the convergence (1.34): $\mathcal{H}(f)(x) = \lim_{r \rightarrow 1_-} Q_r * f(x)$ holds in L^p and almost everywhere for $f \in L^p$, $1 \leq p < \infty$, and the theorem of M. Riesz states that the map $f \mapsto \mathcal{H}(f)$ is bounded on L^p for $p \in (1, \infty)$.

1.4.4 Other related topics

Boundary traces of the Cauchy integral from inside and outside

Formally, the relations among the operators $\mathcal{H}$, Id and the functions P_r, Q_r can be seen more clearly in terms of their Fourier transforms⁴¹:

$$\begin{aligned}\sqrt{2\pi} \widehat{P_r}(n) &= r^{|n|}, \quad \lim_{r \rightarrow 1_-} \widehat{P_r}(n) = 1(n) = \widehat{\text{Id}}, \\ \sqrt{2\pi} \widehat{Q_r}(n) &= \frac{1}{i} \text{sgn}(n) r^{|n|}, \quad \lim_{r \rightarrow 1_-} \widehat{Q_r}(n) = \frac{1}{i} \text{sgn}(n) = \widehat{\mathcal{H}}(n).\end{aligned}$$

We remark that the Riesz projections $P_\pm$:

$$P_\pm = \frac{1}{2} \left(\pm \text{Id} + i\mathcal{H} + \frac{1}{2\pi} \int_{-\pi}^{\pi} \right)$$

with

$$\widehat{P_+}(n) = 1_{\{0,1,\dots\}}(n), \quad \widehat{P_-}(n) = -1_{\{-1,-2,-3,\dots\}}(n),$$

correspond to the traces $\lim_{r \rightarrow 1_\mp}$ of the Cauchy integral of f : $F(z)$ given in (1.35). Indeed, for $|z| < 1$, expanding $\frac{1}{w-z} = \sum_{n=0}^{\infty} \frac{z^n}{w^{n+1}} = \sum_{n=0}^{\infty} e^{-i(n+1)t} z^n$:

$$F_+(z) := F(z) = \sum_{n=0}^{\infty} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{-int} dt \right) z^n = \sum_{n=0}^{\infty} \frac{1}{\sqrt{2\pi}} \widehat{f}(n) z^n, \quad |z| < 1.$$

⁴¹Notice $\widehat{u}(n) = \sqrt{2\pi} \widehat{P_r}(n) \cdot \widehat{f}(n)$ and $\widehat{v}(n) = \sqrt{2\pi} \widehat{Q_r}(n) \cdot \widehat{f}(n)$.

For $|z| > 1$, expanding in $\frac{1}{w-z} = -\sum_{n=1}^{\infty} \frac{w^{n-1}}{z^n} = -\sum_{n=1}^{\infty} e^{i(n-1)t} z^{-n}$:

$$\begin{aligned} F_-(z) := F(z) &= -\sum_{n=1}^{\infty} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} f(t) e^{int} dt \right) z^{-n} = -\sum_{n=1}^{\infty} \frac{1}{\sqrt{2\pi}} \hat{f}(-n) z^{-n} \\ &= -\frac{1}{\sqrt{2\pi}} \sum_{n=-\infty}^{-1} \hat{f}(n) z^n, \quad |z| > 1. \end{aligned}$$

Thus by the definitions (1.27) and (1.28), $P_{\pm}(f)(x) = \lim_{r \rightarrow 1_{\mp}} F_{\pm}(re^{ix})$.

Almost everywhere convergence The L^p -convergence (1.23) does not imply the almost everywhere convergence⁴². It takes a long time for mathematicians to achieve the almost everywhere convergence result of Fourier series for the function $f \in \overline{L^p(-\pi, \pi)}$, $p \in (1, \infty]$.

- Case $p = 1$ is excluded, due to the existence of L^1 -functions whose Fourier series diverge everywhere, which is constructed by Andrey Nikolaevich Kolmogorov in 1926⁴³.
- Case $p = 2$ is included, thanks to Lennart Carleson's work in 1966⁴⁴. Since $L^p(-\pi, \pi) \subset L^2(-\pi, \pi)$ for all $p \in [2, \infty]$, the almost everywhere convergence holds for the cases $p \in [2, \infty]$.
- Case $p \in (1, 2)$ is included, thanks to R.A. Hunt's result in 1967⁴⁵.

Higher dimensional case We can of course the higher dimensional Fourier series. We denote $L_{\text{per}}^p(\mathbb{R}^d)$, $p \in [1, \infty]$ to be the space of measurable functions $f : \mathbb{R}^d \mapsto \mathbb{C}$ which are periodic such that $f \in L^p([-\pi, \pi]^d)$. We identify the two $L_{\text{per}}^p(\mathbb{R}^d)$ functions if they are the same almost everywhere. We define the k , $k = (k_1, \dots, k_d)$ -th Fourier coefficient of a function $f \in L_{\text{per}}^1(\mathbb{R}^d)$ by

$$\hat{f}(k) = \frac{1}{(2\pi)^{\frac{d}{2}}} \int_{[-\pi, \pi]^d} f(x) e^{-ik \cdot x} dx,$$

⁴²It holds only up to a subsequence.

⁴³Andrey Nikolaevich Kolmogorov, Une série de Fourier-Lebesgue divergente partout. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences (Paris)* 183, 1327-1329, 1926.

Andrey Nikolaevich Kolmogorov, Une série de Fourier-Lebesgue divergente presque partout. *Fundamenta Mathematicae* 4, 324-328, 1923.

⁴⁴Lennart Carleson, On convergence and growth of partial sums of Fourier series. *Acta Mathematica* 116, 135-157, 1966.

⁴⁵Richard A. Hunt, On the convergence of Fourier series. In *Orthogonal Expansions and their Continuous Analogues* (Proc. Conf., Edwardsville, Ill., 1967), Southern Illinois University Press, 235-255.

and we call the following sequence $(S_N f(x))_N$ the Fourier series of f :

$$S_N f(x) = \frac{1}{(2\pi)^{\frac{d}{2}}} \sum_{|k| \leq N} \hat{f}(k) e^{ik \cdot x}, \quad N \in \mathbb{N}.$$

From $[-\pi, \pi]$ to $\mathbb{R}$ Intuitively, we go from sums to integrals (by considering T -periodic functions with $T \rightarrow \infty$), to realize the limit of the Fourier series as the inverse Fourier transform of its Fourier transform.

Formally, for $d = 1$, let T be the period, and $\omega = \frac{2\pi}{T}$ be the frequency, then the Fourier coefficient of a function $f \in L^1([-\frac{T}{2}, \frac{T}{2}])$ is defined as

$$\hat{f}(k) = \frac{1}{\sqrt{T}} \int_{-\frac{T}{2}}^{\frac{T}{2}} f(x) e^{-ik\omega x} dx = \frac{\sqrt{\omega}}{\sqrt{2\pi}} \int_{-\frac{\pi}{\omega}}^{\frac{\pi}{\omega}} f(x) e^{-i(k\omega)x} dx,$$

and the representation of f in terms of Fourier series reads as

$$\begin{aligned} f(x) &= \frac{\sqrt{\omega}}{\sqrt{2\pi}} \sum_k \hat{f}(k) e^{i(k\omega)x} \\ &= \frac{1}{\sqrt{2\pi}} \sum_k \left(\frac{1}{\sqrt{2\pi}} \int_{-\frac{\pi}{\omega}}^{\frac{\pi}{\omega}} f(y) e^{-i(k\omega)y} dy \right) e^{i(k\omega)x} \omega. \end{aligned}$$

Let $\omega \rightarrow 0_+$, we formally obtain from the definition of integral that

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \left(\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(y) e^{-iy\xi} dy \right) e^{ix\xi} d\xi.$$

We then derive the continuous version of Fourier coefficients and Fourier series

$$\begin{aligned} \hat{f}(\xi) &= \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} f(x) e^{-ix\xi} dx, \quad \xi \in \mathbb{R}, \\ f(x) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \hat{f}(\xi) e^{ix\xi} d\xi, \quad x \in \mathbb{R}. \end{aligned}$$

This is the Fourier transform on $\mathbb{R}$, and we will discuss it later in Chapter 3 in more detail. To this end, we turn to the introduction of related functional spaces.

References

- [1] H. Bahouri, J.-Y. Chemin and R. Danchin: Fourier analysis and non-linear partial differential equations. Springer, 2011.
- [2] J. Duoandikoetxea: Fourier Analysis. Graduate Studies in Mathematics, Vol. 29. American Mathematical Society, 2001.
- [3] L. Grafakos: Classical Fourier Analysis. Springer, 2014.
- [4] M. Haragus, M.A. Johnson, W.R. Perkins, Linear Modulational and Subharmonic Dynamics of Spectrally Stable Lugiato-Lefever Periodic Waves, *J. Differential Equations*, 280 (2021), 315-354.
- [5] G. Teschl, Ordinary Differential Equations and Dynamical Systems, Version: July 22, 2009.