# Machine Learning

Answer Sheet for Homework 6

### Da-Min HUANG

#### R04942045

Graduate Institute of Communication Engineering, National Taiwan University

December 23, 2015

#### Problem 1

With the definition of  $z_n$ , rewrite the equation

$$\min_{A,B} F(A,B) = \frac{1}{N} \sum_{n=1}^{N} \ln \left( 1 + \exp \left( -y_n \left( A z_n + B \right) \right) \right) \tag{1}$$

So

$$\frac{\partial F}{\partial A} = \frac{1}{N} \sum_{n=1}^{N} -y_n \left( \frac{\exp(-y_n (Az_n + B))}{1 + \exp(-y_n (Az_n + B))} \right)^T z_n = \frac{1}{N} \sum_{n=1}^{N} -y_n p_n^T z_n$$
 (2)

$$\frac{\partial F}{\partial B} = \frac{1}{N} \sum_{n=1}^{N} -y_n \left( \frac{\exp(-y_n (Az_n + B))}{1 + \exp(-y_n (Az_n + B))} \right)^T = \frac{1}{N} \sum_{n=1}^{N} -y_n p_n^T$$
 (3)

Hence

$$\nabla F(A, B) = \frac{1}{N} \sum_{n=1}^{N} \left[ -y_n z_n p_n, -y_n p_n \right]^T$$
 (4)

#### Problem 2

Use the result of Problem 1 and define  $\exp(-y_n(Az_n+B)) = \exp(\xi_n)$ , we have

$$\frac{\partial^2 F}{\partial A^2} = \frac{\partial}{\partial A} \left( \frac{1}{N} \sum_{n=1}^N -y_n p_n^T z_n \right) \tag{5}$$

$$= \frac{1}{N} \sum_{n=1}^{N} -y_n \left( \frac{-y_n \exp(\xi_n) (1 + \exp(\xi_n)) z_n - y_n (\exp(\xi_n))^2 z_n}{(1 + \exp(\xi_n))^2} \right) z_n$$
 (6)

$$= \frac{1}{N} \sum_{n=1}^{N} -y_n \left( \frac{-y_n \exp(\xi_n) z_n}{(1 + \exp(\xi_n))^2} \right) z_n$$
 (7)

$$= \frac{1}{N} \sum_{n=1}^{N} (y_n)^2 \left( \frac{\exp(\xi_n)}{1 + \exp(\xi_n)} \left( 1 - \frac{\exp(\xi_n)}{1 + \exp(\xi_n)} \right) \right) z_n^2$$
 (8)

$$= \frac{1}{N} \sum_{n=1}^{N} z_n^2 p_n (1 - p_n) \tag{9}$$

where  $y_n^2 = 1$  since  $y_n \in \{-1, +1\}$ .

The other term is

$$\frac{\partial^2 F}{\partial A \partial B} = \frac{\partial}{\partial A} \left( \frac{1}{N} \sum_{n=1}^N -y_n p_n^T \right) = \frac{1}{N} \sum_{n=1}^N -y_n \left( \frac{-y_n \exp(\xi_n) z_n}{\left(1 + \exp(\xi_n)\right)^2} \right)$$
(10)

$$= \frac{1}{N} \sum_{n=1}^{N} z_n p_n (1 - p_n) \tag{11}$$

$$\frac{\partial^2 F}{\partial B \partial A} = \frac{\partial}{\partial B} \left( \frac{1}{N} \sum_{n=1}^N -y_n p_n^T z_n \right) = \frac{1}{N} \sum_{n=1}^N -y_n \left( \frac{-y_n \exp(\xi_n)}{\left(1 + \exp(\xi_n)\right)^2} \right) z_n \tag{12}$$

$$= \frac{1}{N} \sum_{n=1}^{N} z_n p_n (1 - p_n) \tag{13}$$

$$\frac{\partial^2 F}{\partial B^2} = \frac{\partial}{\partial B} \left( \frac{1}{N} \sum_{n=1}^N -y_n p_n^T \right) = \frac{1}{N} \sum_{n=1}^N -y_n \left( \frac{-y_n \exp\left(\xi_n\right)}{\left(1 + \exp\left(\xi_n\right)\right)^2} \right)$$
(14)

$$=\frac{1}{N}\sum_{n=1}^{N}p_{n}\left(1-p_{n}\right)\tag{15}$$

Hence, we have

$$H(F) = \frac{1}{N} \sum_{n=1}^{N} \begin{bmatrix} z_n^2 p_n (1 - p_n) & z_n p_n (1 - p_n) \\ z_n p_n (1 - p_n) & p_n (1 - p_n) \end{bmatrix}$$
(16)

#### Problem 3

As  $\gamma \to \infty$ , we have

$$\lim_{\gamma \to \infty} \exp\left(-\gamma \|\mathbf{x}_n - \mathbf{x}_m\|^2\right) = 0 \tag{17}$$

So K should be a zero matrix with size  $N \times N$ , which is  $\mathbf{0}_{N \times N}$ .

And  $\boldsymbol{\beta}$  is

$$\boldsymbol{\beta} = (\lambda I + K)^{-1} \mathbf{y} = \lambda^{-1} \mathbf{y}$$
(18)

#### Problem 4

If  $|y_n - \mathbf{w}^T \phi(\mathbf{x}_n) - b| > \epsilon$ , then

$$\begin{cases} |y_{n} - \mathbf{w}^{T} \phi(\mathbf{x}_{n}) - b| - \epsilon = \xi_{n}^{\wedge} \text{ and } \xi_{n}^{\vee} = 0, & \text{if } y_{n} - \mathbf{w}^{T} \phi(\mathbf{x}_{n}) - b > 0 \\ |y_{n} - \mathbf{w}^{T} \phi(\mathbf{x}_{n}) - b| - \epsilon = \xi_{n}^{\vee} \text{ and } \xi_{n}^{\wedge} = 0, & \text{if } y_{n} - \mathbf{w}^{T} \phi(\mathbf{x}_{n}) - b < 0 \end{cases}$$

$$(19)$$

and if  $|y_n - \mathbf{w}^T \phi(\mathbf{x}_n) - b| \le \epsilon$ , then  $\xi_n^{\wedge} = 0$  and  $\xi_n^{\vee} = 0$ . Hence, we have

$$\left(\xi_{n}^{\wedge}\right)^{2} + \left(\xi_{n}^{\vee}\right)^{2} = \left(\max\left(0, \left|y_{n} - \mathbf{w}^{T}\phi\left(\mathbf{x}_{n}\right) - b\right| - \epsilon\right)\right)^{2}$$
(20)

So  $P_2$  is equivalent to

$$\min_{b,\mathbf{w}} \left( \frac{1}{2} \mathbf{w}^T \mathbf{w} + C \sum_{n=1}^{N} \left( \max \left( 0, \left| y_n - \mathbf{w}^T \phi \left( \mathbf{x}_n \right) - b \right| - \epsilon \right) \right)^2 \right)$$
 (21)

with no constraints.

#### Problem 5

#### Problem 6

Problem 7	
Problem 8	
Problem 9	
Problem 10	
Problem 11	
Problem 12	
Problem 13	
Problem 14	

Problem 15	
Problem 16	
Problem 17	
Problem 18	
Problem 19	
Problem 20	

## Reference

[1] Lecture Notes by Hsuan-Tien LIN, Department of Computer Science and Information Engineering, National Taiwan University, Taipei 106, Taiwan.