
Machine Learning

Answer Sheet for Homework 7

Da-Min HUANG

R04942045

Graduate Institute of Communication Engineering, National Taiwan University

January 6, 2016

Problem 1

Set $\mu_- = 1 - \mu_+$, we have

$$1 - \mu_+^2 - \mu_-^2 = 1 - \mu_+^2 - (1 - \mu_+)^2 = (1 - \mu_+)(1 + \mu_+) - (1 - \mu_+)^2 \quad (1)$$

$$= 2\mu_+(1 - \mu_+) = -2\mu_+^2 + 2\mu_+ = -2\left(\mu_+ - \frac{1}{2}\right)^2 + \frac{1}{2} \quad (2)$$

$$\leq \frac{1}{2} \quad (3)$$

Hence, if $\mu_+ = 1/2 \in [0, 1]$, then the maximum value of Gini index is $1/2$.

□

Problem 2

The normalized Gini index is

$$\frac{(1 - \mu_+^2 - \mu_-^2)}{\left(\frac{1}{2}\right)} = 2(1 - \mu_+^2 - \mu_-^2) \quad (4)$$

The squared error can be rewritten as

$$\mu_+(1 - (\mu_+ - \mu_-))^2 + \mu_-(-1 - (\mu_+ - \mu_-))^2 = 4\mu_+(1 - \mu_+)^2 + 4\mu_+^2(1 - \mu_+) \quad (5)$$

$$= 4\mu_+(1 - \mu_+) \leq 4 \times \frac{1}{4} = 1 \quad (6)$$

Hence the normalized squared error is

$$4\mu_+(1 - \mu_+) = 2(2\mu_+(1 - \mu_+)) = 2((1 - \mu_+)(1 + \mu_+) - (1 - \mu_+)^2) \quad (7)$$

$$= 2(1 - \mu_+^2 - \mu_-^2) \quad (8)$$

which is equal to normalized Gini index.

□

Problem 3

The probability of one example not sampled is

$$\left(1 - \frac{1}{N}\right)^{pN} = \frac{1}{\left(\frac{N}{N-1}\right)^{pN}} = \frac{1}{\left(1 + \frac{1}{N-1}\right)^{pN}} = \left(\frac{1}{\left(1 + \frac{1}{N-1}\right)^N}\right)^p \quad (9)$$

As $N \rightarrow \infty$, we have

$$\lim_{N \rightarrow \infty} \left(\frac{1}{\left(1 + \frac{1}{N-1}\right)^N}\right)^p = \left(\lim_{N \rightarrow \infty} \frac{1}{\left(1 + \frac{1}{N-1}\right)^N}\right)^p = \left(\frac{1}{e}\right)^p = e^{-p} \quad (10)$$

So there approximately $e^{-p} \cdot N$ of the examples not sampled.

□

Problem 4

Since $G = \text{Uniform}(\{g_k\}_{k=1}^3)$, so if at least two terms of $\{g_k\}_{k=1}^3$ output wrong result, then G outputs wrong result. Let $\{E_k\}_{k=1}^3$ be the set of examples that $\{g_k\}_{k=1}^3$ got wrong results. Apparently $|E_3| > |E_2| > |E_1|$ and $|E_1| + |E_2| > |E_3|$. So

1. Maximum of $E_{\text{out}}(G)$ happens at $E_3 \subset (E_1 \cup E_2)$. Then G outputs wrong result in the region of E_3 with $E_{\text{out}}(G) = 0.35$.
2. Minimum of $E_{\text{out}}(G)$ happens at $E_i \cap E_j = \emptyset$, $i \neq j$ and $1 \leq i, j \leq 3$ with $i, j \in \mathbb{N}$.
Then G always outputs the correct result since $(E_1 \cup E_2 \cup E_3) \subset \{\text{all examples}\}$.

Hence, $0 \leq E_{\text{out}}(G) \leq 0.35$.

□

Problem 5

Since $G = \text{Uniform}(\{g_k\}_{k=1}^K)$, so if at least $(K+1)/2$ terms of $\{g_k\}_{k=1}^K$ output wrong result, then G outputs wrong result. Let $\{E_k\}_{k=1}^K$ be the set of examples that $\{g_k\}_{k=1}^K$ got wrong results.

If G outputs wrong result on some example \mathbf{x} , then we have

$$\mathbf{x} \in \bigcap_{i=1}^{((K+1)/2)+m} E_{\alpha_i} \quad (11)$$

where α_i is some index satisfies $1 \leq \alpha_i \leq K$ and $m \in (\mathbb{N} \cup \{0\})$ with $0 \leq m < (K+1)/2$. And

$$\left| \bigcap_{i=1}^{((K+1)/2)+m} E_{\alpha_i} \right| \leq \frac{2}{K+1+2m} \sum_{k=1}^K e_k \leq \frac{2}{K+1} \sum_{k=1}^K e_k \quad (12)$$

(12) holds due to

$$\bigcap_{i=1}^{((K+1)/2)+m} E_{\alpha_i} \subseteq E_{\beta} \quad (13)$$

where β is some index such that $|E_{\beta}| = \min_{\alpha_i} |E_{\alpha_i}|$. So

$$\left(\frac{K+1}{2} + m \right) \left| \bigcap_{i=1}^{((K+1)/2)+m} E_{\alpha_i} \right| \leq \left(\frac{K+1}{2} + m \right) |E_{\beta}| \leq \sum_{k=1}^K |E_k| \quad (14)$$

(14) holds since size of E_{β} is the smallest among $((K+1)/2 + m)$ terms and $\sum_{k=1}^K |E_k|$ must contains the $((K+1)/2 + m)$ terms.

Hence, we have

$$E_{\text{out}}(G) \leq \frac{2}{K+1} \sum_{k=1}^K e_k \quad (15)$$

□

Problem 6

By the definition of U_t , we have

$$U_{t+1} = \frac{1}{N} \sum_{n=1}^N \exp \left(-y_n \sum_{\tau=1}^t \alpha_\tau g_\tau (\mathbf{x}_n) \right) \quad (16)$$

$$= \frac{1}{N} \sum_{n=1}^N \exp \left(-y_n \sum_{\tau=1}^{t-1} \alpha_\tau g_\tau (\mathbf{x}_n) - y_n \alpha_t g_t (\mathbf{x}_n) \right) \quad (17)$$

$$= \frac{1}{N} \sum_{n=1}^N \exp \left(-y_n \sum_{\tau=1}^{t-1} \alpha_\tau g_\tau (\mathbf{x}_n) \right) \exp (-y_n \alpha_t g_t (\mathbf{x}_n)) \quad (18)$$

$$= \sum_{n=1}^N u_n^{(t)} \exp (-y_n \alpha_t g_t (\mathbf{x}_n)) \quad (19)$$

$$= \sum_{\substack{n \\ y_n \neq g_t (\mathbf{x}_n)}} u_n^{(t)} \exp (-y_n \alpha_t g_t (\mathbf{x}_n)) + \sum_{\substack{n \\ y_n = g_t (\mathbf{x}_n)}} u_n^{(t)} \exp (-y_n \alpha_t g_t (\mathbf{x}_n)) \quad (20)$$

$$= \sum_{\substack{n \\ y_n \neq g_t (\mathbf{x}_n)}} u_n^{(t)} \exp (\alpha_t) + \sum_{\substack{n \\ y_n = g_t (\mathbf{x}_n)}} u_n^{(t)} \exp (-\alpha_t) \quad (21)$$

$$= \exp (\alpha_t) (\epsilon_t) \sum_{n=1}^N u_n^{(t)} + \exp (-\alpha_t) (1 - \epsilon_t) \sum_{n=1}^N u_n^{(t)} \quad (22)$$

$$= U_t (\exp (\alpha_t) (\epsilon_t) + \exp (-\alpha_t) (1 - \epsilon_t)) = U_t \cdot 2\sqrt{\epsilon_t (1 - \epsilon_t)} \quad (23)$$

Since

$$U_1 = \sum_{n=1}^N u_n^{(1)} = \sum_{n=1}^N \frac{1}{N} = 1 \quad (24)$$

we have

$$U_3 = U_2 \cdot 2\sqrt{\epsilon_2 (1 - \epsilon_2)} = \left(U_1 \cdot 2\sqrt{\epsilon_1 (1 - \epsilon_1)} \right) \cdot 2\sqrt{\epsilon_2 (1 - \epsilon_2)} \quad (25)$$

$$= 4\sqrt{\epsilon_1 \epsilon_2 (1 - \epsilon_1) (1 - \epsilon_2)} \quad (26)$$

which can be generalized as

$$U_{T+1} = \prod_{t=1}^T \left(2\sqrt{\epsilon_t (1 - \epsilon_t)} \right) \quad (27)$$

□

Problem 7

To compute s_n , we need to find the optimal η of

$$\min_{\eta} \frac{1}{N} \sum_{n=1}^N ((y_n - s_n) - \eta g_t(\mathbf{x}_n))^2 := A \quad (28)$$

From $\partial A / \partial \eta = 0$, we have

$$\eta = \frac{\sum_{n=1}^N g_t(\mathbf{x}_n) (y_n - s_n)}{\sum_{n=1}^N g_t^2(\mathbf{x}_n)} \quad (29)$$

Now $s_n = 0$ and $g_1(\mathbf{x}) = 2$, so

$$\eta = \frac{2 \sum_{n=1}^N y_n}{4 \sum_{n=1}^N 1} = \frac{1}{2N} \sum_{n=1}^N y_n \quad (30)$$

Since $\eta = \alpha_1$, so

$$\alpha_1 g_1(\mathbf{x}_n) = \frac{2}{2N} \sum_{n=1}^N y_n = \frac{1}{N} \sum_{n=1}^N y_n = s_n \quad (31)$$

□

Problem 8

From the equatio of optimal η , we have

$$\eta = \frac{\sum_{n=1}^N g_t(\mathbf{x}_n) (y_n - s'_n)}{\sum_{n=1}^N g_t^2(\mathbf{x}_n)} = \frac{\sum_{n=1}^N y_n g_t(\mathbf{x}_n) - \sum_{n=1}^N s'_n g_t(\mathbf{x}_n)}{\sum_{n=1}^N g_t^2(\mathbf{x}_n)} = \alpha_t \quad (32)$$

so

$$\sum_{n=1}^N y_n g_t(\mathbf{x}_n) - \sum_{n=1}^N s'_n g_t(\mathbf{x}_n) = \alpha_t \sum_{n=1}^N g_t^2(\mathbf{x}_n) = \sum_{n=1}^N \alpha_t g_t^2(\mathbf{x}_n) = \sum_{n=1}^N (s_n - s'_n) g_t(\mathbf{x}_n) \quad (33)$$

where s'_n is defined as the s_n in iteration $(t - 1)$, so

$$\sum_{n=1}^N s_n g_t(\mathbf{x}_n) = \sum_{n=1}^N y_n g_t(\mathbf{x}_n) \quad (34)$$

□

Problem 9

☐

Problem 10

☐

Problem 11

☐

Problem 12

☐

Problem 13

☐

Problem 14

☐

Problem 15

☐

Problem 16

☐

Problem 17



Problem 18



Problem 19



Problem 20



Reference

- [1] Lecture Notes by Hsuan-Tien LIN, Department of Computer Science and Information Engineering, National Taiwan University, Taipei 106, Taiwan.