#### CS 419M Introduction to Machine Learning

Spring 2021-22

Lecture 7: Classification

Lecturer: Abir De Scribe: Group 2

### 7.1 Topics covered in lecture

1. What is classification task

2. What are the classification models

3. Support Vector Machine (not Covered in lecture)

#### 7.2 Classification task

• Given training set  $\mathcal{D} = \{(x_i, y_i) \mid y_i \in \mathcal{G}\}, \mathcal{G} = \{y_1, y_2\}, x_i \in \mathbb{R}^d, \text{ find } m(x) \mapsto y.$ 

• Test set =  $\{x_i \in \mathbb{R}^d\}$ ,  $y_i$  are not known in test set.

### 7.3 Probabilistic Approch

$$P_{m}(y \mid x) = \frac{1}{1 + e^{-w^{T}x \cdot y}}$$

$$\implies \max_{w} \prod_{i \in \mathcal{D}} P_{m}(y_{i} \mid x_{i})$$

$$\implies \max_{w} \sum_{i \in \mathcal{D}} \log P_{m}(y_{i} \mid x_{i})$$

$$\implies \min_{w} \sum_{i \in mathcalD} \log \left(1 + e^{w^{T}x_{i} \cdot y_{i}}\right)$$

# 7.4 Simpler way to classify

Let  $\mathcal{G} = \{+1, -1\}$ , for some other labels we can convert that to +1, -1. We want some linear boundary to classify given points into  $\mathcal{G}$ .

$$w^{T}x + b \ge \Delta, \quad y = 1$$
  
$$w^{T}x + b \le -\Delta, \quad y = -1$$
  
$$\min_{w} f(w)$$

where f(w) is any regularizer. This problem can be solved when there is no overlapping between the classification variables but there isn't a solution for something like This




Figure 7.1: no overlapping




Figure 7.2: Overlapping

We can ignore the overlapping points,

Let 
$$I^+ = \{i \mid y_i = 1\}, I^- = \{i \mid y_i = -1\}, S^+ \in I^+, S^- \in I^- \text{ such that } |S^+ \cup S^-| = n,$$

$$\min_{\zeta_i} f(w) - \left( \sum_{i \in S^+} \mathbb{I}\left( w^T x_i + b \ge \Delta \right) + \sum_{i \in S^-} \mathbb{I}\left( w^T x_i + b \le \Delta \right) \right)$$

### 7.5 Adding Slack variable to solve the overlapping case

Modifying optimisation problem to include overlapping points,

$$w^{T}x_{i} + b \ge \Delta - \zeta_{i}, \quad y_{i} = 1$$
$$w^{T}x_{i} + b \le -\Delta + \zeta_{i}, \quad y_{i} = -1$$
$$y_{i} \cdot (w^{T}x_{i} + b) \ge \Delta - \zeta_{i}$$
$$\zeta_{i} \ge 0$$

with above conditions, we have to solve following

$$\min_{w,b,\zeta_i} C \sum_{i \in \mathcal{D}} \zeta_i + \lambda \mid\mid w \mid\mid^2$$

## 7.6 Group Details and Individual Contribution

Name	Roll number	contribution
Dadhichi Telwadkar	20D070083	7.1,7.2,7.3,7.4,7.5
Sameer Parwani	20D180029	7.1,7.2,7.3,7.4,7.5
Pratham Kingar	200040075	7.1,7.2,7.3,7.4,7.5
Samyak Ajmera	19B080023	7.1,7.2,7.3,7.4,7.5
Aakash Jha	200040002	7.1,7.2,7.3,7.4,7.5