

EXISTENCE AND SIMULATION OF GIBBS-LAGUERRE-DELAUNAY TETRAHEDRIZATIONS

DANIEL JAHN

We present revolutionary unprecedented results using newly developed methods. Songs will be sung about this paper.

Keywords: Gibbs point process, Laguerre-Delaunay triangulation, tetrahedralization, MCMC simulation

Classification: 60K35, 60G55

1. INTRODUCTION

This is an interesting field. We did some things in this interesting field. Here they are.

2. PRELIMINARIES

See [1].

- Laguerre geometry
- Gibbs point process - already tie in [2].

3. EXISTENCE

Basically [2].

- Specification of the model
- Proof of existence

Smooth interaction: For $\eta \in \mathcal{LD}(\mathbf{x})$ define the potential φ_S as a unary potential such that

$$\varphi_S(\eta, \mathbf{x}) \leq K_0 + K_1 \chi(\eta)^\beta$$

for some $K_0, K_1 \geq 0, \beta > 0$

Hard-core interaction: For $\eta \in \mathcal{LD}(\mathbf{x})$ define the potential φ_{HC} as a unary potential such that

$$\sup_{\eta: d_0 \leq \chi(\eta) \leq d_1} \varphi_{HC}(\eta, \mathbf{x}) < \infty \text{ and } \varphi_{HC}(\eta, \mathbf{x}) = \infty \text{ if } \chi(\eta) > \alpha$$

How exactly
does this
look? Why?

for some $0 \leq d_0 < d_1 \leq \alpha$.

3.1. Existence theorems

Theorem 3.1. For every hypergraph structure \mathcal{E} , hyperedge potential φ and activity $z > 0$ satisfying **(S)**, **(R)** and **(U)** there exists at least one Gibbs measure.

Theorem 3.2. For every hypergraph structure \mathcal{E} , hyperedge potential φ and activity $z > 0$ satisfying **(S)**, **(R)** and **(Ū)** there exists at least one Gibbs measure.

Proofs of both theorems can be found in [2], see also Remark 3.7. in the same paper about the marked case.

(S) Stability. The energy function H is called *stable* if there exists a constant $c_S \geq 0$ such that

$$H_{\Lambda, \mathbf{x}}(\zeta) \geq -c_S \cdot \text{card}(\zeta \cup \partial_{\Lambda} \mathbf{x})$$

for all $\Lambda \in \mathcal{B}_0, \zeta \in \mathbf{N}_{\Lambda}, \mathbf{x} \in \mathbf{N}_{\text{cr}}^{\Lambda}$.

(R) Range condition. There exist constants $\ell_R, n_R \in \mathbb{N}$ and $\chi_R < \infty$ such that for all $(\eta, \mathbf{x}) \in \mathcal{E}$ there exists a finite horizon Δ satisfying: For every $x, y \in \Delta$ there exist ℓ open balls B_1, \dots, B_{ℓ} (with $\ell \leq \ell_R$) such that

- the set $\cup_{i=1}^{\ell} \bar{B}_i$ is connected and contains x and y , and
- for each i , either $\text{diam} B_i \leq \chi_R$ or $N_{B_i}(\mathbf{x}) \leq n_R$.

(U) Upper regularity. M and Γ can be chosen so that the following holds.

(U1) *Uniform confinement:* $\bar{\Gamma} \subset \mathbf{N}_{\text{cr}}^{\Lambda}$ for all $\Lambda \in \mathcal{B}_0$ and

$$r_{\Gamma} := \sup_{\Lambda \in \mathcal{B}_0} \sup_{\mathbf{x} \in \bar{\Gamma}} r_{\Lambda, \mathbf{x}} < \infty. \quad (1)$$

(U2) *Uniform summability:*

$$c_{\Gamma} := \sup_{\mathbf{x} \in \bar{\Gamma}} \sum_{\eta \in \mathcal{E}(\mathbf{x}) : \eta' \cap C \neq \emptyset} \frac{\varphi(\eta, \mathbf{x})}{\#(\hat{\eta})} < \infty,$$

where $\hat{\eta} := \{k \in \mathbb{Z}^3 : \eta \cap C(k) \neq \emptyset\}$.

(U3) *Strong non-rigidity:* $e^{z|C|} \Pi_{\bar{C}}^z(\Gamma) > e^{c_{\Gamma}}$.

For some models it is possible to replace the upper regularity assumptions by their alternative and prove the existence for all $z > 0$.

(Ū) Alternative upper regularity. M and Γ can be chosen so that the following holds.

(Ū1) *Lower density bound:* There exist constants $c, d > 0$ such that

$$\text{card}(\zeta) \geq c|\Lambda| - d$$

whenever $\zeta \in \mathbf{N}_f \cap \mathbf{N}_{\Lambda}$ is such that $H_{\Lambda, \mathbf{x}}(\zeta) < \infty$ for some $\Lambda \in \mathcal{B}_0$ and some $\mathbf{x} \in \bar{\Gamma}$.

(Ū2) = (U2) *Uniform summability.*

(Ū3) *Weak non-rigidity:* $\Pi_{\bar{C}}^z(\Gamma) > 0$.

3.2. Verification

Fix some $A \subset C \times S$ and define

$$\Gamma^b = \{\zeta \in \mathbf{N}_C : \zeta = \{p\}, p \in B(0, b) \times \left[0, \sqrt{\frac{a}{2}(1-2\rho)}\right]\},$$

the set of configurations consisting of exactly one point in the set A . The set of pseudo-periodic configurations $\bar{\Gamma}$ thus contains only one point in each $C(k), k \in \mathbb{Z}^3$.

Let M be such that $|M_i| = a > 0$ for $i = 1, 2, 3$ and $\angle(M_i, M_j) = \pi/3$ for $i \neq j$. We further assume $\rho < 1/4$, see Appendix ??.

4. SIMULATION

Basically [3].

- Specification of the simulated model.
- Results

ACKNOWLEDGEMENT

This work was partially supported by GACR 17-00393J (investigator V. Beneš). This work was fully supported by Panda.

(Received ????)

REFERENCES

-
- [1] D. Dereudre: Introduction to the theory of Gibbs point processes. ArXiv 1701.08105 [math.PR] (2017).
 - [2] D. Dereudre, R. Drouilhet, H.O. Georgii: Existence of Gibbsian point processes with geometry-dependent interactions. Probability Theory and Related Fields 153 (3) (2012), 643–670.
 - [3] D. Dereudre and F. Lavancier: Practical simulation and estimation for Gibbs Delaunay-Voronoi tessellations with geometric hardcore interaction. Computational Statistics and Data Analysis 55 (1) (2011), 498–519.

Daniel Jahn, Department of Probability and Mathematical Statistics, Charles University, Sokolovská 83, 186 75 Praha 8, Czech Republic
e-mail: jahn@karlin.mff.cuni.cz