

# $P$ -adic Borel hyperbolicity of $A_g$

Xinwen Zhu

Stanford University

PKU Mathematics Forum

August 2, 2023

# Picard's theorem

## Theorem (Fundamental theorem of algebra)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant polynomial function. Then  $f(\mathbb{C}) = \mathbb{C}$ .*

# Picard's theorem

## Theorem (Fundamental theorem of algebra)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant polynomial function. Then  $f(\mathbb{C}) = \mathbb{C}$ .*

## Theorem (Little Picard theorem)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant entire function. Then  $\#(\mathbb{C} - f(\mathbb{C})) \leq 1$ .*

# Picard's theorem

## Theorem (Fundamental theorem of algebra)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant polynomial function. Then  $f(\mathbb{C}) = \mathbb{C}$ .*

## Theorem (Little Picard theorem)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant entire function. Then  $\#(\mathbb{C} - f(\mathbb{C})) \leq 1$ .*

Now, for  $z_0 \in \mathbb{C}$  and  $r \in \mathbb{R}_{>0}$ , let

$$D(z_0, r) = \{z \in \mathbb{C} \mid |z - z_0| < r\}, \quad D(z_0, r)^\times = D(z_0, r) - \{z_0\}.$$

# Picard's theorem

## Theorem (Fundamental theorem of algebra)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant polynomial function. Then  $f(\mathbb{C}) = \mathbb{C}$ .*

## Theorem (Little Picard theorem)

*Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be a non-constant entire function. Then  $\sharp(\mathbb{C} - f(\mathbb{C})) \leq 1$ .*

Now, for  $z_0 \in \mathbb{C}$  and  $r \in \mathbb{R}_{>0}$ , let

$$D(z_0, r) = \{z \in \mathbb{C} \mid |z - z_0| < r\}, \quad D(z_0, r)^\times = D(z_0, r) - \{z_0\}.$$

## Theorem (Great Picard theorem)

*Let  $f : D(0, r)^\times \rightarrow \mathbb{C}$  be a holomorphic function with essential singularity at 0, then  $\sharp(\mathbb{C} - f(D(0, r)^\times)) \leq 1$ .*

Note that the exponential function  $\exp : \mathbb{C} \rightarrow \mathbb{C}^\times := \mathbb{C} - \{0\}$  shows that Picard's theorems are sharp.

# Hyperbolic Riemann surfaces

Recall the Riemann sphere, also known as 1-dim. projective space, is

$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\} = (\mathbb{C}^2 - \{(0, 0)\}) / (z_1, z_2) \sim (\lambda z_1, \lambda z_2).$$

# Hyperbolic Riemann surfaces

Recall the Riemann sphere, also known as 1-dim. projective space, is

$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\} = (\mathbb{C}^2 - \{(0,0)\}) / (z_1, z_2) \sim (\lambda z_1, \lambda z_2).$$

Picard's theorems are equivalent to

## Theorem

- 1 Every holomorphic map  $f : \mathbb{C} \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  is constant.
- 2 Every holomorphic map  $f : D(0, r)^\times \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  extends to a holomorphic map  $\tilde{f} : D(0, r) \rightarrow \mathbb{P}^1$ .

# Hyperbolic Riemann surfaces

Recall the Riemann sphere, also known as 1-dim. projective space, is

$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\} = (\mathbb{C}^2 - \{(0,0)\}) / (z_1, z_2) \sim (\lambda z_1, \lambda z_2).$$

Picard's theorems are equivalent to

## Theorem

- ① *Every holomorphic map  $f : \mathbb{C} \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  is constant.*
- ② *Every holomorphic map  $f : D(0, r)^\times \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  extends to a holomorphic map  $\tilde{f} : D(0, r) \rightarrow \mathbb{P}^1$ .*

A Riemann surface  $\Sigma$  is called hyperbolic if it can be uniformized by  $D(0, 1)$ , i.e.  $\exists$  a holomorphic covering map  $D(0, 1) \rightarrow \Sigma$ .



# Hyperbolic Riemann surfaces

Recall the Riemann sphere, also known as 1-dim. projective space, is

$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\} = (\mathbb{C}^2 - \{(0,0)\}) / (z_1, z_2) \sim (\lambda z_1, \lambda z_2).$$

Picard's theorems are equivalent to

## Theorem

- ① Every holomorphic map  $f : \mathbb{C} \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  is constant.
- ② Every holomorphic map  $f : D(0, r)^\times \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  extends to a holomorphic map  $\tilde{f} : D(0, r) \rightarrow \mathbb{P}^1$ .

A Riemann surface  $\Sigma$  is called hyperbolic if it can be uniformized by  $D(0, 1)$ , i.e.  $\exists$  a holomorphic covering map  $D(0, 1) \rightarrow \Sigma$ . E.g.

- $\mathbb{P}^1 - \{0, 1, \infty\}$ ;
- Compact Riemann surfaces of genus  $g \geq 2$ .

# Hyperbolic Riemann surfaces

Recall the Riemann sphere, also known as 1-dim. projective space, is

$$\mathbb{P}^1 = \mathbb{C} \cup \{\infty\} = (\mathbb{C}^2 - \{(0,0)\}) / (z_1, z_2) \sim (\lambda z_1, \lambda z_2).$$

Picard's theorems are equivalent to

## Theorem

- ① *Every holomorphic map  $f : \mathbb{C} \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  is constant.*
- ② *Every holomorphic map  $f : D(0, r)^\times \rightarrow \mathbb{P}^1 - \{0, 1, \infty\}$  extends to a holomorphic map  $\tilde{f} : D(0, r) \rightarrow \mathbb{P}^1$ .*

A Riemann surface  $\Sigma$  is called hyperbolic if it can be uniformized by  $D(0, 1)$ , i.e.  $\exists$  a holomorphic covering map  $D(0, 1) \rightarrow \Sigma$ . E.g.

- $\mathbb{P}^1 - \{0, 1, \infty\}$ ;
- Compact Riemann surfaces of genus  $g \geq 2$ .

The above form of Picard's theorem holds for all hyperbolic Riemann surfaces.

# Algebraic varieties

A complex algebraic variety is a space *locally* defined as the set of zeros of a system of polynomial equations in  $\mathbb{C}^n$ . E.g.

# Algebraic varieties

A complex algebraic variety is a space *locally* defined as the set of zeros of a system of polynomial equations in  $\mathbb{C}^n$ . E.g.

- Affine  $n$ -space  $\mathbb{C}^n$ , also denoted by  $\mathbb{A}^n$ .
- $\mathbb{C}^\times = \{(x, y) \in \mathbb{C}^2 \mid xy = 1\}$ , also denoted by  $\mathbb{G}_m$ .
- (Partial) flag varieties  $\mathcal{F}\ell$  parameterizing chains of subspaces in  $\mathbb{C}^n$  with given dimensions. A special case is the projective  $n$ -space  $\mathbb{P}^n = (\mathbb{C}^{n+1} - \{(0, \dots, 0)\})/\mathbb{G}_m$ , parameterizing lines in  $\mathbb{C}^{n+1}$ .
- $\Sigma \setminus \{z_1, \dots, z_r\}$  with  $\Sigma$  compact Riemann surfaces, called algebraic curves. E.g.  $(\mathbb{C} - \Lambda)/\Lambda \cong \{(x, y) \in \mathbb{C}^2 \mid 4y^2 = x^3 - g_2x - g_3\}$ .
- Abelian varieties  $A = \mathbb{C}^n/\Lambda$  by certain lattices.

# Algebraic varieties

A complex algebraic variety is a space *locally* defined as the set of zeros of a system of polynomial equations in  $\mathbb{C}^n$ . E.g.

- Affine  $n$ -space  $\mathbb{C}^n$ , also denoted by  $\mathbb{A}^n$ .
- $\mathbb{C}^\times = \{(x, y) \in \mathbb{C}^2 \mid xy = 1\}$ , also denoted by  $\mathbb{G}_m$ .
- (Partial) flag varieties  $\mathcal{F}\ell$  parameterizing chains of subspaces in  $\mathbb{C}^n$  with given dimensions. A special case is the projective  $n$ -space  $\mathbb{P}^n = (\mathbb{C}^{n+1} - \{(0, \dots, 0)\})/\mathbb{G}_m$ , parameterizing lines in  $\mathbb{C}^{n+1}$ .
- $\Sigma \setminus \{z_1, \dots, z_r\}$  with  $\Sigma$  compact Riemann surfaces, called algebraic curves. E.g.  $(\mathbb{C} - \Lambda)/\Lambda \cong \{(x, y) \in \mathbb{C}^2 \mid 4y^2 = x^3 - g_2x - g_3\}$ .
- Abelian varieties  $A = \mathbb{C}^n/\Lambda$  by certain lattices.

On the other hand,  $D(0, r)$  and  $D(0, r)^\times$  are not algebraic.

# Algebraic varieties

A complex algebraic variety is a space *locally* defined as the set of zeros of a system of polynomial equations in  $\mathbb{C}^n$ . E.g.

- Affine  $n$ -space  $\mathbb{C}^n$ , also denoted by  $\mathbb{A}^n$ .
- $\mathbb{C}^\times = \{(x, y) \in \mathbb{C}^2 \mid xy = 1\}$ , also denoted by  $\mathbb{G}_m$ .
- (Partial) flag varieties  $\mathcal{F}\ell$  parameterizing chains of subspaces in  $\mathbb{C}^n$  with given dimensions. A special case is the projective  $n$ -space  $\mathbb{P}^n = (\mathbb{C}^{n+1} - \{(0, \dots, 0)\})/\mathbb{G}_m$ , parameterizing lines in  $\mathbb{C}^{n+1}$ .
- $\Sigma \setminus \{z_1, \dots, z_r\}$  with  $\Sigma$  compact Riemann surfaces, called algebraic curves. E.g.  $(\mathbb{C} - \Lambda)/\Lambda \cong \{(x, y) \in \mathbb{C}^2 \mid 4y^2 = x^3 - g_2x - g_3\}$ .
- Abelian varieties  $A = \mathbb{C}^n/\Lambda$  by certain lattices.

On the other hand,  $D(0, r)$  and  $D(0, r)^\times$  are not algebraic.

The notion of algebraic varieties makes sense by replacing  $\mathbb{C}$  by any field  $k$ . In addition, if  $k \rightarrow k'$  is a field extension, an algebraic variety over  $k$  gives an algebraic variety over  $k'$ .

# Analytic v.s. Algebraic

Polynomial/algebraic maps between algebraic varieties induce holomorphic maps between complex analytic spaces. It is natural to ask the converse: which holomorphic maps between complex analytic spaces are induced from algebraic maps between algebraic varieties.

# Analytic v.s. Algebraic

Polynomial/algebraic maps between algebraic varieties induce holomorphic maps between complex analytic spaces. It is natural to ask the converse: which holomorphic maps between complex analytic spaces are induced from algebraic maps between algebraic varieties.

## Theorem (Chow)

*Every closed analytic subspaces in  $\mathbb{P}^n$  is algebraic.*



# Analytic v.s. Algebraic

Polynomial/algebraic maps between algebraic varieties induce holomorphic maps between complex analytic spaces. It is natural to ask the converse: which holomorphic maps between complex analytic spaces are induced from algebraic maps between algebraic varieties.

## Theorem (Chow)

*Every closed analytic subspaces in  $\mathbb{P}^n$  is algebraic.*

The great Picard theorem and Chow's theorem together  $(+\cdots)$  imply

## Theorem

*Every holomorphic map  $f : S \rightarrow X$  from an algebraic variety to a hyperbolic algebraic curve is algebraic.*

# Analytic v.s. Algebraic

Polynomial/algebraic maps between algebraic varieties induce holomorphic maps between complex analytic spaces. It is natural to ask the converse: which holomorphic maps between complex analytic spaces are induced from algebraic maps between algebraic varieties.

## Theorem (Chow)

*Every closed analytic subspace in  $\mathbb{P}^n$  is algebraic.*

The great Picard theorem and Chow's theorem together  $(+\cdots)$  imply

## Theorem

*Every holomorphic map  $f : S \rightarrow X$  from an algebraic variety to a hyperbolic algebraic curve is algebraic.*

This algebraicity theorem in turn implies the little Picard theorem. (Hint: if both  $f : \mathbb{C} \rightarrow X$  and  $f \circ \exp$  are polynomial maps, then  $f$  is constant.)

# Hyperbolicity in higher dimensions

The notion of hyperbolicity for higher dimensional algebraic varieties has connections with different branches of mathematics: complex analysis, differential geometry, algebraic geometry and diophantine geometry.

# Hyperbolicity in higher dimensions

The notion of hyperbolicity for higher dimensional algebraic varieties has connections with different branches of mathematics: complex analysis, differential geometry, algebraic geometry and diophantine geometry.

## Definition

A complex algebraic variety is called Brody hyperbolic if every holomorphic map  $f : \mathbb{C} \rightarrow X$  is constant.

# Hyperbolicity in higher dimensions

The notion of hyperbolicity for higher dimensional algebraic varieties has connections with different branches of mathematics: complex analysis, differential geometry, algebraic geometry and diophantine geometry.

## Definition

A complex algebraic variety is called Brody hyperbolic if every holomorphic map  $f : \mathbb{C} \rightarrow X$  is constant.

When  $X$  is compact, this is equivalent to the so-called Kobayashi hyperbolicity, defined using metric and distance.

# Hyperbolicity in higher dimensions

The notion of hyperbolicity for higher dimensional algebraic varieties has connections with different branches of mathematics: complex analysis, differential geometry, algebraic geometry and diophantine geometry.

## Definition

A complex algebraic variety is called Brody hyperbolic if every holomorphic map  $f : \mathbb{C} \rightarrow X$  is constant.

When  $X$  is compact, this is equivalent to the so-called Kobayashi hyperbolicity, defined using metric and distance.

## Conjecture (Green-Griffith-Lang)

*Assume that  $X$  is compact. Then TFAE:*

- *$X$  is Brody hyperbolic;*
- *Every closed subvariety of  $X$  is of general type;*
- *There is no non-constant rational map from an abelian variety to  $X$ .*

# Arithmetic manifolds

Let  $D$  be a bounded symmetric domain in a complex vector space, also known as hermitian symmetric domain. It is of the form

$$D = G/K$$

where  $G$  is the group of holomorphic automorphism of  $D$ , which is a real (semisimple) Lie group and  $K$  is a maximal compact subgroup of  $G$ .

# Arithmetic manifolds

Let  $D$  be a bounded symmetric domain in a complex vector space, also known as hermitian symmetric domain. It is of the form

$$D = G/K$$

where  $G$  is the group of holomorphic automorphism of  $D$ , which is a real (semisimple) Lie group and  $K$  is a maximal compact subgroup of  $G$ . E.g.

- $D(0, 1) \cong \mathfrak{H} = \{z \in \mathbb{C} \mid \text{Im}z > 0\} \cong \text{SL}_2(\mathbb{R})/\text{SO}_2$ ;
- $\mathfrak{H}_g = \{Z \in M_g(\mathbb{C}) \mid Z = Z^T, \text{Im}Z > 0\} \cong \text{Sp}_{2g}(\mathbb{R})/\text{U}_g$ .



# Arithmetic manifolds

Let  $D$  be a bounded symmetric domain in a complex vector space, also known as hermitian symmetric domain. It is of the form

$$D = G/K$$

where  $G$  is the group of holomorphic automorphism of  $D$ , which is a real (semisimple) Lie group and  $K$  is a maximal compact subgroup of  $G$ . E.g.

- $D(0, 1) \cong \mathfrak{H} = \{z \in \mathbb{C} \mid \text{Im}z > 0\} \cong \text{SL}_2(\mathbb{R})/\text{SO}_2$ ;
- $\mathfrak{H}_g = \{Z \in M_g(\mathbb{C}) \mid Z = Z^T, \text{Im}Z > 0\} \cong \text{Sp}_{2g}(\mathbb{R})/\text{U}_g$ .

There is a notion of arithmetic subgroup  $\Gamma \subset G$ . E.g.,  $\text{SL}_2(\mathbb{Z}) \subset \text{SL}_2(\mathbb{R})$ .

# Arithmetic manifolds

Let  $D$  be a bounded symmetric domain in a complex vector space, also known as hermitian symmetric domain. It is of the form

$$D = G/K$$

where  $G$  is the group of holomorphic automorphism of  $D$ , which is a real (semisimple) Lie group and  $K$  is a maximal compact subgroup of  $G$ . E.g.

- $D(0, 1) \cong \mathfrak{H} = \{z \in \mathbb{C} \mid \text{Im}z > 0\} \cong \text{SL}_2(\mathbb{R})/\text{SO}_2$ ;
- $\mathfrak{H}_g = \{Z \in M_g(\mathbb{C}) \mid Z = Z^T, \text{Im}Z > 0\} \cong \text{Sp}_{2g}(\mathbb{R})/\text{U}_g$ .

There is a notion of arithmetic subgroup  $\Gamma \subset G$ . E.g.,  $\text{SL}_2(\mathbb{Z}) \subset \text{SL}_2(\mathbb{R})$ .

## Theorem (Bailey-Borel)

*For an arithmetic subgroup  $\Gamma \subset G$ ,  $X = \Gamma \backslash D$  has a natural algebraic variety structure. Indeed,  $X$  admits a canonical compactification  $X^*$ , usually called the Bailey-Borel (or minimal) compactification of  $X$ , and  $X^*$  can be embedded into some projective space  $\mathbb{P}^n$ .*

# Examples

- If  $D = D(0, 1)$ , we recover many (but not all) hyperbolic algebraic curves.

# Examples

- If  $D = D(0, 1)$ , we recover many (but not all) hyperbolic algebraic curves.
- For appropriate  $\Gamma \subset \mathrm{Sp}_{2g}(\mathbb{R})$ , the space

$$A_g := \Gamma \backslash \mathfrak{H}_g$$

can be identified as the moduli space of principally polarized abelian variety (with sufficiently small level). This is an algebraic variety defined over  $\mathbb{Q}$ .

# Examples

- If  $D = D(0, 1)$ , we recover many (but not all) hyperbolic algebraic curves.
- For appropriate  $\Gamma \subset \mathrm{Sp}_{2g}(\mathbb{R})$ , the space

$$A_g := \Gamma \backslash \mathfrak{H}_g$$

can be identified as the moduli space of principally polarized abelian variety (with sufficiently small level). This is an algebraic variety defined over  $\mathbb{Q}$ .

This means that a map  $f : S \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow S$ .

# Examples

- If  $D = D(0, 1)$ , we recover many (but not all) hyperbolic algebraic curves.
- For appropriate  $\Gamma \subset \mathrm{Sp}_{2g}(\mathbb{R})$ , the space

$$A_g := \Gamma \backslash \mathfrak{H}_g$$

can be identified as the moduli space of principally polarized abelian variety (with sufficiently small level). This is an algebraic variety defined over  $\mathbb{Q}$ .

This means that a map  $f : S \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow S$ .

- For general  $D$  and  $\Gamma$  congruent subgroup of  $G$ , the space  $\Gamma \backslash D$  is a (connected component of a) Shimura variety.

# Hyperbolicity

It follows immediately from Liouville's theorem that  $\Gamma \backslash D$  is Brody hyperbolic.

# Hyperbolicity

It follows immediately from Liouville's theorem that  $\Gamma \backslash D$  is Brody hyperbolic. It is much more difficult to prove the analogous of great Picard theorem for  $X$ .

## Theorem (Borel)

*Every holomorphic map  $f : D(0, r)^\times \rightarrow X$  extends to a holomorphic map  $f : D(0, r) \rightarrow X^*$ .*



# Hyperbolicity

It follows immediately from Liouville's theorem that  $\Gamma \backslash D$  is Brody hyperbolic. It is much more difficult to prove the analogous of great Picard theorem for  $X$ .

## Theorem (Borel)

*Every holomorphic map  $f : D(0, r)^\times \rightarrow X$  extends to a holomorphic map  $f : D(0, r) \rightarrow X^*$ .*

## Corollary

*Every holomorphic map  $f : S \rightarrow X$  from an algebraic variety (over  $\mathbb{C}$ ) is algebraic.*

# Hyperbolicity

It follows immediately from Liouville's theorem that  $\Gamma \backslash D$  is Brody hyperbolic. It is much more difficult to prove the analogous of great Picard theorem for  $X$ .

## Theorem (Borel)

*Every holomorphic map  $f : D(0, r)^\times \rightarrow X$  extends to a holomorphic map  $f : D(0, r) \rightarrow X^*$ .*

## Corollary

*Every holomorphic map  $f : S \rightarrow X$  from an algebraic variety (over  $\mathbb{C}$ ) is algebraic.*

## Corollary

*The algebraic structure on  $X$  is unique.*

This is extremely important for the arithmetic theory of Shimura varieties.

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\};$

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

They satisfy triangle inequalities: for  $a, b \in \mathbb{Q}$ ,

$$|a + b|_v \leq |a|_v + |b|_v,$$

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

They satisfy triangle inequalities: for  $a, b \in \mathbb{Q}$ ,

$$|a + b|_v \leq |a|_v + |b|_v, \quad |a + b|_p \leq \max\{|a|_p, |b|_p\}.$$

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

They satisfy triangle inequalities: for  $a, b \in \mathbb{Q}$ ,

$$|a + b|_v \leq |a|_v + |b|_v, \quad |a + b|_p \leq \max\{|a|_p, |b|_p\}.$$

The completion of  $\mathbb{Q}$  with respect to  $|\cdot|_v$  is  $\mathbb{Q}_v$ , which is the set of Cauchy sequences  $(a_n)_n$  in  $\mathbb{Q}$  (w.r.t.  $|\cdot|_v$ ) modulo null sequence.

# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

They satisfy triangle inequalities: for  $a, b \in \mathbb{Q}$ ,

$$|a + b|_v \leq |a|_v + |b|_v, \quad |a + b|_p \leq \max\{|a|_p, |b|_p\}.$$

The completion of  $\mathbb{Q}$  with respect to  $|\cdot|_v$  is  $\mathbb{Q}_v$ , which is the set of Cauchy sequences  $(a_n)_n$  in  $\mathbb{Q}$  (w.r.t.  $|\cdot|_v$ ) modulo null sequence.

- $\mathbb{Q}_\infty = \mathbb{R}$ , and  $\overline{\mathbb{R}} = \mathbb{C}$  is complete, and  $\dim_{\mathbb{R}} \mathbb{C} = 2$ .



# A little bit about $p$ -adics

There are different absolute values  $|\cdot|_v$  of  $\mathbb{Q}$ :

- $|\cdot|_\infty$  = the Archimedean absolute value:  $|\frac{m}{n}|_\infty = \max\{\frac{m}{n}, -\frac{m}{n}\}$ ;
- $|\cdot|_p$  = the  $p$ -adic absolute value:  
if  $\frac{m}{n} = p^\nu \frac{m_1}{n_1}$  with  $(m_1 n_1, p) = 1$ , then  $|\frac{m}{n}|_p = \frac{1}{p^\nu}$ .

They satisfy triangle inequalities: for  $a, b \in \mathbb{Q}$ ,

$$|a + b|_v \leq |a|_v + |b|_v, \quad |a + b|_p \leq \max\{|a|_p, |b|_p\}.$$

The completion of  $\mathbb{Q}$  with respect to  $|\cdot|_v$  is  $\mathbb{Q}_v$ , which is the set of Cauchy sequences  $(a_n)_n$  in  $\mathbb{Q}$  (w.r.t.  $|\cdot|_v$ ) modulo null sequence.

- $\mathbb{Q}_\infty = \mathbb{R}$ , and  $\overline{\mathbb{R}} = \mathbb{C}$  is complete, and  $\dim_{\mathbb{R}} \mathbb{C} = 2$ .
- $\mathbb{Q}_p$ ,  $\dim_{\mathbb{Q}_p} \overline{\mathbb{Q}_p} = \infty$ . The completion  $\mathbb{C}_p$  of  $\overline{\mathbb{Q}_p}$  is algebraically closed.

In each case, the absolute value  $|\cdot|_v$  extends uniquely to these fields.

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object.

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

There are also non-archimedean analogue of open disc

$$D(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p < r\} = \cup_{r' < r} D^+(z_0, r')$$

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

There are also non-archimedean analogue of open disc

$$D(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p < r\} = \cup_{r' < r} D^+(z_0, r')$$

Some pathologies (the first two are due to strong triangle inequality):

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

There are also non-archimedean analogue of open disc

$$D(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p < r\} = \cup_{r' < r} D^+(z_0, r')$$

Some pathologies (the first two are due to strong triangle inequality):

- If  $|a|_p < r$ , then  $D^+(0, r) = D^+(a, r)$ ;

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

There are also non-archimedean analogue of open disc

$$D(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p < r\} = \cup_{r' < r} D^+(z_0, r')$$

Some pathologies (the first two are due to strong triangle inequality):

- If  $|a|_p < r$ , then  $D^+(0, r) = D^+(a, r)$ ;
- If  $|a|_p = r$ , then  $D(a, r) \subset D^+(0, r)$ ;

# A little bit $p$ -adic geometry

There are several ways to make sense of  $p$ -adic plane  $\mathbb{C}_p$  as a geometric object. (Slightly) imprecisely, we may think it as a topological space with a base of topology given by closed disc

$$D^+(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p \leq r\}.$$

There are also non-archimedean analogue of open disc

$$D(z_0, r) = \{z \in \mathbb{C}_p \mid |z - z_0|_p < r\} = \cup_{r' < r} D^+(z_0, r')$$

Some pathologies (the first two are due to strong triangle inequality):

- If  $|a|_p < r$ , then  $D^+(0, r) = D^+(a, r)$ ;
- If  $|a|_p = r$ , then  $D(a, r) \subset D^+(0, r)$ ;
- The (Berkovich space of)  $D(0, r)^\times$  is contractible.



# A little bit $p$ -adic analysis

“Analytic/holomorphic” functions  $\mathcal{O}(D^+(z_0, r))$  on the closed disc are

$$\mathbb{C}_p\langle z \rangle := \{f(z) = \sum a_i(z - z_0)^i \mid a_i \in \mathbb{C}_p, |a_i|_p r^i \rightarrow 0 \text{ as } i \rightarrow \infty\}.$$

# A little bit $p$ -adic analysis

“Analytic/holomorphic” functions  $\mathcal{O}(D^+(z_0, r))$  on the closed disc are

$$\mathbb{C}_p\langle z \rangle := \{f(z) = \sum a_i(z - z_0)^i \mid a_i \in \mathbb{C}_p, |a_i|_p r^i \rightarrow 0 \text{ as } i \rightarrow \infty\}.$$

In general, a function on  $U$  is analytic if its restriction to every closed disc inside it is analytic.

# A little bit $p$ -adic analysis

“Analytic/holomorphic” functions  $\mathcal{O}(D^+(z_0, r))$  on the closed disc are

$$\mathbb{C}_p\langle z \rangle := \{f(z) = \sum a_i(z - z_0)^i \mid a_i \in \mathbb{C}_p, |a_i|_p r^i \rightarrow 0 \text{ as } i \rightarrow \infty\}.$$

In general, a function on  $U$  is analytic if its restriction to every closed disc inside it is analytic.

Some pathologies:

- In general, the anti derivative of an analytic function on  $D^+(0, r)$  is only an analytic function on a smaller disc.

# A little bit $p$ -adic analysis

“Analytic/holomorphic” functions  $\mathcal{O}(D^+(z_0, r))$  on the closed disc are

$$\mathbb{C}_p\langle z \rangle := \{f(z) = \sum a_i(z - z_0)^i \mid a_i \in \mathbb{C}_p, |a_i|_p r^i \rightarrow 0 \text{ as } i \rightarrow \infty\}.$$

In general, a function on  $U$  is analytic if its restriction to every closed disc inside it is analytic.

Some pathologies:

- In general, the anti derivative of an analytic function on  $D^+(0, r)$  is only an analytic function on a smaller disc.
- The exponential function  $\exp(z) = \sum_{i \geq 0} z^i / i!$  only converges for  $|z|_p < \frac{1}{p^{1/(p-1)}}$ .

# Non-archimedean Picard's theorem

Theorem (non-archimedean little Picard theorem)

*Every non-constant analytic function  $f : \mathbb{C}_p \rightarrow \mathbb{C}_p$  is surjective.*

# Non-archimedean Picard's theorem

## Theorem (non-archimedean little Picard theorem)

*Every non-constant analytic function  $f : \mathbb{C}_p \rightarrow \mathbb{C}_p$  is surjective.*

## Theorem (non-archimedean great Picard theorem)

*Every analytic function  $f : D(0, r)^\times \rightarrow \mathbb{C}_p$  with essential singularity at 0 is surjective.*

# Non-archimedean Picard's theorem

## Theorem (non-archimedean little Picard theorem)

*Every non-constant analytic function  $f : \mathbb{C}_p \rightarrow \mathbb{C}_p$  is surjective.*

## Theorem (non-archimedean great Picard theorem)

*Every analytic function  $f : D(0, r)^\times \rightarrow \mathbb{C}_p$  with essential singularity at 0 is surjective.*

As before, we can reformulate the above results as

## Theorem

- ① *Every analytic function  $\mathbb{C}_p \rightarrow \mathbb{C}_p^\times$  is constant.*
- ② *Every analytic function  $f : D(0, r)^\times \rightarrow \mathbb{C}_p^\times$  extends to an analytic function  $\tilde{f} : D(0, r) \rightarrow \mathbb{P}^1$ .*
- ③ *Every analytic map  $S \rightarrow \mathbb{G}_m$  from an algebraic curve  $S$  is algebraic.*

# $p$ -adic Brody hyperbolicity

We do not want to claim  $\mathbb{G}_m$  to be hyperbolic. For this reason,

## Definition (Javanpeykar-Vezzani)

A  $p$ -adic variety  $X$  is called Brody hyperbolic if every analytic map  $f : G \rightarrow X$  is constant, where  $G$  is an algebraic group over  $\mathbb{C}_p$ .



# $p$ -adic Brody hyperbolicity

We do not want to claim  $\mathbb{G}_m$  to be hyperbolic. For this reason,

## Definition (Javanpeykar-Vezzani)

A  $p$ -adic variety  $X$  is called Brody hyperbolic if every analytic map  $f : G \rightarrow X$  is constant, where  $G$  is an algebraic group over  $\mathbb{C}_p$ .

There is the analogue of GGL conjecture. Instead giving the formulation, we mention some evidences (even for the original GGL conjecture).

# $p$ -adic Brody hyperbolicity

We do not want to claim  $\mathbb{G}_m$  to be hyperbolic. For this reason,

## Definition (Javanpeykar-Vezzani)

A  $p$ -adic variety  $X$  is called Brody hyperbolic if every analytic map  $f : G \rightarrow X$  is constant, where  $G$  is an algebraic group over  $\mathbb{C}_p$ .

There is the analogue of GGL conjecture. Instead giving the formulation, we mention some evidences (even for the original GGL conjecture).

## Theorem (Cherry, Kawamata, Ueno)

*Let  $K = \overline{K}$  with  $\text{char} K = 0$ . Let  $X$  be a closed subvariety of an abelian variety  $A$  over  $K$ . Then the following are equivalent.*

- ①  *$X$  does not contain the translate of a positive-dimensional abelian subvariety of  $A$ .*
- ② *Every closed integral subvariety of  $X$  is of general type.*
- ③ *If  $K = \mathbb{C}$  or  $\mathbb{C}_p$ , the projective variety is Brody hyperbolic.*

# Main theorem

## Theorem (Oswal-Shankar-Z.)

*Every analytic map  $f : D(0, r)^\times \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  can be extended to an analytic map  $\tilde{f} : D(0, r) \rightarrow A_g^*$ .*

# Main theorem

## Theorem (Oswal-Shankar-Z.)

*Every analytic map  $f : D(0, r)^\times \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  can be extended to an analytic map  $\tilde{f} : D(0, r) \rightarrow A_g^*$ .*

## Corollary

*Every analytic map  $f : S \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  with  $S$  an algebraic variety is algebraic.*

## Corollary

*Every analytic map  $f : G \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  with  $G$  an algebraic group is constant.*

# Main theorem

## Theorem (Oswal-Shankar-Z.)

*Every analytic map  $f : D(0, r)^\times \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  can be extended to an analytic map  $\tilde{f} : D(0, r) \rightarrow A_g^*$ .*

## Corollary

*Every analytic map  $f : S \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  with  $S$  an algebraic variety is algebraic.*

## Corollary

*Every analytic map  $f : G \rightarrow A_g$  defined over some finite extension  $K/\mathbb{Q}_p$  with  $G$  an algebraic group is constant.*

- Our proof requires  $K/\mathbb{Q}_p$  as above. Unfortunately,  $K = \mathbb{C}_p$  is currently not allowed.
- By some standard arguments, the results hold with  $A_g$  replaced by Shimura varieties of abelian type.

# Step I

The minimal compactification of  $A_g$  looks like

$$A_g^* = A_g \sqcup \partial A_g^* = \sqcup_{g' \leq g} A_{g'}.$$

Here  $A_{g'}$  could appear in several boundary components and are equipped with appropriate level structures.

# Step I

The minimal compactification of  $A_g$  looks like

$$A_g^* = A_g \sqcup \partial A_g^* = \sqcup_{g' \leq g} A_{g'}.$$

Here  $A_{g'}$  could appear in several boundary components and are equipped with appropriate level structures. Such decomposition extend over  $\mathbb{Z}_p$ .

Then for each  $A_{g', \mathbb{F}_p}$ , one can consider its tubular neighborhood in  $A_{g, \mathbb{Q}_p}^*$ . (These are  $p$ -adic analytic spaces, but not algebraic varieties.)

# Step I

The minimal compactification of  $A_g$  looks like

$$A_g^* = A_g \sqcup \partial A_g^* = \sqcup_{g' \leq g} A_{g'}.$$

Here  $A_{g'}$  could appear in several boundary components and are equipped with appropriate level structures. Such decomposition extend over  $\mathbb{Z}_p$ .

Then for each  $A_{g', \mathbb{F}_p}$ , one can consider its tubular neighborhood in  $A_{g, \mathbb{Q}_p}^*$ . (These are  $p$ -adic analytic spaces, but not algebraic varieties.)

E.g. the tubular neighborhood of  $A_{g, \mathbb{F}_p}$  is contained in  $A_{g, \mathbb{Q}_p}$ , parameterizing those abelian varieties with (potentially) good reduction. It is the rigid analytic variety  $\mathcal{A}_g^{\text{rig}}$  associated to the formal completion of  $A_g/\mathbb{Z}_p$  (along  $p = 0$ ).



# Step I

The minimal compactification of  $A_g$  looks like

$$A_g^* = A_g \sqcup \partial A_g^* = \sqcup_{g' \leq g} A_{g'}.$$

Here  $A_{g'}$  could appear in several boundary components and are equipped with appropriate level structures. Such decomposition extend over  $\mathbb{Z}_p$ .

Then for each  $A_{g', \mathbb{F}_p}$ , one can consider its tubular neighborhood in  $A_{g, \mathbb{Q}_p}^*$ . (These are  $p$ -adic analytic spaces, but not algebraic varieties.)

E.g. the tubular neighborhood of  $A_{g, \mathbb{F}_p}$  is contained in  $A_{g, \mathbb{Q}_p}$ , parameterizing those abelian varieties with (potentially) good reduction. It is the rigid analytic variety  $\mathcal{A}_g^{\text{rig}}$  associated to the formal completion of  $A_g/\mathbb{Z}_p$  (along  $p = 0$ ).

## Proposition

*Every analytic map  $D(0, r)^\times \rightarrow A_g^*$  is contained in one of the above tubular neighborhoods.*

# Step II

The variety  $A_{g, \mathbb{F}_p}$  parameterizes abelian varieties (with polarization) over  $\mathbb{F}_p$ . It admits a decomposition

$$A_{g, \mathbb{F}_p} = \bigsqcup_{\phi} S_{\phi},$$

where each  $S_{\phi}$  is locally closed and two points  $x, y \in A_g(\overline{\mathbb{F}}_p)$  belong to the same  $S_{\phi}$  if the corresponding abelian varieties (with polarization)  $A_x$  and  $A_y$  are (quasi-)isogenous.

# Step II

The variety  $A_{g, \mathbb{F}_p}$  parameterizes abelian varieties (with polarization) over  $\mathbb{F}_p$ . It admits a decomposition

$$A_{g, \mathbb{F}_p} = \bigsqcup_{\phi} S_{\phi},$$

where each  $S_{\phi}$  is locally closed and two points  $x, y \in A_g(\overline{\mathbb{F}}_p)$  belong to the same  $S_{\phi}$  if the corresponding abelian varieties (with polarization)  $A_x$  and  $A_y$  are (quasi-)isogenous.

As before, for each  $S_{\phi}$ , one can consider its tubular neighborhood  $]S_{\phi}[\mathbf{\text{ in }} \mathcal{A}_g^{\text{rig}}$ .

# Step II

The variety  $A_{g, \mathbb{F}_p}$  parameterizes abelian varieties (with polarization) over  $\mathbb{F}_p$ . It admits a decomposition

$$A_{g, \mathbb{F}_p} = \bigsqcup_{\phi} S_{\phi},$$

where each  $S_{\phi}$  is locally closed and two points  $x, y \in A_g(\overline{\mathbb{F}_p})$  belong to the same  $S_{\phi}$  if the corresponding abelian varieties (with polarization)  $A_x$  and  $A_y$  are (quasi-)isogenous.

As before, for each  $S_{\phi}$ , one can consider its tubular neighborhood  $]S_{\phi}[$  in  $\mathcal{A}_g^{\text{rig}}$ .

## Theorem

*Every analytic map  $D(0, r)^{\times} \rightarrow \mathcal{A}_g^{\text{rig}}$  is contained in one of  $]S_{\phi}[$  as above.*

# Step II

The variety  $A_{g, \mathbb{F}_p}$  parameterizes abelian varieties (with polarization) over  $\mathbb{F}_p$ . It admits a decomposition

$$A_{g, \mathbb{F}_p} = \bigsqcup_{\phi} S_{\phi},$$

where each  $S_{\phi}$  is locally closed and two points  $x, y \in A_g(\overline{\mathbb{F}_p})$  belong to the same  $S_{\phi}$  if the corresponding abelian varieties (with polarization)  $A_x$  and  $A_y$  are (quasi-)isogenous.

As before, for each  $S_{\phi}$ , one can consider its tubular neighborhood  $]S_{\phi}[$  in  $\mathcal{A}_g^{\text{rig}}$ .

## Theorem

*Every analytic map  $D(0, r)^{\times} \rightarrow \mathcal{A}_g^{\text{rig}}$  is contained in one of  $]S_{\phi}[$  as above.*

The proof uses the Tate conjecture for abelian varieties over global function fields.

# Step III

Fixing a point  $x \in ]S_\phi[$ , Rapoport-Zink constructed a uniformization map (i.e. an analytic map which is a topological covering)

$$RZ_x \rightarrow ]S_\phi[ ,$$

where  $RZ_x$  is certain rigid analytic space parameterizing abelian varieties with (quasi-)isogeny to the abelian variety  $A_x$  at  $x$ .

# Step III

Fixing a point  $x \in ]S_\phi[$ , Rapoport-Zink constructed a uniformization map (i.e. an analytic map which is a topological covering)

$$RZ_x \rightarrow ]S_\phi[ ,$$

where  $RZ_x$  is certain rigid analytic space parameterizing abelian varieties with (quasi-)isogeny to the abelian variety  $A_x$  at  $x$ . In addition, there exists an analytic map

$$\pi_{GM} : RZ_x \rightarrow \mathcal{F}\ell,$$

called the (Grothendieck-Messing) period map.

# Step III

Fixing a point  $x \in ]S_\phi[$ , Rapoport-Zink constructed a uniformization map (i.e. an analytic map which is a topological covering)

$$RZ_x \rightarrow ]S_\phi[ ,$$

where  $RZ_x$  is certain rigid analytic space parameterizing abelian varieties with (quasi-)isogeny to the abelian variety  $A_x$  at  $x$ . In addition, there exists an analytic map

$$\pi_{GM} : RZ_x \rightarrow \mathcal{F}\ell,$$

called the (Grothendieck-Messing) period map.

We may summarize the last (and the main) step of the proof as filling out dotted arrows in the following commutative

$$\begin{array}{ccccc}
 & D(0, r)^\times & \longrightarrow & D(0, r) & \\
 & \swarrow & & \searrow & \\
 & (1) \downarrow & & (3) \swarrow & \downarrow (2) \\
 \mathcal{A}_g^{\text{rig}} & \longleftarrow RZ_x & \xrightarrow{\pi_{GM}} & \mathcal{F}\ell. & 
 \end{array}$$



# Cohomology of (algebraic) variety

Let  $X$  be a smooth (projective) algebraic variety over a field  $k$  ( $\text{char} k = 0$ ). There are various cohomology theory attached to  $X$

- $\sigma : k \subset \mathbb{C}$ , the singular/Betti cohomology  $H_B^*(\sigma X, \mathbb{Z})$ ;
- The étale cohomology  $H_{\text{et}}^*(X_{\bar{k}}, \mathbb{Q}_p)$ ;
- The de Rham cohomology  $H_{\text{dR}}^*(X/k)$ .

# Cohomology of (algebraic) variety

Let  $X$  be a smooth (projective) algebraic variety over a field  $k$  ( $\text{char } k = 0$ ). There are various cohomology theory attached to  $X$

- $\sigma : k \subset \mathbb{C}$ , the singular/Betti cohomology  $H_B^*(\sigma X, \mathbb{Z})$ ;
- The étale cohomology  $H_{\text{et}}^*(X_{\bar{k}}, \mathbb{Q}_p)$ ;
- The de Rham cohomology  $H_{\text{dR}}^*(X/k)$ .

Now let  $f : X \rightarrow S$  be a family of smooth projective varieties. Then various cohomology theories of  $\{X_s\}_s$  also vary in family, giving:

- a  $\mathbb{Z}$ -local system on  $\sigma S$ ;
- a étale local system on  $S_{\text{et}}$ ;
- a vector bundle  $\mathcal{E}$  on  $S$  with a flat connection (the Gauss-Manin connection), and a decreasing filtration of  $\mathcal{E}$  by subbundles  $\text{Fil}^\bullet$ .

# Cohomology of (algebraic) variety

Let  $X$  be a smooth (projective) algebraic variety over a field  $k$  ( $\text{char } k = 0$ ). There are various cohomology theory attached to  $X$

- $\sigma : k \subset \mathbb{C}$ , the singular/Betti cohomology  $H_B^*(\sigma X, \mathbb{Z})$ ;
- The étale cohomology  $H_{\text{et}}^*(X_{\bar{k}}, \mathbb{Q}_p)$ ;
- The de Rham cohomology  $H_{\text{dR}}^*(X/k)$ .

Now let  $f : X \rightarrow S$  be a family of smooth projective varieties. Then various cohomology theories of  $\{X_s\}_s$  also vary in family, giving:

- a  $\mathbb{Z}$ -local system on  $\sigma S$ ;
- a étale local system on  $S_{\text{et}}$ ;
- a vector bundle  $\mathcal{E}$  on  $S$  with a flat connection (the Gauss-Manin connection), and a decreasing filtration of  $\mathcal{E}$  by subbundles  $\text{Fil}^\bullet$ .

There are various comparison isomorphisms between different cohomology theories.

# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathbb{R}Z_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathbb{R}Z_x$ ,

# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathrm{RZ}_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathrm{RZ}_x$ , whose GM connection is canonically trivial by the theory of crystalline cohomology.

# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathrm{RZ}_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathrm{RZ}_x$ , whose GM connection is canonically trivial by the theory of crystalline cohomology. The variation of the filtration  $\mathrm{Fil}^\bullet$  (on  $\mathrm{RZ}_x$ ) defines the period map  $\mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ .

# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathrm{RZ}_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathrm{RZ}_x$ , whose GM connection is canonically trivial by the theory of crystalline cohomology. The variation of the filtration  $\mathrm{Fil}^\bullet$  (on  $\mathrm{RZ}_x$ ) defines the period map  $\mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ .

Now we have  $D(0, r)^\times \rightarrow \mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ . The following theorem, which can be regarded as the  $p$ -adic analogue of Schmid's theorem on limit Hodge structure, implies that this map is meromorphic and therefore extends to an analytic map  $D(0, r) \rightarrow \mathcal{F}\ell$ .

# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathrm{RZ}_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathrm{RZ}_x$ , whose GM connection is canonically trivial by the theory of crystalline cohomology. The variation of the filtration  $\mathrm{Fil}^\bullet$  (on  $\mathrm{RZ}_x$ ) defines the period map  $\mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ .

Now we have  $D(0, r)^\times \rightarrow \mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ . The following theorem, which can be regarded as the  $p$ -adic analogue of Schmid's theorem on limit Hodge structure, implies that this map is meromorphic and therefore extends to an analytic map  $D(0, r) \rightarrow \mathcal{F}\ell$ .

## Theorem (Diao-Lan-Liu-Zhu)

*Let  $\mathbb{L}$  be a de Rham  $p$ -adic local system on  $D(0, r)^\times$ , with the associated filtered connection  $(\mathcal{E}, \nabla, \mathrm{Fil}^\bullet)$ . Then  $(\mathcal{E}, \nabla)$  admits a canonical extension (in the sense of Deligne) to a vector bundle  $\bar{\mathcal{E}}$  on  $D(0, r)$  with logarithmic pole. In addition, the filtration  $\mathrm{Fil}^\bullet$  extends to a filtration of  $\bar{\mathcal{E}}$  by vector bundles.*



# Limit of $p$ -adic Hodge structure

Now the uniformization map  $\mathrm{RZ}_x \rightarrow A_g$  corresponds to a family of abelian varieties  $A \rightarrow \mathrm{RZ}_x$ , whose GM connection is canonically trivial by the theory of crystalline cohomology. The variation of the filtration  $\mathrm{Fil}^\bullet$  (on  $\mathrm{RZ}_x$ ) defines the period map  $\mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ .

Now we have  $D(0, r)^\times \rightarrow \mathrm{RZ}_x \rightarrow \mathcal{F}\ell$ . The following theorem, which can be regarded as the  $p$ -adic analogue of Schmid's theorem on limit Hodge structure, implies that this map is meromorphic and therefore extends to an analytic map  $D(0, r) \rightarrow \mathcal{F}\ell$ .

## Theorem (Diao-Lan-Liu-Zhu)

*Let  $\mathbb{L}$  be a de Rham  $p$ -adic local system on  $D(0, r)^\times$ , with the associated filtered connection  $(\mathcal{E}, \nabla, \mathrm{Fil}^\bullet)$ . Then  $(\mathcal{E}, \nabla)$  admits a canonical extension (in the sense of Deligne) to a vector bundle  $\bar{\mathcal{E}}$  on  $D(0, r)$  with logarithmic pole. In addition, the filtration  $\mathrm{Fil}^\bullet$  extends to a filtration of  $\bar{\mathcal{E}}$  by vector bundles.*

That  $D(0, r) \rightarrow \mathcal{F}\ell$  lifts to  $D(0, r) \rightarrow \mathrm{RZ}_x$  uses some theory of crystalline representations.

# Thank You!