
Study on Lamprey: Sex Ratio and Its Impact

Summary

Resource availability was found to influence whether lampreys end up as males or females, which makes the sex ratio of lampreys change in response to the external environment. This adaptive sex ratio has significant effects on ecosystems. We simulated the evolution of a community containing lamprey and analyzed the impact of adaptive sex ratios at multiple levels.

First, we introduced **the male advantageous coefficient** and developed the individual development model to analyze the sex determination process of lamprey. Based on Logistic equation, we developed the intraspecific competition model. Results of those models indicate basic regulation of the sex ratio. We also used the lotka-voterra model to preliminarily analyze lamprey interactions with other species. To interpret the advantages and disadvantages of adaptive sex ratio, we defined the fitness function. After conducting simulations under different conditions, we found that adaptive sex ratio improves population fitness. Furthermore, we calculated the optimal sex ratio for different resource availability. We found that lamprey's sex ratio is very close to the optimal ratio.

What's more, **the complex community dynamical model** was developed by the introduction of **the interspecific interaction matrix**. On the basis of a established food network, we conducted **controlled community simulation** to further study the evolution of communities. In doing so, to study the stability of ecosystems, we defined **the stability function** based on the number of species, the average interaction strength, and the connectivity of the food network. Through controlled community simulations we found that adaptive sex ratio not only increase the competitive advantage of lamprey, but also promote the survival of other species and improve community stability.

Besides, we developed the dynamical model of parasites population and coupled it with complex community dynamical model. Similarly, the results of controlled community simulation suggest that lamprey's adaptive sex ratio can offer advantages for parasites.

Additionally, we conducted sensitivity analysis to unpredictable factors and lethality due to infection. The Gaussian noise was applied to simulate the natural fluctuation of resources availability. Although the communities no longer have a steady state, random noise has little impact on the stability of community, and our conclusions still holds. Studying on lethality due to infection, we found that excessive lethality would disrupt lamprey's sex ratio regulation mechanisms, which would weaken its competitive advantage, with consequences for all ecosystem members.

Finally, based on our study, we recommend that more measurements and studies on the lamprey should be conducted to better protect and recognize this amazing species.

Keywords:Lamprey, Sex ratio, Resource availability, Dynamical model, Ecosystem stability, Optimal strategy, Lotka-Voterra model, population interaction, Logistic equation

Contents

1	Introduction	1
1.1	Background	1
1.2	Restatement of the Problem	1
1.3	Our Work	1
2	Notations and Assumptions	3
2.1	Notations	3
2.2	Assumptions	4
3	Impact on the Large Ecosystem	4
3.1	Model Overview	4
3.2	Sex Determination Mechanism	4
3.3	the Dynamics Model of Lamprey Population	5
3.4	Lampreys and Prey	6
3.5	Result Analysis	7
4	Advantages and Disadvantages	7
4.1	Fitness Function	7
4.2	Different Supply Conditions	8
4.3	Fitness-based Optimization Strategies	9
5	Impact on the Stability of ecosystem	9
5.1	the Dynamics Model of Complex ecosystem	9
5.2	the Stability Function of Community	10
5.3	Community Simulation	10
6	Interaction With Parasites	12
6.1	the Dynamics model of Parasites	12
6.2	Community Simulation	13
7	Sensitivity Analysis	13
7.1	Models Review	13
7.2	Sensitivity to Unpredictable Factor	14
7.3	Sensitivity to Lethality due to infection	15
8	Strengths and Weaknesses	15
8.1	Strengths	15
8.2	Weaknesses	16
9	Conclusion	16

1 Introduction

1.1 Background

While some animal species exist outside of the usual male or female sexes, most species are substantially either male or female. Although many species exhibit a 1:1 sex ratio at birth, other species deviate from an even sex ratio. This is called adaptive sex ratio variation. For example, the temperature of the nest incubating eggs of the American alligator influences the sex ratios at birth.

The role of lampreys is complex. In some lake habitats, they are seen as parasites with a significant impact on the ecosystem, whereas lampreys are also a food source in some regions of the world.

The sex ratio of sea lampreys can vary based on external circumstances. Sea lampreys become male or female depending on how quickly they grow during the larval stage. These larval growth rates are influenced by the availability of food. The sex ratio in sea lampreys can influence other species and the whole ecosystem.



Figure 1: The Lampreys

1.2 Restatement of the Problem

Focus on the question of sex ratios and their dependence on local conditions, specifically for sea lampreys. Examine the advantages and disadvantages of the ability for a species to alter its sex ratio depending on resource availability. Develop and examine a model to provide insights into the resulting interactions in an ecosystem.

- Study the impact on the larger ecological system when the population of lampreys can alter its sex ratio.
- Study the advantages and disadvantages to the population of lampreys
- Study the impact on the stability of the ecosystem given the changes in the sex ratios of lampreys
- Study advantages offered by the variable sex ratio in the lamprey population to other members in the ecosystem.

1.3 Our Work

- We developed a sex determination function based on larval growth rate. Applying the normal distribution function, we introduced male advantageous coefficient to simulate the changes of sex ratio with respect to the resources availability.

- On the basis of the Logistic equation, we developed the intraspecific competition model to illustrate the implication of adaptive sex ratio. The model contains the resource constraints and competitive pressures between the sexes.
- We defined the fitness function to evaluate the advantages of a swarm. This function contains the reproduction success and advantages in predation. Male advantageous coefficient was introduced to indicate the resource availability. It accounts for lampreys' behavior that changing the sex ratio. Two simulation with different conditions further studied changes in fitness.
- Through developing the differential equation, we found the optimal sex ratio with respect to male advantageous coefficient and comparing it with actual sex ratio.
- After establishing a food network containing lampreys, we developed a group of differential equations based on the Lotka-Volterra model. By introducing the interspecific interaction matrix, we successfully coupled them together to form the complex community dynamical model. Then we conducted community simulations to study the interspecific interaction.
- We developed the stability function to evaluate the stability of community. It incorporates the number of species, average interaction strength, and food network connectivity. Then we studied the impacts of adaptive sex ratio on the stability of the food network.
- We developed the dynamical model of parasites and coupled it with complex community dynamical model. Then we used community simulation to explain the interaction between parasites and lampreys, and the whole community.
- Gaussian noise was applied to simulate the unpredictable fluctuation of resources. We tested the robustness of our model under random noise.
- We moderated the lethality due to infection of lamprey to test the sensibility of our models.




Figure 2: The flow chart of our work

2 Notations and Assumptions

2.1 Notations

Symbols	Description
N	Total population of the sea lampreys
t	Time
F	Population of female lampreys
M	Population of male lampreys
r	Birth rate of lampreys
rs	Sex Ratio in lampreys
s	Growth rate of lampreys
S	Average growth rate of lampreys
R	Natural increase rate of prey
C	Total popular of prey
W	Interspecific interaction matrix
K	Resources availability
H	Population of parasites
r_A	Natural increasing rate of species A
r_B	Natural increasing rate of species B
r_C	Natural increasing rate of species C
r_H	Natural increasing rate of parasites
$g(S)$	Sex determination function
P_f	Percentage of female lampreys
R_0	Basic reproduction rate of lamprey
R_H	Basic reproduction rate of parasites
γ	Basic reproduction rate scale coefficient
μ	Actual reproduction rate of parasites
δ	Infection rate scale factor
ρ	Actual infection rate
ρ_0	Basic infection rate
λ	Lethality due to infection
Ω	Stability of a community
ϵ	Degree of interspecific connectivity
ξ	Average strength of action
σ	Sex determination threshold
η	Male advantageous coefficient
α	Female mortality rate
β	Competition coefficient of female with respect to male
θ	Male mortality rate
ω	Competition coefficient of male with respect to female

Table 1: Symbols and description

2.2 Assumptions

In order to simplify the problem, we have made some assumptions and provided reasonable justifications for them.

- **Assumption1:**Lampreys are affected by predators that feed on them in certain proportions.
Justification:Lamprey's predators include many kinds of birds and fish. In a stable ecosystem, mortality ratio tends to be constant.
- **Assumption2:**Males are dominant in intraspecific competition, and this dominance is evident in times of food scarcity
Justification:The sex ratio is higher in times of food scarcity, which means male individuals have advantages over females in surviving hard times[1].
- **Assumption3:**Populations of lampreys with higher sex ratios have higher reproductive success, interspecific competitiveness, and predation, while mating frequency decreases.
Justification:For the vast majority of animals, including lamprey, males catch more food. A surplus of males allows only the strongest individuals to get a chance to mate, thus increasing reproductive success. However, unlike other animals, the lamprey's parent individuals will die after mating.This makesThis allows enoughThis allows enough females toThis leaves enough females forThis makes it important to have enough females to ensure that the mating frequencyThis makes a sufficient number of females essential to ensure mating frequency.This makes a sufficient number of females essential to ensure mating frequency.
- **Assumption4:**The growth rate of lampreys follow the Gaussian distribution.
Justification:The Gaussian distribution, also known as the normal distribution, is widely used to describe the distribution of biological properties. Applying the Gaussian distribution has consistently proven to be a reliable approach

3 Impact on the Large Ecosystem

3.1 Model Overview

To examine the interaction between lampreys and the ecosystem, we firstly developed the dynamical model of lamprey population based on the modified Logistic equations. Then the Lotka-Voterra model was used to simulate the interspecific competition.

3.2 Sex Determination Mechanism

Lampreys gonads are not yet developed at an early age. After juvenile age, they undergo metamorphosis to become adults[2]. The growth rates of lampreys during the juvenile period determines the sex of the adult lampreys. Therefore, the sex determination function can be defined as a step function.

$$g(s) = \begin{cases} 1 & s < \sigma \\ 0 & s \geq \sigma \end{cases} \quad (1)$$

If $g(S)$ of a lamprey larvae equals to 1, it will become a female adult after metamorphosis, otherwise a male adult will be generated. To determine the degree of food shortage, the male advantageous coefficient is defined.

$$\eta = \frac{N}{C} \quad (2)$$

When η tends to 0, the food supply is so enough that male have no advantages over female. Bigger the η , the scarcer the food. Therefore, η directly influences the average growth rate. We assuming that the growth rate of lampreys follows a Gaussian distribution.

$$\frac{dr_s}{d\eta} = \frac{\int_0^{\frac{3\sigma}{\sqrt{2}S}} \exp(-t^2) dt}{\sqrt{\pi} - \int_0^{\frac{3\sigma}{\sqrt{2}S}} \exp(-t^2) dt} \quad (3)$$

It's hard to find analytic solution. Using finite differential equations, we can find the numerical solution.




Figure 3: Adaptive Sex Ratio of lampreys

3.3 the Dynamical Model of Lamprey Population

To illustrate the implications of the mechanism of change in the sex ratio of lampreys, we envision a group of lampreys in which female individuals compete with male individuals. In such a group, the sex ratio will changes over time. To simulate the changes, we use the modified logistic equation to represent the population of both sexes.

$$\frac{dF}{dt} = rF(1 - \alpha F - \beta M) \quad (4)$$

$$\frac{dM}{dt} = rM(1 - \theta M - \omega F) \quad (5)$$

$$\frac{dN}{dt} = \frac{dM}{dt} + \frac{dF}{dt} \quad (6)$$

Besides, the correction terms are added to represent the intraspecific competition between two sexes. It is important to note that due to their unique reproduction, lampreys have no real birth rate. Here, birth

rate is defined as the ratio of the difference between the number of successfully hatched lampreys in the offspring and the population of the previous generation to the population of the previous generation. The birth ratio can be represented by the Under certain parameter settings, the model predicts changes in population size and sex ratio.




Figure 4: Population Dynamics of Lampreys




Figure 5: Sex Ratio of Lampreys

3.4 Lampreys and Prey

Lampreys burrows into the sand or gravel during the day and comes out at night to feed. It adsorbs on other fish with its sucker-like mouth and files through the fish with the horny teeth on both sides of the mouth and on the tongue, sucking their blood and flesh, and sometimes only a skeleton remains of the sucked fish in the end.

The population of prey is represented by R , and the possibility of death caused by lampreys attack is ζ . The Lotka-Volterra equation is a classical differential equation model that gives an indication of how changes in the population sizes of predators and prey depend on the numbers and interactions between them. Using Lotka-Volterra model, we can calculate population of the prey.

$$\frac{dC}{dt} = RC - \zeta CN \quad (7)$$

Solving this equation with the dynamical model of lampreys, the result is shown in Figure.5.

We can see it clearly that sharp decline of the population of prey was caused by lampreys, and after a short-lived rise, the population of lampreys finally comes to zero.




Figure 6: Population Dynamics of Lampreys and Prey

3.5 Result Analysis

Through our model, the population dynamics of lampreys and its interaction with community are described. Alerting parametries in our moedl, we can an further improve the accuracy of the model and provide in-depth observation and analysis of the dynamics of interactions between groups of organisms. Through our analysis, we found that sex ratio plays a important role in the survival of lampreys. The lampreys gouns with more male individuals might predate more prey and behave higher success of reproduction. However, the female individuals are essential to the reproductive frequency. So the sex ratio will converge to a stable value depending on food availability. In our simulation, the highest value of sex ratio about 2.3 was observed. When food supply is sufficient, the value may approximately approach 1.

4 Advantages and Disadvantages

Analyzing the interaction between sex ratio and environment, it's obvious that for a given environment, there exists a sex ratio that is best adapted to it. To find the best value, we defined the objective function $f(rs)$ to describe the fitness of the sex ratio.

4.1 Fitness Function

We can use reproductive success to estimate the fitness of a population. Considering the unique breeding method, the parental individuals will all die after reproduction.

$$f(x) = \frac{R_0}{1 + rs} \frac{rs - \eta}{1 + rs} \quad (8)$$

The function is composed by two parts. First part is the reproductive term, representing the number of individuals reproduced. Second part represents the advantage of predation brought by surplus male individuals.

4.2 Different Supply Conditions

The ability of lampreys to alert the sex ratio plays an important role in fitting the environment. We studied two different supply conditions. When food supply is sufficient, C tends to infinity, the male advantageous coefficient tends to 0. In that case, the best sex ratio is 1.




Figure 7: Population Dynamics of Lampreys

When food supply is scarce, that is, population of prey is 5 times bigger than the population of lampreys. In that case, we can find that the sex ratio tends to 1.8 approximately.




Figure 8: Population Dynamics of Lampreys

In both conditions, the fitness of group increase rapidly and approach the theoretical maximum.




Figure 9: Fitness changes over time

4.3 Fitness-based Optimization Strategies

Combining the dynamical model of lampreys, we can express population of lampreys with respect to the food supply. Solving the equations, we have the differential equation of the best sex ratio with respect to η .

$$\frac{dr_s}{d\eta} = \frac{2(rs - \eta)}{\eta + 1} + 1 \quad (9)$$

We can see that the best sex ratio increase with food supply becoming scarce.




Figure 10: The Best Sex Ratio and Actual Sex Ratio

5 Impact on the Stability of ecosystem

5.1 the Dynamical Model of Complex ecosystem

To examine the impact of lampreys on the stability of ecosystem, we established a community composed by four species including lampreys. Its food network is shown in Fig.9.




Figure 11: The Food Network of Community

We use the Lotka-Voterra model to describe the interspecific interaction. The dynamical model of complex community is shown below.

$$\frac{dA}{dt} = r_A A + W_{AN}N + W_{AB}B \quad (10)$$

$$\frac{dN}{dt} = rN + W_{NC}C + W_{NB}B + W_{NA}A \quad (11)$$

$$\frac{dB}{dt} = r_B B + W_{BC}C + W_{BN}N + W_{BA}A \quad (12)$$

$$\frac{dC}{dt} = r_C C(1 - KC) + W_{CN}N + W_{CB}B \quad (13)$$

in which W is the interspecific interaction matrix.

$$W = \begin{bmatrix} r_A & W_{AN} & W_{AB} & 0 \\ W_{NA} & r & W_{NB} & W_{NC} \\ W_{BA} & W_{BN} & r_B & W_{BC} \\ 0 & W_{CN} & W_{CB} & r_C \end{bmatrix} \quad (14)$$

For elements in the matrix, r_A , r_B , and r_C represent environment capacity of each species respectively. W_{AN} , W_{AB} , W_{NC} , and W_{BC} indicate the suppression of population growth by environmental capacity. W_{NB} and W_{BN} express competition between lampreys and species B. W_{CN} and W_{CB} denote the suppression of population growth by predation.

5.2 the Stability Function of Community

Based on previous research, we developed the stability function of community. It is related to the number of species in the community, the degree of interspecific connectivity, the average strength of action.

$$\xi = \frac{1}{2} I^T [I(W^T + W)] \quad (15)$$

$$\Omega = 2\xi\sqrt{\epsilon} \quad (16)$$

$$\Omega < 1 \quad (17)$$

Equation (16) is the equilibrium condition for community. Communities that meet the equilibrium condition are stable, which means such communities can spontaneously return to their initial state after receiving a disturbance[3].

5.3 Community Simulation

We carried out two simulations with the same initial parameters, the difference being that one of them, lamprey, had the ability to change the sex ratio, while the other did not. lamprey's strategy for changing the sex ratio is to choose the optimal value based on the male advantageous coefficient.

The results are shown in Fig.13 and Fig.14.

Comparing the results we find that lamprey's ability to alter sex ratios not only increases its environmental capacity, but also increases its competitive advantage over species B. At the same time, the populations of all three species rose to some extent.




Figure 12: Linear Fitting for the Best Sex Ratio




Figure 13: Results of Community Simulation




Figure 14: Sex Ratio of lampreys




Figure 15: Stability of Community

Lamprey's sex ratio has stabilized over time. Changes in sex ratios significantly reduced the community stability function and increased the stability of the community. In summary, lamprey's ability to vary sex ratios not only facilitates its own survival, but also promotes the survival of other species, and is a way for the community to become more stable.

6 Interaction With Parasites

Parasites in lamprey mainly include *Diplostomum* sp., *Triaenophorus crassus*, *Echinorhynchus salmoneis*, and *Ergasilus* spp. It was found that males and females do not differ significantly in their resistance to parasites[4]. However, males are heavier and merely more numerous and are more likely to be infected with parasites[5]. There is a lack of research on the dangers of parasitic infections, and it has been found that the diseases that cause lamprey deaths are primarily fungal infections rather than parasites.

6.1 the Dynamical model of Parasites

Similarly, we use modified logistic equation to develop the dynamical model of parasites.

$$\frac{dH}{dt} = r_H H \left(1 - \frac{H}{\mu \rho N}\right) \quad (18)$$

$$\mu = R_H + \gamma r s \quad (19)$$

$$\rho = \rho_0 + \delta r s \quad (20)$$

This model describes community-parasite interactions and considers the effect of sex ratio on parasite populations.

The dynamical model of lampreys can be rewritten as:

$$\frac{dN}{dt} = (r - \lambda \rho) N + W_{NC} C + W_{NB} B + W_{NA} A \quad (21)$$

6.2 Community Simulation

After the introduction of parasites, we conducted two simulations. As in the previous study of community stability, the sex ratio of lamprey was fixed in one of the two simulations and the optimal value based on the MALE advantageous coefficient was taken in the other. Trough the simulations, we




Figure 16: Population Dynamics of parasites

found that adaptive sex ratio can increase the environment capacity of parasites, while have little impact on the whole community. That accounts for the fact that lampreys always carry tons of parasites.

7 Sensitivity Analysis

7.1 Models Review

First, here are our final mathematical models. The sex-determination model:

$$\frac{drs}{d\eta} = \frac{\int_0^{\frac{3\sigma}{\sqrt{2}S}} \exp(-t^2) dt}{\sqrt{\pi} - \int_0^{\frac{3\sigma}{\sqrt{2}S}} \exp(-t^2) dt} \quad (22)$$

Intraspecific competition model:

$$\frac{dF}{dt} = rF(1 - \alpha F - \beta M) \quad (23)$$

$$\frac{dM}{dt} = rM(1 - \theta M - \omega F) \quad (24)$$

$$\frac{dN}{dt} = \frac{dM}{dt} + \frac{dF}{dt} \quad (25)$$

The fitness optimization model:

$$\begin{aligned} \max \quad & f(x) = \frac{R_0}{1 + rs} \frac{rs - \eta}{1 + rs} \\ \text{s.t.} \quad & rs = N/F \\ & \frac{dF}{dt} = rF(1 - \alpha F - \beta M) \\ & \frac{dM}{dt} = rM(1 - \theta M - \omega F) \\ & \beta = \omega + \eta \end{aligned}$$

The complex community dynamical model:

$$\frac{dA}{dt} = r_A A + W_{AN}N + W_{AB}B \quad (26)$$

$$\frac{dN}{dt} = (r - \lambda\rho)N + W_{NC}C + W_{NB}B + W_{NA}A \quad (27)$$

$$\frac{dB}{dt} = r_B B + W_{BC}C + W_{BN}N + W_{BA}A \quad (28)$$

$$\frac{dC}{dt} = r_C C(1 - KC) + W_{CN}N + W_{CB}B \quad (29)$$

$$\frac{dH}{dt} = r_H H(1 - \frac{H}{\mu\rho N}) \quad (30)$$

Evaluation of community stability:

$$\Omega = 2\xi\sqrt{\epsilon} \quad (31)$$

7.2 Sensitivity to Unpredictable Factor

In order to simulate fluctuations in total environmental resources, we impose Gaussian noise on the natural growth rate of a first-tier consumer (species C). The interval of Gaussian noise needs to be kept within a reasonable range. Too small an interval will not simulate the effect of the unpredictable factor on the community, while too large an interval will affect the stability of the community or even cause species extinction.




Figure 17: Results of Community Simulation Under Gaussian Noise

We find that the introduction of noise makes the model unstable and no longer converges to a steady state. However, the community metrics still fluctuate around the stabilization point, and our previous conclusions still hold. This suggests that our model is correct and robust to some extent. It is also important to note that the Gaussian noise was limited to be very small in this simulation, one to two orders of magnitude smaller than r_C . Under greater noise, the community will move slowly toward extinction. In our simulation, the Gaussian noise acts only on r_C , which is not perfectly consistent with the real situation. Therefore, more complex models are needed to simulate the community under noise.

7.3 Sensitivity to Lethality due to infection

There are many different types of parasites that parasitize lamprey, and some of them can cause loss of vitality, mechanical damage, blindness, and even death. We adjusted the infection lethality of lamprey, performed separate colony simulations and measured various metrics of the community when it reached a steady state. The results are shown in Fig.18.




Figure 18: Results of Community Simulation with different lethality

We found that lethality increased when lamprey's environmental capacity and sex ratio declined, along with a loss of competitive advantage over species B. Parasite abundance also declined.

8 Strengths and Weaknesses

8.1 Strengths

- The sex-determination model reflects the real mechanism of lampreys' sex determination. Combining it with intraspecific competition model and fitness optimization model can provides comprehensive insights into the lampreys' behavior of changing sex ratio.
- With interspecific interaction matrix, the complex community dynamical model can describe the evolution of a community excellently. It's also flexible and can be applied into different conditions conveniently. With the help of stability functions, it is able to quantify the stability of communities and model their evolution.
- The introduction of Gaussian noise not only proves the robustness of model, but also simulate the natural changes in resources availability, which is more realistic.
- Our entire model comprehensively characterizes the effects of adaptive sex ratios at the individual, population, community, and ecosystem levels. The role of other parts of the ecosystem on lampreys is also considered.

8.2 Weaknesses

- Due to the lack of accurate experimental data on lamprey in terms of feeding and hunting, health status, and sex differentiation, our simulation parameters may differ from the real world in their settings.
- Parasitism is relatively independent in our model, and the interspecific action matrix only describes interspecific relationships within the food network; parasitism is excluded.

9 Conclusion

- The sex of the lamprey is determined by the rate of development of the hatchlings, which results in a sex ratio that is close to the optimal sex ratio given the availability of resources at the time.
- lamprey's adaptive sex ratio allows it to achieve a dominant position in the ecosystem. Since most farmed fish belong to their prey or hosts, in some cases the invasive lamprey has the potential to deal a severe blow to local fish farming.
- Adaptive aliasing in lamprey not only improves its own competitive advantage, but also promotes the survival of other species in the community. And it substantially improves ecosystem stability.
- Parasites can benefit from lamprey's adaptive sex ratio, however excessive infection lethality can disrupt the sex ratio regulation mechanism, ultimately causing damage to all members of the ecosystem.

References

- [1] Docker, Margaret F. et al. "Age, growth, and sex ratio among populations of least brook lamprey, *Lampetra aepyptera*, larvae: an argument for environmental sex determination", *Environmental Biology of Fishes* 41.1 (1994): 191-205.
- [2] Jun, LI et al. "THE EARLY DEVELOPMENT OF *LAMPETRA MORII*", *Acta Hydrobiologica Sinica* 43.4 (2019): 814-824.
- [3] May, Robert M.. "Relationships Among Various Types Of Population Models", *American Naturalist* 107.953 (1973): 46-57.
- [4] Shavalier, Megan A. et al. "Parasites and microbial infections of lamprey (order Petromyzontiformes Berg 1940): A review of existing knowledge and recent studies", *Journal of Great Lakes Research* 47 (2021): S90-S111.
- [5] Johnson, Nicholas S. et al. "Reproductive Ecology Of Lampreys", *LAMPREYS: BIOLOGY, CONSERVATION AND CONTROL*, VOL 1 37. (2015): 265-303.

Report on Use of AI:

ZHIPUAI ChatGLM (v 1.0)

Query 1: Please help me interpret this paper: Parasites and microbial infections of lamprey (order Petromyzontiformes Berg 1940): A review of existing knowledge and recent studies

Output: Regarding this paper "Parasitic and microbial infections in blind eels (Petromyzontiformes Berg 1940): a review of existing knowledge and recent studies", from the title we can understand that this is a literature review on parasitic and microbial infections in blind eels. The authors aim to summarize existing research findings and present recent research advances.

Since I do not have access to the latest database of academic papers, the following interpretation will be based on general knowledge and understanding of the literature review:

Background of the study: Blind eels are an ancient and primitive group of fishes belonging to the jawless class, which is an important branch in the evolutionary history of fishes. Due to its specialized habits and physiology, the study of parasitic and microbial infections in blind eels is important for understanding microbe-host interactions in ecosystems.

Summary of existing knowledge: The paper may start with a review of the basics of parasitic and microbial infections in blind eels, such as the types of infections (e.g., parasites such as protozoa, hem-morhynchids, nematodes, and microorganisms such as bacteria, fungi, etc.), the modes of infections, and the immune responses of the hosts, etc. The authors may then summarize the current knowledge in the paper.

Recent Research Advances: Next, the author may present recent research findings. This may include new discoveries such as new understanding of the life cycle of a particular parasite or microorganism, infection strategies, or revelations about the molecular mechanisms of host-parasite interactions.

Research Implications: The paper may discuss the implications of these studies for basic ecology, pathogenesis, and biodiversity. In particular, studies to understand the unique physiological and ecological characteristics of the blind eel may shed light on the development of new drugs or treatments.

Directions for future research: Finally, based on the existing knowledge and research progress, the authors may suggest directions for future research. For example, in-depth studies on the parasite adaptation mechanisms in the special host environment of blind eels, or the development of novel microbial drugs.

In summary, this paper provides researchers and interested readers with a comprehensive overview of the field of parasites and microbial infections in blind eels, as well as the latest research developments in this field. It is an invaluable reference for those wishing to conduct research or deepen their understanding in this field.

Query 2: Please describe the feeding patterns and pathogen exposure of lampreys at different life stages.

Output: According to the text, the feeding mode and pathogen exposure of different life stages of sevendill eels are as follows:

Larval stage (ammocoetes): sevendill larvae live mainly in sediments during this stage and ingest organic matter dissolved in river water, such as diatoms and other plankton, by filter feeding. This may limit their exposure to external pathogens.

Transformation Period: During the transformation process, sevendill eels stop eating and rely on their body reserves of nutrients for survival. Their transition from larvae to adults and their high metabolism during this period may make them more susceptible to pathogens.

Adult stage: Adults feed by sucking on the blood and muscle of the host fish via suckers. This para-

sitism brings them into direct contact with the body fluids and tissues of the host, increasing the risk of exposure to a wide range of pathogens. In particular, sevengill eels parasitize a wide range of fish species, greatly increasing their exposure to a variety of pathogens.

Migratory and spawning phases: during these phases, the sevengill eel essentially stops eating and once again relies primarily on its body reserves of nutrients for survival. Altered physiological status during these stages may affect their susceptibility to pathogens.

Overall, sevengill eels at different life stages vary greatly in their feeding patterns and lifestyles, which likely dictate different opportunities for exposure to and transmission of pathogens. The transition from filter-feeding in the larval stage to direct access to host body fluids and tissues in the adult stage, to relying primarily on internal reserves of nutrients again in the migratory and spawning stages, has a significant impact on the physiological state and immune system of the sevengill, and may have different implications for pathogen infection.

Query 3: Is there a difference in the risk of exposure to pathogens between female and male sevengill eels during the adult stage?

Output: According to the article, females and males of adult sevengill eels do not show a significant difference in the risk of infection with pathogens. Although the article mentions some examples of parasites and microorganisms infecting sevengill eels, it does not clearly indicate differences in infection risk between males and females. In addition, the abundance of these infection cases was analyzed and did not reflect differences in infection risk between males and females. Therefore, based on the results of the current study, no differences between males and females were found in the assessment of infection risk in sevengill eels. However, the article also mentions the need for comprehensive parasite and microbial monitoring of sevengill eels, and that more systematic monitoring and research is necessary for a more accurate assessment of infection risk.

Query 4: What are the parasites in a lamprey?

Output: According to the article, the parasites carried by seven-gill eels include at least the following:

****Circular Hookworm Order Polychaeta (Trematoda)**:** e.g. *Diplostomum* sp. corneal larvae, which primarily parasitize the eye.

****Cestoda (Cestoda)**:** *Triaenophorus crassus*, *Proteocephalus longicollis*, parasitic in the digestive tract.

****Echinodermata (Acanthocephala)**:** *Echinorhynchus salmonis*, parasitic in the digestive tract.

****Nematoda (Nematoda)**:** *Camallanus* sp., *Truttaedacnitis stelmioides*, parasitizes the abdominal cavity, liver, etc.

****Microsporidia**:** *Heterosporis* sp., parasitizes muscle tissue.

****Myxozoa (Myxozoa)**:** some species parasitize the kidneys.

****Parasitic Protozoa (Protozoa)**:** *Trichodina* spp., *Ergasilus* spp., parasitize gills and body surface, respectively.

****Fungi (Oomycetes)**:** *Saprolegnia* spp. that cause water mold.

****Acraniata (Acraniata)**:** Sea sevengill eels have been found to contain *Anisakis simplex* larvae, which may be a secondary host.

As mentioned above, sevengill eels can carry a wide range of parasites, mainly digestive tract parasites. However, it is important to note that the relationship between most of these parasites and disease in sevengill eels is unclear.