



SOEN 6011 : SOFTWARE ENGINEERING PROCESSES
SUMMER 2021

SUPER CALCULATOR

PROBLEM - 1
Function Description

Authors

Rokeya Begum Keya

Kyle Taylor Lange

Sijie Min

Manimaran Palani

Contents

a) Description of Functions	2
b) Context of Use Model	8

Description of Functions

PROBLEM 1 - F2: $\tan(x)$

SOEN 6011 - Summer 2021

Software Engineering Processes

Repository address : <https://github.com/Dakatsu/SOEN6011Calculator>

Rokeya Begum Keya

40183615


Tangent Function, $\tan(x)$

$\tan(x)$ is a trigonometric function which relate a right-angled triangle to ratios of two side lengths. In addition, it also uses to find the slope of a line. The tangent function is define as below:

$$\tan(x) = \frac{\sin(x)}{\cos(x)}$$

Graph

In the graph of a tangent function, there are no high or low points, so this function does not have an amplitude.



Graph of tangent function $y = \tan(x)$

Where, x is real number. The tangent function is undefined when $\cos(x) = 0$, therefore, tangent function has a vertical asymptote whenever $\cos(x) = 0$.

Range

The range of $\tan(x)$ is all real number (negative to positive infinity), except 0. The $\tan(x)$ does not have amplitude therefore, it has vertical asymptotes. As, tangent function increases and decreases without bound between vertical asymptotes, there is no horizontal asymptotes exist for it.

Domain and Co-domain

The domain of tangent function is all real number except the value where $\cos(x) = 0$ because if $\cos(x) = 0$ then $\tan(x)$ will be undefined. The co-domain of $\tan(x)$ is $(-\infty, +\infty)$.

PROBLEM 1 - F3: Hyperbolic Sine, $\sinh(x)$

SOEN 6011 - Summer 2021

Software Engineering Processes

Repository address : <https://github.com/Dakatsu/SOEN6011Calculator>

Kyle Taylor Lange

27627696

The function $\sinh(x)$ is known as the hyperbolic sine, and it has the following formula [1].

$$\sinh(x) \equiv \frac{1}{2}(e^x - e^{-x})$$

The character x is the only variable, as e is a constant roughly equivalent to 2.71828 [4]. Both the domain and co-domain of $\sinh(x)$ is \mathbb{R} , i.e. $(-\infty, +\infty)$. Positive values of x trend toward $+\infty$, negative values of x trend toward $-\infty$, and an input of 0 outputs 0.

Just as the trigonometric functions sine and cosine relate to a unit circle [3], their hyperbolic equivalents relate to the right branch of a unit hyperbola [2]. The x-coordinate of the right branch of the hyperbola at a given hyperbolic angle a can be calculated with $\cosh(a)$, and $\sinh(a)$ will return the y-coordinate [5].

Bibliography

- [1] Hyperbolic Sine, Wolfram MathWorld.
<http://mathworld.wolfram.com/HyperbolicSine.html>
- [2] Hyperbolic Functions, Wolfram MathWorld.
<http://mathworld.wolfram.com/HyperbolicFunctions.html>
- [3] Circular Functions, Wolfram MathWorld.
<http://mathworld.wolfram.com/CircularFunctions.html>
- [4] e , Wolfram MathWorld.
<http://mathworld.wolfram.com/e.html>
- [5] Rectangular Hyperbola, Wolfram MathWorld.
<https://mathworld.wolfram.com/RectangularHyperbola.html>

PROBLEM 1 - F*

SOEN 6011 - Summer 2021

Software Engineering Processes

Repository address : <https://github.com/Dakatsu/SOEN6011Calculator>

Sijie Min

401*****

Team please add your content here

PROBLEM 1 - F7 : x^y

SOEN 6011 - Summer 2021

Software Engineering Processes

Repository address : <https://github.com/Dakatsu/SOEN6011Calculator>

Manimaran Palani

40167543

Definition of x^y

[1] Exponentiation is a mathematical operation, denoted as x^y , If y is a positive integer and x is any real number, then x^y corresponds to repeated multiplication.

$$x^y = x * x * x * \dots * x \text{ of } y \text{ times}$$

The expression can be called as “ x raised to the power of y ,” “ x to the power of y ,” or simply “ x to the y .” Here, x is the base and y is the exponent or the power.

Domain

[2] All the real numbers from -infinite to +infinite. $(-\infty \text{ to } \infty)$

$$(x, y) \in \mathbb{R}^2 : (x \geq 0 \wedge y \neq 0) \vee x > 0$$

Co-domain


A set of all positive real numbers from zero to infinite $(0 \text{ to } \infty)$ is known as the Exponentiation function

Characteristics

Graph -

[2][3] The **rate of change** increases (or decreases) across the graph in an exponential graph.

- ★ The exponential graph crosses the y-axis at $(0,1)$.
- ★ The exponential graph increases, when $x > 1$.
- ★ The exponential graph decreases, when $0 < x < 1$.
- ★ The exponential graph is asymptotic to the x-axis - gets very, very close to the x-axis but, in this case, does not touch it or cross it.



Graph crosses the y-axis at $(0,1)$

Bibliography

- [1] Nykamp DQ: Basic rules for exponentiation
<https://mathbitsnotebook.com/Algebra2/Exponential/EXExpFunctions.html>
- [2] MathBits Teacher: Exponential Functions,
<https://mathbitsnotebook.com/Algebra2/Exponential/EXExpFunctions.html>
- [3] WolframAlpha: Domain of exponentiation function,
<https://www.wolframalpha.com/input/?i=x%5Ey>

Context of Use Model

