


Module 04: Interpolation

Starting Monday, February 10, 2014

Consider the population of Canada (in millions):

Year	1971	1976	1981	1986	1991	1996	2001	2006	2011
Pop	21.57	22.99	24.34	25.31	27.30	28.85	30.01	31.61	33.48



Interpolation

- Given points (x_i, y_i) , for $i=0:n$, find a function $f(x)$ such that $f(x_i) = y_i$.
- Why?
 - Answer questions about values in the range
 - Answer questions about derivatives

Interpolation

- Issues to consider regarding $f(x)$?
 - Cost of finding f
 - Difficulty of evaluation
 - Cost of evaluation of f
 - Continuity
 - Estimation of derivatives

→ We will use polynomials of degree:

$$p(x) = a_0 + a_1x + a_2x^2 + \cdots a_nx^n$$

Vandermonde Matrix

$$\begin{bmatrix} 1 & x_0 & x_0^2 & \cdots & x_0^n \\ 1 & x_1 & x_1^2 & \cdots & x_1^n \\ 1 & x_2 & x_2^2 & \cdots & x_2^n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & x_n & x_n^2 & \cdots & x_n^n \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} y_0 \\ y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} \rightarrow Va = y$$

It can be shown that

$$\det(V) = \prod_{0 \leq i < j \leq n} (x_i - x_j)$$

Note: $\det(V) = 0 \iff x_i = x_j$ for some $i \neq j$

So, this system has a solution for distinct x_j values.

Revisiting population

Consider the data:

Years since 1995	1	6	11	16
Pop (millions)	28.85	30.01	31.61	33.48

Vandermonde matrix

$$\begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 6 & 36 & 216 \\ 1 & 11 & 121 & 1331 \\ 1 & 16 & 256 & 4096 \end{bmatrix} \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix} = \begin{bmatrix} 28.85 \\ 30.01 \\ 31.61 \\ 33.48 \end{bmatrix}$$
$$p(x) = -0.000227x^3 + \mathbf{0.01288}x^2 + 0.1516x + 28.6858$$

Solving for coefficients a

- System has $n+1$ equations and $n+1$ unknowns
- Can solve using GEPP or find QR factors
- However,
 - Either approach required $O(n^3)$ operations
 - $\kappa_2(V)$ grows exponentially \rightarrow Ill-conditioned system
- Let's consider a faster approach which is not as sensitive to condition of V .

Motivating example

- Suppose we have points (x_0, y_0) and (x_1, y_1)
- Consider $p(x) = a_0 + a_1x$
 - $p(x_0) = a_0 + a_1x_0 = y_0$
 - $p(x_1) = a_0 + a_1x_1 = y_1$
- Solve the 2x2 system:
 - $a_0 = (y_1 - y_0)/(x_1 - x_0)$
 - $a_1 = (x_1y_0 - x_0y_1)/(x_1 - x_0)$

Rewriting $p(x)$ gives

$$p(x) = \frac{(x - x_1)}{(x_0 - x_1)} y_0 + \frac{(x - x_0)}{(x_1 - x_0)} y_1$$
$$p(x) = \ell_0(x) y_0 + \ell_1(x) y_1$$

where ℓ_0, ℓ_1 (degree-1 polynomials) are called the Lagrange polynomials

→ $p(x)$ is a sum of degree-1 polynomial

→ In this format, we don't need to solve a system of equations – just calculate the coefficients directly.

Lagrange polynomials for larger n

For $k=0:n$, define

$$\ell_k(\mathbf{x}) = \prod_{j=0, j \neq k}^n \frac{x - x_j}{x_k - x_j}$$

The interpolating polynomial of degree $n+1$ is given by

$$p_n(x) = \sum_{k=0}^n \ell_k(x) y_k$$

Lagrange basis

- Let $P_n(x) = \{y_n(x): y_n(x) \text{ is a poly of deg } \leq n\}$
- $P_n(x)$ is a vector space with the basis
$$B = \{1, x, x^2, \dots, x^n\}.$$
- It is not the only basis
$$B_\ell = \{\ell_0(x), \ell_1(x), \dots, \ell_n(x)\}$$
 is also a basis
- Any polynomial in $P_n(x)$ can be written as a linear combination of $\ell_k(x)$ in B_ℓ
- The interpolating polynomial is unique – same polynomial regardless of way it is calculated.
- How many flops are required?