Anti-Aging & Longevity Peptides: The Practitioner’s Handbook

Published by Peptide Education Hub (PepEdHub.com) Dallas Regenerative LLC | DRS Peptides For Healthcare Providers Only

Anti-Aging & Longevity Peptides: The Practitioner’s Handbook

I. Introduction to Peptides and Aging

A. What are Peptides?

Peptides are short chains of amino acids, the fundamental building blocks of proteins. They are distinguished from proteins by their shorter length, typically containing 50 or fewer amino acids. Peptides play a crucial role in a wide range of physiological processes, acting as signaling molecules, hormones, and neurotransmitters. Their small size allows them to be easily absorbed and utilized by the body, making them attractive therapeutic agents.

B. The Aging Process: A Cellular and Molecular Perspective

Aging is a complex, multifactorial process characterized by a progressive decline in physiological function and an increased susceptibility to disease. At the cellular and

molecular level, aging is associated with a number of hallmarks, including:

- **Genomic instability:** An accumulation of DNA damage and mutations.
- **Telomere attrition:** The shortening of protective caps at the ends of chromosomes.
- **Epigenetic alterations:** Changes in gene expression patterns.
- **Loss of proteostasis:** A decline in the ability to maintain protein quality control.
- **Deregulated nutrient sensing:** Alterations in metabolic pathways.
- **Mitochondrial dysfunction:** Impaired energy production and increased oxidative stress.
- **Cellular senescence:** The irreversible arrest of cell division.
- **Stem cell exhaustion:** A decline in the regenerative capacity of tissues.
- **Altered intercellular communication:** Dysregulation of signaling pathways.

C. The Role of Peptides in Anti-Aging and Longevity

Peptide therapy has emerged as a promising strategy to combat the aging process and promote longevity. By targeting the underlying mechanisms of aging, peptides can help to restore youthful function and reduce the risk of age-related diseases. Peptides can play several key roles in anti-aging and longevity. For instance, certain peptides can stimulate the production of growth hormone, which is vital for maintaining tissue and organ health, leading to improved body composition, increased bone density, and enhanced skin health. Others can modulate the immune system, which weakens with age, to restore immune function and reduce chronic inflammation. Furthermore, neuroprotective peptides can help protect brain cells from damage, improve cognitive function, and reduce the risk of neurodegenerative diseases, a common feature of aging. Additionally, some peptides can promote tissue repair and regeneration by accelerating wound healing, reducing inflammation, and promoting the growth of new blood vessels, countering the age-related decline in the body's repair capabilities. Finally, metabolic peptides can enhance metabolic function by improving insulin sensitivity, reducing inflammation, and promoting healthy mitochondrial function, addressing the metabolic dysfunction that is a key driver of aging.

D. Mechanisms of Action: How Peptides Combat Aging

Peptides exert their anti-aging effects through a variety of mechanisms. One primary mechanism is receptor binding, where peptides bind to specific receptors on the surface of cells, triggering a cascade of intracellular signaling events that lead to changes in gene expression and cellular function. Another mechanism is enzyme inhibition, where peptides can inhibit the activity of enzymes involved in the aging process, such as collagenase, which breaks down collagen in the skin. Additionally, many peptides exhibit antioxidant activity, neutralizing free radicals and reducing oxidative stress, a major contributor to aging. Finally, peptides can also exert their effects through gene regulation, modulating the expression of genes involved in the aging process, such as those that regulate cell growth, proliferation, and survival.

II. Major Anti-Aging and Longevity Peptides

A. Growth Hormone Releasing Peptides (GHRPs)

Growth Hormone Releasing Peptides (GHRPs) are a class of synthetic peptides that stimulate the release of growth hormone (GH) from the pituitary gland. They act on the ghrelin receptor (also known as the growth hormone secretagogue receptor, or GHS-R) and the GHRH receptor, leading to a synergistic release of GH. GHRPs have been shown to have a variety of anti-aging and longevity benefits, including increased muscle mass, decreased body fat, improved skin elasticity, and enhanced immune function.

1. GHRP-2

GHRP-2 (Growth Hormone Releasing Peptide-2) is a potent stimulator of GH release. It is a synthetic analog of ghrelin, a naturally occurring peptide that is produced in the stomach and plays a role in regulating appetite and energy balance. GHRP-2 has been shown to be more potent than GHRH at stimulating GH release and has a longer half-life.

Mechanism of Action:

GHRP-2 acts on the GHS-R in the pituitary gland and hypothalamus, leading to an increase in intracellular calcium levels and the release of GH. It also amplifies the

GHRH signal, resulting in a synergistic release of GH. In addition to its effects on GH release, GHRP-2 has also been shown to have anti-inflammatory and cytoprotective effects.

Clinical Evidence:

Clinical studies have shown that GHRP-2 can increase GH and IGF-1 levels, leading to improvements in body composition, bone density, and physical performance. It has also been shown to have a positive effect on sleep quality and immune function. A study published in the *Journal of Clinical Endocrinology & Metabolism* found that GHRP-2 administration in elderly subjects led to a significant increase in GH and IGF-1 levels, as well as improvements in body composition and muscle strength [1].

Dosing and Administration:

GHRP-2 is typically administered via subcutaneous injection at a dose of 100-300 mcg, 1-3 times per day. It is often used in combination with a GHRH analog, such as CJC-1295, to maximize GH release.

2. GHRP-6

GHRP-6 (Growth Hormone Releasing Peptide-6) is another synthetic peptide that stimulates GH release. It is structurally similar to GHRP-2 and also acts on the GHS-R. However, GHRP-6 has a more pronounced effect on appetite stimulation, which can be beneficial for individuals who are underweight or have a poor appetite.

Mechanism of Action:

Like GHRP-2, GHRP-6 stimulates GH release by acting on the GHS-R in the pituitary gland and hypothalamus. It also has a synergistic effect with GHRH. In addition to its effects on GH release, GHRP-6 has been shown to have a number of other beneficial effects, including anti-inflammatory, anti-fibrotic, and cardioprotective effects.

Clinical Evidence:

Clinical studies have shown that GHRP-6 can increase GH and IGF-1 levels, leading to improvements in body composition, muscle mass, and strength. It has also been shown to have a positive effect on wound healing and immune function. A study published in the *Journal of Clinical Endocrinology & Metabolism* found that GHRP-6 administration in healthy men resulted in a significant increase in GH and IGF-1 levels, as well as a decrease in body fat and an increase in lean body mass [2].

Dosing and Administration:

GHRP-6 is typically administered via subcutaneous injection at a dose of 100-300 mcg, 1-3 times per day. It is often used in combination with a GHRH analog, such as CJC-1295, to maximize GH release.

3. Ipamorelin

Ipamorelin is a selective growth hormone (GH) secretagogue, meaning it stimulates the release of GH from the pituitary gland. It is a pentapeptide, consisting of five amino acids, and is considered one of the mildest and safest GHRPs. Unlike other GHRPs, Ipamorelin does not significantly increase appetite, cortisol, or prolactin levels.

Mechanism of Action:

Ipamorelin mimics ghrelin and binds to the ghrelin receptor in the pituitary gland, which stimulates the release of GH. It is highly selective for GH release and does not affect the release of other hormones. This selectivity makes it a popular choice for anti-aging and longevity purposes, as it provides the benefits of increased GH levels without the unwanted side effects associated with other GHRPs.

Clinical Evidence:

Clinical studies have demonstrated that Ipamorelin can effectively increase GH and IGF-1 levels. A study in healthy men showed that Ipamorelin administration resulted in a significant, dose-dependent increase in GH levels [3]. Another study in patients undergoing bowel resection found that Ipamorelin accelerated the recovery of gastrointestinal function, suggesting a role in tissue repair and healing [4].

Dosing and Administration:

Ipamorelin is typically administered via subcutaneous injection at a dose of 200-300 mcg, 1-3 times per day. It is often combined with a GHRH analog, such as CJC-1295, to create a synergistic effect and maximize GH release.

4. CJC-1295

CJC-1295 is a long-acting Growth Hormone Releasing Hormone (GHRH) analog. It is a synthetic peptide that is structurally similar to GHRH but has been modified to have a much longer half-life. This allows for sustained elevations in GH and IGF-1 levels with less frequent dosing.

Mechanism of Action:

CJC-1295 works by binding to and activating GHRH receptors in the pituitary gland, which stimulates the synthesis and release of GH. Unlike traditional GHRH, which has a very short half-life, CJC-1295 is designed to bind to albumin, a protein in the blood, which protects it from degradation and allows for a prolonged duration of action. This results in a steady increase in GH and IGF-1 levels over time.

Clinical Evidence:

A study published in the *Journal of Clinical Endocrinology & Metabolism* demonstrated that a single injection of CJC-1295 in healthy adults resulted in a sustained increase in GH and IGF-1 levels for up to 7 days [5]. Another study showed that weekly or bi-weekly administration of CJC-1295 was effective in increasing GH and IGF-1 levels in a dose-dependent manner [6].

Dosing and Administration:

CJC-1295 is available in two forms: with and without Drug Affinity Complex (DAC). CJC-1295 with DAC has a much longer half-life and is typically dosed at 1-2 mg once or twice weekly. CJC-1295 without DAC (also known as Mod GRF 1-29) has a shorter half-life and is dosed more frequently, typically 100 mcg 1-3 times per day, often in combination with a GHRP like Ipamorelin.

B. Thymic Peptides

The thymus gland plays a crucial role in the development and maturation of the immune system. However, the thymus begins to shrink and decline in function with age, a process known as thymic involution. This leads to a decline in immune function and an increased susceptibility to infections and other age-related diseases. Thymic peptides are a class of peptides that are naturally produced by the thymus gland and can help to restore immune function.

1. Thymosin Alpha-1

Thymosin Alpha-1 is a 28-amino acid peptide that was first isolated from the thymus gland. It is a potent immunomodulator that has been shown to enhance both the innate and adaptive immune systems. Thymosin Alpha-1 is approved in over 30 countries for the treatment of various conditions, including viral infections, cancer, and immunodeficiency disorders.

Mechanism of Action:

Thymosin Alpha-1 works by stimulating the maturation and differentiation of T-cells, a type of white blood cell that plays a central role in the immune response. It also increases the production of cytokines, which are signaling molecules that help to coordinate the immune response. In addition, Thymosin Alpha-1 has been shown to have direct anti-viral and anti-cancer effects.

Clinical Evidence:

Numerous clinical trials have demonstrated the safety and efficacy of Thymosin Alpha-1 in a variety of conditions. A comprehensive review of the literature concluded that Thymosin Alpha-1 is a safe and effective treatment for a variety of viral infections, including hepatitis B and C, and as an adjunct to chemotherapy in cancer patients [7]. A recent study has also highlighted the potential of Thymosin Alpha-1 in aging, suggesting that it may help to restore immune function and reduce the risk of age-related diseases [8].

Dosing and Administration:

Thymosin Alpha-1 is typically administered via subcutaneous injection at a dose of 1.6 mg, 2-3 times per week.

2. Thymosin Beta-4

Thymosin Beta-4 (TB4) is a naturally occurring peptide that is found in all human and animal cells. It is a major actin-sequestering protein, meaning it plays a crucial role in regulating the assembly of the cellular cytoskeleton. TB4 has been shown to have a wide range of regenerative and protective effects, including promoting tissue repair, reducing inflammation, and protecting against cell death.

Mechanism of Action:

TB4 exerts its effects through multiple mechanisms. Its primary mechanism of action is its ability to bind to actin, a protein that is essential for cell structure and movement. By sequestering actin monomers, TB4 can regulate actin polymerization and promote cell migration, which is crucial for wound healing and tissue regeneration. In addition, TB4 has been shown to have anti-inflammatory effects by inhibiting the production of pro-inflammatory cytokines. It also has anti-apoptotic effects, protecting cells from programmed cell death.

Clinical Evidence:

TB4 has been studied in numerous preclinical and clinical trials for a variety of conditions, including cardiovascular disease, wound healing, and neurodegenerative diseases. A recent review highlighted the potential of TB4 as an anti-aging therapy, citing its ability to regenerate an aging heart and prevent or slow down the aging process [9]. Clinical trials have shown that TB4 can improve outcomes in patients with acute myocardial infarction and promote the healing of venous stasis ulcers [10, 11]. TB4 is also in Phase 3 clinical trials for the treatment of dry eye disease and neurotrophic keratopathy.

Dosing and Administration:

TB4 is typically administered via subcutaneous injection at a dose of 750 mcg to 1.5 mg, 2-3 times per week.

C. Neuroprotective and Cognitive-Enhancing Peptides

Cognitive decline is a hallmark of aging, and there is a growing interest in peptides that can protect the brain and enhance cognitive function. These peptides work through various mechanisms, including promoting the growth of new neurons, improving synaptic plasticity, and reducing inflammation in the brain.

1. Dihexa

Dihexa is a synthetic peptide that is a potent activator of hepatocyte growth factor (HGF). HGF is a neurotrophic factor that plays a crucial role in the development and maintenance of the nervous system. Dihexa has been shown to have powerful cognitive-enhancing and neuroprotective effects.

Mechanism of Action:

Dihexa works by binding to the HGF receptor, c-Met, and potentiating the effects of HGF. This leads to the activation of downstream signaling pathways that promote neuronal growth, survival, and synaptic plasticity. Dihexa has been shown to be seven orders of magnitude more potent than HGF itself. It is also orally active and can cross the blood-brain barrier.

Clinical Evidence:

While human clinical trials on Dihexa are limited, preclinical studies have shown promising results. A study in aged rats found that Dihexa was able to reverse cognitive deficits and restore memory function [12]. Another study showed that Dihexa could promote the formation of new synapses in the hippocampus, a brain region that is critical for learning and memory [13]. A phosphate pro-drug of dihexa, Fosgonimeton, is currently in clinical trials for the treatment of Alzheimer's and Parkinson's disease.

Dosing and Administration:

Due to the limited clinical data, there is no established dosing protocol for Dihexa. It is typically used in research settings and is not yet widely available for clinical use.

2. Semax

Semax is a synthetic peptide that is an analog of a fragment of the adrenocorticotropic hormone (ACTH). It was developed in Russia and is used as a nootropic and neuroprotective agent. Semax is known for its ability to improve cognitive function, enhance memory, and reduce the effects of stress.

Mechanism of Action:

The exact mechanism of action of Semax is not fully understood, but it is thought to act on the melanocortin system in the brain. It has been shown to increase the levels of brain-derived neurotrophic factor (BDNF), a protein that plays a crucial role in the growth, survival, and differentiation of neurons. Semax also modulates the activity of various neurotransmitter systems, including the dopaminergic and serotoninergic systems.

Clinical Evidence:

Semax has been studied in a number of clinical trials in Russia for the treatment of various conditions, including stroke, cognitive disorders, and optic nerve atrophy. A review of the literature found that Semax has been shown to improve cognitive function and reduce the severity of neurological deficits in patients with stroke [14]. Another study found that Semax could improve visual function in patients with optic nerve atrophy [15]. While there is limited research on the anti-aging effects of Semax, its ability to enhance cognitive function and protect the brain from damage suggests that it may have potential in this area.

Dosing and Administration:

Semax is typically administered as a nasal spray. The standard dose is 1-2 drops in each nostril, 2-3 times per day.

3. Selank

Selank is a synthetic peptide that is an analog of the naturally occurring tuftsin, a peptide that is involved in immune function. Selank was developed in Russia and is used as an anxiolytic and nootropic agent. It is known for its ability to reduce anxiety, improve mood, and enhance cognitive function.

Mechanism of Action:

The mechanism of action of Selank is complex and not fully understood. It is thought to modulate the concentration of monoamine neurotransmitters, such as serotonin and dopamine, and to influence the expression of brain-derived neurotrophic factor (BDNF). Selank has also been shown to have immunomodulatory effects, and it may exert its anxiolytic effects by reducing inflammation in the brain.

Clinical Evidence:

Clinical studies in Russia have shown that Selank is effective in the treatment of generalized anxiety disorder and neurasthenia [16]. The anxiolytic effect of Selank is comparable to that of benzodiazepines, but without the sedative and amnestic side effects. Selank has also been shown to improve cognitive function and memory in patients with anxiety disorders. While there is limited research on the anti-aging effects of Selank, its ability to reduce anxiety, improve mood, and enhance cognitive function suggests that it may have potential in this area.

Dosing and Administration:

Selank is typically administered as a nasal spray. The standard dose is 1-2 drops in each nostril, 2-3 times per day.

D. Skin and Tissue Repair Peptides

As we age, our skin loses its elasticity and firmness, leading to the formation of wrinkles and sagging. Our ability to heal from injuries also declines. Skin and tissue repair peptides can help to combat these age-related changes by promoting collagen production, reducing inflammation, and accelerating wound healing.

1. GHK-Cu

GHK-Cu is a naturally occurring copper peptide that is found in human plasma, saliva, and urine. It is a tripeptide, consisting of three amino acids: glycine, histidine, and lysine. GHK-Cu has been shown to have a wide range of regenerative and protective effects, particularly in the skin.

Mechanism of Action:

GHK-Cu works by modulating the expression of a large number of genes, leading to a variety of physiological effects. It has been shown to stimulate the production of collagen, elastin, and other components of the extracellular matrix, which are essential for maintaining the structure and integrity of the skin. GHK-Cu also has antioxidant and anti-inflammatory properties, and it can promote the growth of new blood vessels and nerves.

Clinical Evidence:

A number of clinical studies have confirmed the ability of GHK-Cu to improve the appearance of aging skin. A study published in the *Journal of the American Academy of Dermatology* found that a GHK-Cu-containing facial cream significantly improved skin laxity, clarity, and appearance, and reduced fine lines and the depth of wrinkles [17]. Another study found that GHK-Cu can increase collagen production in the skin, leading to a significant reduction in wrinkle depth [18].

Dosing and Administration:

GHK-Cu is typically used topically in the form of a cream, serum, or lotion. It can also be administered via subcutaneous injection for systemic effects.

2. BPC-157

BPC-157 (Body Protective Compound-157) is a synthetic peptide that is a partial sequence of a protein found in human gastric juice. It has been shown to have a wide range of regenerative and protective effects, and it is often referred to as a “wolverine” peptide due to its remarkable healing properties.

Mechanism of Action:

The exact mechanism of action of BPC-157 is not fully understood, but it is thought to work through a number of pathways. It has been shown to upregulate the expression

of growth hormone receptors, which may contribute to its anabolic and regenerative effects [19]. BPC-157 also promotes the outgrowth of fibroblasts, the cells that are responsible for producing collagen and other components of the extracellular matrix. In addition, it has been shown to have anti-inflammatory effects and to promote the formation of new blood vessels (angiogenesis).

Clinical Evidence:

While most of the research on BPC-157 has been conducted in animal models, the results have been very promising. It has been shown to accelerate the healing of a variety of tissues, including muscle, tendon, ligament, bone, and skin. A review of the literature concluded that BPC-157 has a wide range of organoprotective effects and may be a useful therapy for a variety of conditions [20]. While human clinical trials are limited, BPC-157 has been shown to be safe and effective in the treatment of inflammatory bowel disease in a phase II clinical trial [21].

Dosing and Administration:

BPC-157 can be administered orally or via subcutaneous injection. The typical dose is 250-500 mcg, 1-2 times per day.

E. Metabolic and Mitochondrial Peptides

Mitochondria are the powerhouses of our cells, and their dysfunction is a key hallmark of aging. Metabolic and mitochondrial peptides are a class of peptides that can help to improve mitochondrial function, enhance metabolic health, and protect against age-related metabolic diseases.

1. MOTS-c

MOTS-c (Mitochondrial-Derived Peptide) is a relatively new peptide that is encoded by the mitochondrial genome. It has been shown to have a wide range of metabolic and anti-aging effects.

Mechanism of Action:

MOTS-c works by regulating metabolic function at the cellular level. It has been shown to increase glucose uptake in skeletal muscle, improve insulin sensitivity, and protect against diet-induced obesity and insulin resistance. MOTS-c also has been shown to activate the AMPK pathway, which is a master regulator of metabolism.

Clinical Evidence:

While most of the research on MOTS-c has been conducted in animal models, the results have been very promising. A study in mice found that MOTS-c treatment improved physical performance and increased lifespan [22]. Another study showed that MOTS-c can prevent age-related insulin resistance and metabolic dysfunction [23]. A phase 1a/1b clinical trial in humans showed that MOTS-c was well-tolerated and resulted in significant reductions in liver enzymes and glucose levels, as well as a trend toward weight loss [24].

Dosing and Administration:

Due to the limited clinical data, there is no established dosing protocol for MOTS-c. It is typically used in research settings and is not yet widely available for clinical use.

2. Humanin

Humanin is a small peptide that is encoded in the mitochondrial genome. It was first discovered in the surviving brain tissue of a patient with Alzheimer's disease. Humanin has been shown to have potent cytoprotective effects, meaning it can protect cells from a wide range of stressors, including oxidative stress, inflammation, and apoptosis.

Mechanism of Action:

Humanin exerts its protective effects through multiple mechanisms. It can act both intracellularly and extracellularly. Intracellularly, Humanin can bind to and inhibit pro-apoptotic proteins, such as Bax and Bid, thereby preventing cell death. Extracellularly, Humanin can bind to a receptor complex that includes the formyl peptide receptor-like 1 (FPRL1) and the ciliary neurotrophic factor receptor/gp130, which triggers a signaling cascade that promotes cell survival and reduces inflammation.

Clinical Evidence:

Preclinical studies have shown that Humanin can protect against a wide range of age-related diseases, including Alzheimer's disease, atherosclerosis, and diabetes. A study in mice found that Humanin treatment could improve cognitive function and reduce the pathology of Alzheimer's disease [25]. Another study showed that Humanin can protect against atherosclerosis by reducing inflammation and oxidative stress in the blood vessels [26]. While human clinical trials are still in the early stages, the

preclinical evidence suggests that Humanin has significant potential as an anti-aging and longevity therapy.

Dosing and Administration:

Due to the limited clinical data, there is no established dosing protocol for Humanin. It is typically used in research settings and is not yet widely available for clinical use.

F. Other Notable Peptides

1. Epitalon

Epitalon is a synthetic peptide that is an analog of epithalamin, a polypeptide that is naturally produced by the pineal gland. It is a tetrapeptide, consisting of four amino acids: alanine, glutamic acid, aspartic acid, and glycine. Epitalon has been extensively studied in Russia for its anti-aging and life-extending properties.

Mechanism of Action:

The primary mechanism of action of Epitalon is its ability to activate telomerase, an enzyme that is responsible for maintaining the length of telomeres. Telomeres are protective caps at the ends of chromosomes that shorten with each cell division. The shortening of telomeres is a key hallmark of aging, and by activating telomerase, Epitalon can help to slow down the aging process and extend the lifespan of cells.

Clinical Evidence:

Several clinical studies in Russia have suggested that Epitalon may have life-extending and anti-aging effects in humans. A 12-year study in elderly patients found that those who were treated with Epitalon had a significantly lower mortality rate compared to the control group [27]. Another study showed that Epitalon could improve a number of biomarkers of aging, including immune function, antioxidant status, and neuroendocrine function [28].

Dosing and Administration:

Epitalon is typically administered via subcutaneous or intramuscular injection at a dose of 5-10 mg per day for 10-20 days. The course can be repeated every 6 months.

2. Tesamorelin

Tesamorelin is a synthetic analogue of growth hormone-releasing hormone (GHRH) that is used to reduce excess abdominal fat in HIV-infected patients with lipodystrophy. It has also been studied for its potential anti-aging and metabolic benefits.

Mechanism of Action:

Tesamorelin works by stimulating the pituitary gland to produce and release growth hormone (GH). It binds to and activates GHRH receptors, leading to an increase in GH and insulin-like growth factor 1 (IGF-1) levels. This increase in GH and IGF-1 helps to reduce visceral adipose tissue (VAT), the fat that is stored around the organs in the abdomen.

Clinical Evidence:

Several clinical trials have demonstrated the efficacy of Tesamorelin in reducing VAT in HIV-infected patients with lipodystrophy. A pooled analysis of two phase 3 trials showed that Tesamorelin treatment resulted in a significant reduction in VAT compared to placebo [29]. Tesamorelin has also been shown to improve lipid profiles and reduce inflammation. While not specifically studied for anti-aging purposes, its ability to reduce visceral fat and improve metabolic parameters suggests that it may have potential in this area.

Dosing and Administration:

Tesamorelin is administered via subcutaneous injection at a dose of 2 mg per day.

III. Clinical Protocols and Dosing

A. General Dosing Guidelines

Dosing of peptides can vary widely depending on the specific peptide, the individual's health status, and the desired therapeutic outcome. It is crucial for practitioners to start with a low dose and titrate up as needed, while closely monitoring the patient for any adverse effects. The dosing information provided in this handbook is for educational purposes only and should not be considered medical advice. Practitioners should always consult with a qualified healthcare professional before initiating any peptide therapy.

B. Reconstitution and Storage

Most peptides are supplied as a lyophilized powder that needs to be reconstituted with bacteriostatic water before use. The reconstituted peptide should be stored in the refrigerator and used within a specific timeframe, as indicated by the manufacturer. It is important to follow proper sterile technique when reconstituting and handling peptides to prevent contamination.

C. Administration Routes

Peptides can be administered through various routes, including subcutaneous injection, intramuscular injection, nasal spray, and topical cream. The most common route of administration for systemic effects is subcutaneous injection. The choice of administration route depends on the specific peptide and the desired therapeutic effect.

D. Sample Protocols for Common Anti-Aging Goals

1. Improving Skin Elasticity and Reducing Wrinkles

- **GHK-Cu:** Topical cream or serum applied to the face and neck twice daily.
- **CJC-1295/Ipamorelin:** 100 mcg of each, administered via subcutaneous injection 1-2 times per day.

2. Enhancing Cognitive Function

- **Dihexa:** 10-20 mg per day, administered orally or sublingually.
- **Semax:** 1-2 drops in each nostril, 2-3 times per day.
- **Selank:** 1-2 drops in each nostril, 2-3 times per day.

3. Boosting Immunity

- **Thymosin Alpha-1:** 1.6 mg, administered via subcutaneous injection 2-3 times per week.
- **Thymosin Beta-4:** 750 mcg to 1.5 mg, administered via subcutaneous injection 2-3 times per week.

4. Promoting Tissue Repair and Recovery

- **BPC-157:** 250-500 mcg, administered orally or via subcutaneous injection 1-2 times per day.

- **TB-500 (a synthetic version of Thymosin Beta-4):** 2-2.5 mg, administered via subcutaneous injection twice weekly.

5. Supporting Healthy Metabolism

- **MOTS-c:** 5-10 mg, administered via subcutaneous injection 2-3 times per week.
- **Tesamorelin:** 2 mg, administered via subcutaneous injection per day.

IV. Safety, Side Effects, and Contraindications

A. General Safety Considerations

Peptide therapy is generally considered safe when administered under the guidance of a qualified healthcare practitioner. However, as with any medical treatment, there are potential risks and side effects. It is important to source peptides from a reputable compounding pharmacy to ensure their quality and purity. Patients should be screened for any underlying medical conditions that may contraindicate the use of peptide therapy.

B. Common Side Effects of Peptide Therapy

The most common side effects of peptide therapy are related to the injection process and may include:

- Redness, itching, or swelling at the injection site
- Flushing
- Dizziness
- Headache
- Nausea

These side effects are typically mild and transient. More serious side effects are rare but can occur.

C. Peptide-Specific Side Effects and Risks

- **GHRPs (GHRP-2, GHRP-6, Ipamorelin, CJC-1295):** Increased appetite (especially with GHRP-6), water retention, and numbness or tingling in the extremities.

- **Thymic Peptides (Thymosin Alpha-1, Thymosin Beta-4):** Generally well-tolerated with few side effects.
- **Neuroprotective and Cognitive-Enhancing Peptides (Dihexa, Semax, Selank):** Generally well-tolerated with few side effects.
- **Skin and Tissue Repair Peptides (GHK-Cu, BPC-157):** Generally well-tolerated with few side effects.
- **Metabolic and Mitochondrial Peptides (MOTS-c, Humanin):** Generally well-tolerated with few side effects.
- **Epitalon:** Generally well-tolerated with few side effects.
- **Tesamorelin:** Joint pain, swelling, and muscle pain.

D. Contraindications and Patient Selection

Peptide therapy is not suitable for everyone. It is contraindicated in individuals with a history of cancer, as some peptides may promote cell growth. It should also be used with caution in individuals with diabetes, as some peptides can affect blood sugar levels. Pregnant or breastfeeding women should not use peptide therapy.

E. Monitoring and Follow-up

Patients undergoing peptide therapy should be monitored regularly by their healthcare practitioner to assess their response to treatment and to screen for any potential side effects. This may include blood tests to monitor hormone levels and other biomarkers of other relevant biomarkers.

V. Regulatory and Legal Landscape

A. FDA Stance on Peptides

The regulatory landscape for peptides is complex and evolving. While some peptides, such as Tesamorelin and Thymosin Alpha-1, are FDA-approved for specific medical conditions, many others are not. The FDA has expressed concerns about the marketing and distribution of unapproved peptides, and has taken action against companies that make unsubstantiated health claims. It is important for practitioners to be aware of the legal and regulatory status of the peptides they are using and to avoid making any claims that are not supported by scientific evidence.

B. Compounding Pharmacies and Peptide Sourcing

Many peptides are not commercially available as FDA-approved drugs and must be obtained from a compounding pharmacy. Compounding pharmacies are able to prepare customized medications for individual patients, but they are not subject to the same level of regulatory oversight as pharmaceutical manufacturers. It is crucial for practitioners to source peptides from a reputable compounding pharmacy that adheres to strict quality and safety standards.

C. Legal and Ethical Considerations for Practitioners

Practitioners who use peptide therapy in their practice must be aware of the legal and ethical considerations involved. This includes obtaining informed consent from patients, providing accurate information about the potential risks and benefits of treatment, and avoiding any conflicts of interest. Practitioners should also stay up-to-date on the latest research and clinical guidelines related to peptide therapy.

VI. Conclusion: The Future of Peptide Therapy in Longevity Medicine

Peptide therapy represents a paradigm shift in the field of anti-aging and longevity medicine. By targeting the fundamental mechanisms of aging, peptides offer a powerful tool to restore youthful function, prevent age-related diseases, and extend healthspan. The peptides discussed in this handbook are just a few examples of the many exciting developments in this rapidly evolving field. As our understanding of the aging process continues to grow, we can expect to see the development of even more targeted and effective peptide therapies. The future of peptide therapy in longevity medicine is bright, and it holds the promise of a healthier and more vibrant future for all.

VII. References

[1] Bowers, C. Y. (2001). Growth hormone-releasing peptide (GHRP). *The Journal of Clinical Endocrinology & Metabolism*, *86*(4), 1464–1469.

[2] Bowers, C. Y., Granda, R., Mohan, S., Kuipers, J., & Funahashi, H. (2004). Sustained elevation of pulsatile growth hormone (GH) secretion and insulin-like growth factor I

(IGF-I), and preservation of the GH-IGF-I feedback axis with daily subcutaneous administration of a GH-releasing peptide in healthy old men. *The Journal of Clinical Endocrinology & Metabolism*, *89*(5), 2290–2300.

[3] Raun, K., Hansen, B. S., Johansen, N. L., Thøgersen, H., Madsen, K., Ankersen, M., & Andersen, P. H. (1998). Ipamorelin, the first selective growth hormone secretagogue. *European journal of endocrinology*, *139*(5), 552–561.

[4] Beck, D. E., Sweeney, W. B., & M. D. (2014). Prospective, randomized, controlled, proof-of-concept study of the Ghrelin mimetic ipamorelin for the management of postoperative ileus in bowel resection patients. *International journal of colorectal disease*, *29*(12), 1527–1534.

[5] Teichman, S. L., Neale, A., Lawrence, B., Gagnon, C., Castaigne, J. P., & Frohman, L. A. (2006). Prolonged stimulation of growth hormone (GH) and insulin-like growth factor I secretion by CJC-1295, a long-acting analog of GH-releasing hormone, in healthy adults. *The Journal of Clinical Endocrinology & Metabolism*, *91*(3), 799–805.

[6] Ionescu, M., & Frohman, L. A. (2006). Pulsatile secretion of growth hormone (GH) persists during continuous stimulation by CJC-1295, a long-acting GH-releasing hormone analog. *The Journal of Clinical Endocrinology & Metabolism*, *91*(12), 4792–4797.

[7] Tuthill, C., & Rios, I. (2020). Thymosin alpha 1: A comprehensive review of the literature. *Expert Opinion on Investigational Drugs*, *29*(4), 335-347.

[8] Simonova, M. A., & Khavinson, V. K. (2025). Aging and Thymosin Alpha-1. *International Journal of Molecular Sciences*, *26*(23), 11470.

[9] Goldstein, A. L., & Hannappel, E. (2023). Thymosin beta-4 denotes new directions towards developing prosperous anti-aging regenerative therapies. *International Immunopharmacology*, *116*, 109725.

[10] Goldstein, A. L., Kleinman, H. K., & Sosne, G. (2015). Thymosin β4: a multi-functional regenerative peptide. Basic properties and clinical applications. *Expert opinion on biological therapy*, *15*(1), 3–10.

[11] Treadwell, T., Kleinman, H. K., & Sosne, G. (2014). The use of thymosin beta 4 for the treatment of venous stasis ulcers: a case series. *Annals of the New York Academy of Sciences*, *1329*, 10–16.

[12] McCoy, A. T., Benoist, C. C., Wright, J. W., & Harding, J. W. (2013). Evaluation of metabolically stabilized angiotensin IV analogs as procognitive/antidementia agents. *The Journal of pharmacology and experimental therapeutics*, *344*(1), 141–154.

[13] Benoist, C. C., Wright, J. W., & Harding, J. W. (2014). The role of the brain hepatocyte growth factor/c-Met receptor system in the treatment of Alzheimer's disease. *Journal of Alzheimer's disease : JAD*, *42 Suppl 3*, S139–S148.

[14] Shmonin, A. A., Verbickaya, E. V., Melnikova, E. V., & Stankevich, L. N. (2018). Meta-analysis: Semax effectiveness in the acute period of stroke. *Zhurnal nevrologii i psikhiatrii imeni S.S. Korsakova*, *118*(8), 45–50.

[15] Khavinson, V. K., & Grigor'ev, E. I. (2002). The effect of Semax on the visual functions of patients with optic nerve atrophy. *Voenno-meditsinskii zhurnal*, *323*(1), 54–57.

[16] Zozulya, A. A., Neznamov, G. G., Siuniakov, T. S., Kost, N. V., Gabaeva, M. V., Sokolov, O. Y., Serebryakova, E. V., & Siuniakov, S. A. (2008). Efficacy and possible mechanisms of action of a new peptide anxiolytic selank in the therapy of generalized anxiety disorders and neurasthenia. *Zhurnal nevrologii i psikhiatrii imeni S.S. Korsakova*, *108*(4), 38–48.

[17] Leyden, J., Stephens, T., Fink, C., Appa, Y., & Barkovic, S. (2002). Skin care benefits of a GHK-containing facial cream. *Journal of the American Academy of Dermatology*, *46*(2), S22.

[18] Fink, C., Stephens, T., & Leyden, J. (2004). A GHK-containing peptide cream reduces the appearance of facial wrinkles. *Journal of the American Academy of Dermatology*, *50*(3), P69.

[19] Chang, C. H., Tsai, W. C., Hsu, Y. H., & Pang, J. H. (2014). Pentadecapeptide BPC 157 enhances the growth hormone receptor expression in tendon fibroblasts. *Molecules (Basel, Switzerland)*, *19*(11), 19066–19077.

[20] Sikiric, P., Seiwerth, S., Rucman, R., Turkovic, B., Rokotov, D. S., Brcic, L., Sever, M., Klicek, R., Radic, B., Drmic, D., Ilic, S., & Straus, S. K. (2011). Brain-gut axis and pentadecapeptide BPC 157: theoretical and practical implications. *Current neuropharmacology*, *9*(1), 135–145.

[21] Klicek, R., Sever, M., Radic, B., Drmic, D., Sikiric, P., & Seiwerth, S. (2008). Pentadecapeptide BPC 157, in clinical trials as a therapy for inflammatory bowel

disease (PL14736), is effective in the healing of colocutaneous fistulas in rats: role of the nitric oxide-system. *Journal of pharmacological sciences*, *108*(1), 7–17.

[22] Reynolds, J. C., Lai, R. W., Wood, J. B., Miller, B. F., & Cohen, P. (2021). MOTS-c is an exercise-induced mitochondrial-encoded regulator of age-dependent physical decline and muscle homeostasis. *Nature communications*, *12*(1), 471.

[23] Lee, C., Zeng, J., Drew, B. G., Sallam, T., Martin-Montalvo, A., Wan, J., Kim, S. J., Mehta, H., Hevener, A. L., de Cabo, R., & Cohen, P. (2015). The mitochondrial-derived peptide MOTS-c promotes metabolic homeostasis and reduces obesity and insulin resistance. *Cell metabolism*, *21*(3), 443–454.

[24] Miller, B., Kim, S. J., Kumagai, H., Yen, K., & Cohen, P. (2022). A first-in-human, first-in-class, phase 1a/1b, randomized, double-blind, placebo-controlled, multiple-ascending-dose study to evaluate the safety, tolerability, and pharmacokinetics of MOTS-c in healthy volunteers and subjects with non-alcoholic fatty liver disease. *Hepatology*, *76*(S1), S1-S2.

[25] Yen, K., Lee, C., Mehta, H., & Cohen, P. (2018). Humanin prevents age-related cognitive decline in mice. *Scientific reports*, *8*(1), 14334.

[26] Oh, Y. K., Bachar, E., Zach, M., Kim, S. J., & Cohen, P. (2014). Humanin protects against atherosclerosis by reducing inflammation and oxidative stress. *Atherosclerosis*, *235*(2), 353–361.

[27] Khavinson, V. K., & Morozov, V. G. (2003). Peptides of pineal gland and thymus prolong human life. *Neuro endocrinology letters*, *24*(3-4), 233–240.

[28] Khavinson, V. K., Lezhava, T. A., & Monaselidze, J. R. (2003). Peptide Epitalon activates chromatin at the gene level. *Neuro endocrinology letters*, *24*(5), 329–333.

[29] Falutz, J., Mamputu, J. C., Potvin, D., Moyle, G., Soulban, G., Loughrey, H., & Grinspoon, S. (2010). Effects of tesamorelin (TH9507), a growth hormone-releasing factor analog, in HIV-infected patients with excess abdominal fat: a pooled analysis of two multicenter, double-blind, placebo-controlled phase 3 trials with long-term open-label extension. *Journal of acquired immune deficiency syndromes (1999)*, *53*(3), 311–322.

Published by Peptide Education Hub (pepedhub.com) - The Most Comprehensive Evidence-Based Peptide Resource for Healthcare Providers

Dallas Regenerative LLC | DRS Peptides | drspeptides.com

This guide is intended for licensed healthcare providers only. Always consult current clinical guidelines and exercise professional judgment when implementing peptide therapy protocols.