

Tarea 06

Nombre: David Alejandro Díaz Pineda

Determine el orden de la mejor aproximación para las siguientes funciones, usando la Serie de Taylor y el Polinomio de Lagrange:

1.
$$f(x) = \frac{1}{25x^2 + 1}, x_0 = 0$$

```
In [1]: import numpy as np
import matplotlib.pyplot as plt
```

- Serie de Taylor:

$$f'(x) = -\frac{50x}{(25x^2+1)^2}$$

$$f''(x) = -\frac{50}{(25x^2+1)^2} + \frac{5000x^2}{(25x^2+1)^3}$$

$$f'''(x) = \frac{15000x}{(25x^2+1)^3} - \frac{750000x^3}{(25x^2+1)^4}$$

$$f^{(4)}(x) = \frac{15000}{(25x^2+1)^3} - \frac{4500000x^2}{(25x^2+1)^4} + \frac{150000000x^4}{(25x^2+1)^5}$$

Quedándonos como serie de Taylor de orden 4 con $x_0 = 0$

$$f(0) \approx 1 - 25x^2 + 625x^4$$

así, graficamos cada aproximación

$$T_0(x) = 1$$


```
In [2]: def f(x):
        return 1/(25*x**2 + 1)

def T0(x):
    return np.ones_like(x)

x = np.linspace(-1, 1, 400)

plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="T0(x) = 1", linestyle="--")
plt.title("Aproximación de Taylor - Orden 0")
plt.xlabel("x")
plt.ylabel("Valor")
```

```
plt.legend()
plt.grid(True)
plt.show()
```




$$T_2 = 1 - 25x^2$$

```
In [3]: def f(x):
        return 1/(25*x**2 + 1)

        def T2(x):
            return 1 - 25*x**2

        x = np.linspace(-1, 1, 400)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, T2(x), label="T2(x) = 1 - 25x²", linestyle="--")
        plt.title("Aproximación de Taylor - Orden 2")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.ylim(0,1)
        plt.legend()
        plt.grid(True)
        plt.show()
```




$$T_4 = 1 - 25x^2 + 625x^4$$

```
In [4]: def f(x):
        return 1/(25*x**2 + 1)

        def T4(x):
            return 1 - 25*x**2 + 625*x**4

        x = np.linspace(-1, 1, 400)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, T4(x), label="T4(x) = 1 - 25x^2 + 625x^4", linestyle="--")
        plt.title("Aproximación de Taylor - Orden 4")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.ylim(0,1)
        plt.legend()
        plt.grid(True)
        plt.show()
```



- Polinomio de LaGrange

Como en la serie de Taylor usamos un polinomio de orden 4, para tener un polinomio de orden 4 en el polinomio de LaGrange debemos usar 5 puntos:

$$(0, 1) \quad (0.2, 0.5) \quad (-0.2, 0.5) \quad (0.6, 0.1) \quad (-0.6, 0.1)$$

$$P(x) = L_0(x) + 0.5L_1(x) + 0.5L_2(x) + 0.1L_3(x) + 0.1L_4(x)$$

$$L_0 = \frac{(x-0.2)(x+0.2)(x-0.6)(x+0.6)}{0.0144}$$

$$L_1 = -\frac{x(x+0.2)(x-0.6)(x+0.6)}{0.0256}$$

$$L_2 = -\frac{x(x-0.2)(x-0.6)(x+0.6)}{0.0256}$$

$$L_3 = \frac{x(x-0.2)(x+0.2)(x+0.6)}{0.2304}$$

$$L_4 = \frac{x(x-0.2)(x+0.2)(x-0.6)}{0.2304}$$

$$P(x) = \frac{(x-0.2)(x+0.2)(x-0.6)(x+0.6)}{0.0144} - \frac{1}{2} \left(\frac{x(x+0.2)(x-0.6)(x+0.6)}{0.0256} \right) - \frac{1}{2} \left(\frac{x(x-0.2)(x-0.6)(x+0.6)}{0.0256} \right) +$$

$$P(x) = 31.25x^4 - 13.75x^2 + 1$$




$$P_0(x) = 1$$

```
In [5]: def f(x):
        return 1/(25*x**2 + 1)

        def T0(x):
            return np.ones_like(x)

        x = np.linspace(-1, 1, 400)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, T0(x), label="P0(x) = 1", linestyle="--")
        plt.title("Polinomio de Lagrange grado 0")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.legend()
        plt.grid(True)
        plt.show()
```




$$P_2(x) = 1 - 13.75x^2$$

```
In [6]: def f(x):
        return 1/(25*x**2 + 1)

        def T2(x):
            return 1 - 13.75*x**2

        x = np.linspace(-1, 1, 400)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, T2(x), label="P0(x) = 1 - 13.75x²", linestyle="--")
        plt.title("Polinomio de Lagrange grado 2")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.ylim (0,1)
        plt.legend()
        plt.grid(True)
        plt.show()
```




$$P_4(x) = 1 - 13.75x^2 + 31.25x^4$$

```
In [7]: def f(x):
        return 1/(25*x**2 + 1)

        def P(x):
            return 31.25*x**4 - 13.75*x**2 + 1

        x = np.linspace(-1, 1, 400)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, P(x), label="P(x) = 1 - 13.75x^2 + 31.25x^4", linestyle="--")
        plt.title("Polinomio de Interpolación de Lagrange")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.axhline(y=0, color='black')
        plt.ylim (-1,1)
        plt.legend()
        plt.grid(True)
        plt.show()
```



El orden de la mejor aproximación es de 4

2. $f(x) = \arctan x, x_0 = 1$

- Serie de Taylor

$$f'(x) = \frac{1}{1+x^2}$$

$$f''(x) = -\frac{2x}{(1+x^2)^2}$$

$$f'''(x) = -\frac{2}{(1+x^2)^2} + \frac{8x^2}{(1+x^2)^3}$$

$$f^{(4)}(x) = \frac{24x}{(1+x^2)^3} - \frac{48x^3}{(1+x^2)^4}$$

$$f^{(5)}(x) = \frac{24}{(1+x^2)^3} - \frac{288x^2}{(1+x^2)^4} + \frac{384x^4}{(1+x^2)^5}$$

Quedándonos como serie de Taylor con orden 5:

$$f(1) \approx \frac{\pi}{4} + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2 + \frac{1}{12}(x-1)^3 - \frac{1}{40}(x-1)^5$$

Así, graficamos cada aproximación


$$T_0(x) = \frac{\pi}{4}$$

```
In [8]: def f(x):
        return np.arctan(x)

        def T0(x):
            return np.ones_like(x) * np.pi/4

        x = np.linspace(-5, 5, 2000)

        plt.figure()
        plt.plot(x, f(x), label="f(x)")
        plt.plot(x, T0(x), label="T0(x) ", linestyle="--")
        plt.title("Aproximación de Taylor - Orden 0")
        plt.xlabel("x")
        plt.ylabel("Valor")
        plt.axhline(y=0, color='black')
        plt.ylim(-2, 2)
        plt.legend()
        plt.grid(True)
        plt.show()
```




$$T_1(x) = \frac{\pi}{4} + \frac{1}{2}(x - 1)$$

```
In [9]: def f(x):
        return np.arctan(x)

        def T0(x):
            return np.pi/4 + 1/2*(x-1)

        x = np.linspace(-5, 5, 2000)
```

```
plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="T1(x) ", linestyle="--")
plt.title("Aproximación de Taylor - Orden 1")
plt.xlabel("x")
plt.ylabel("Valor")
plt.ylim(-2, 2)
plt.axhline(y=0, color='black')
plt.legend()
plt.grid(True)
plt.show()
```



$$T_2(x) = \frac{\pi}{4} + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2$$


```
In [10]: def f(x):
          return np.arctan(x)

def T0(x):
    return np.pi/4 + 1/2*(x-1) - 1/4*(x-1)**2

x = np.linspace(-5, 5, 2000)

plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="T2(x)", linestyle="--")
plt.title("Aproximación de Taylor - Orden 2")
plt.xlabel("x")
plt.ylabel("Valor")
plt.ylim(-2, 2)
plt.axhline(y=0, color='black')
plt.legend()
```

```
plt.grid(True)
plt.show()
```




$$T_3(x) = \frac{\pi}{4} + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2 + \frac{1}{12}(x-1)^3$$

```
In [11]: def f(x):
          return np.arctan(x)

          def T0(x):
              return np.pi/4 + 1/2*(x-1) - 1/4*(x-1)**2 + 1/12*(x-1)**3

          x = np.linspace(-5, 5, 2000)

          plt.figure()
          plt.plot(x, f(x), label="f(x)")
          plt.plot(x, T0(x), label="T3(x)", linestyle="--")
          plt.title("Aproximación de Taylor - Orden 3")
          plt.xlabel("x")
          plt.ylabel("Valor")
          plt.ylim(-2, 2)
          plt.axhline(y=0, color='black')
          plt.legend()
          plt.grid(True)
          plt.show()
```




$$T_5(x) = \frac{\pi}{4} + \frac{1}{2}(x-1) - \frac{1}{4}(x-1)^2 + \frac{1}{12}(x-1)^3 - \frac{1}{40}(x-1)^5$$

```
In [12]: def f(x):
          return np.arctan(x)

          def T0(x):
              return np.pi/4 + 1/2*(x-1) - 1/4*(x-1)**2 + 1/12*(x-1)**3 - 1/40*(x-1)**5

          x = np.linspace(-5, 5, 2000)

          plt.figure()
          plt.plot(x, f(x), label="f(x)")
          plt.plot(x, T0(x), label="T5(x) ", linestyle="--")
          plt.title("Aproximación de Taylor - Orden 5")
          plt.xlabel("x")
          plt.ylabel("Valor")
          plt.ylim(-2, 2)
          plt.axhline(y=0, color='black')
          plt.legend()
          plt.grid(True)
          plt.show()
```



- Polinomio de LaGrange

Como en la serie de Taylor usamos un polinomio de orden 5, para tener un polinomio de orden 5 en el polinomio de LaGrange debemos usar 6 puntos:

$$\left(-1, -\frac{\pi}{4}\right) \quad (0, 0) \quad (0.5, 0.4636) \quad \left(1, \frac{\pi}{4}\right) \quad (1.5, 0.9828) \quad (2, 1.1071)$$

$$L_0(x) = -\frac{2}{45}x^5 + \frac{2}{9}x^4 - \frac{7}{18}x^3 + \frac{5}{18}x^2 - \frac{1}{15}x$$

$$L_1(x) = \frac{2}{3}x^5 - \frac{8}{3}x^4 + \frac{5}{2}x^3 + \frac{5}{3}x^2 - \frac{19}{6}x + 1$$

$$L_2(x) = -\frac{16}{9}x^5 + \frac{56}{9}x^4 - \frac{32}{9}x^3 - \frac{56}{9}x^2 + \frac{16}{3}x$$

$$L_3(x) = 2x^5 - 6x^4 + \frac{3}{2}x^3 + \frac{13}{2}x^2 - 3x$$

$$L_4(x) = -\frac{16}{15}x^5 + \frac{8}{3}x^4 - \frac{8}{3}x^2 + \frac{16}{15}x$$

$$L_2(x) = \frac{2}{9}x^5 - \frac{4}{9}x^4 - \frac{1}{18}x^3 + \frac{4}{9}x^2 - \frac{1}{16}x$$


$$P(x) = -0.0208397605651139x^5 + 0.126713715616397x^4 - 0.226503229907357x^3 -$$

$$P_1(x) = 1.03274115386992x$$

```
In [13]: def f(x):
          return np.arctan(x)

def T0(x):
    return 1.03274115386992 * x

x = np.linspace(-2, 3, 400)
x_points = np.array([-1, 0, 0.5, 1, 1.5, 2])
y_points = np.arctan(x_points)
plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="P1(x) ", linestyle="--")
plt.title("Polinomio de Lagrange grado 1")
plt.xlabel("x")
plt.ylabel("Valor")
plt.scatter(x_points, y_points, color='red', label='Puntos dados')
plt.legend()
plt.grid(True)
plt.show()
```



$$P_2(x) = -0.126713715616396x^2 + 1.03274115386992x$$

```
In [14]: def f(x):
          return np.arctan(x)


def T0(x):
    return 1.03274115386992 * x - 0.126713715616396 * x**2

x = np.linspace(-2, 3, 400)
```

```

x_points = np.array([-1, 0, 0.5, 1, 1.5, 2])
y_points = np.arctan(x_points)
plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="P2(x) ", linestyle="--")
plt.title("Polinomio de Lagrange grado 2")
plt.xlabel("x")
plt.ylabel("Valor")
plt.scatter(x_points, y_points, color='red', label='Puntos dados')
plt.legend()
plt.grid(True)
plt.show()

```



$$P_3(x) = -0.226503229907357x^3 - 0.126713715616396x^2 + 1.03274115386992x$$

```


In [15]: def f(x):
          return np.arctan(x)

def T0(x):
    return 1.03274115386992 * x - 0.126713715616396 * x**2 - 0.226503229907357 *

x = np.linspace(-2, 3, 400)
x_points = np.array([-1, 0, 0.5, 1, 1.5, 2])
y_points = np.arctan(x_points)
plt.figure()
plt.plot(x, f(x), label="f(x)")
plt.plot(x, T0(x), label="P3(x) ", linestyle="--")
plt.title("Polinomio de Lagrange grado 3")
plt.xlabel("x")
plt.ylabel("Valor")
plt.scatter(x_points, y_points, color='red', label='Puntos dados')
plt.legend()

```

```
plt.grid(True)
plt.show()
```




$$P_4(x) = 0.126713715616397x^4 - 0.226503229907357x^3 - 0.126713715616396x^2 + 1.0$$

```
In [16]: def f(x):
          return np.arctan(x)

          def T0(x):
              return 1.03274115386992 * x - 0.126713715616396 * x**2 - 0.226503229907357 *

          x = np.linspace(-2, 3, 400)
          x_points = np.array([-1, 0, 0.5, 1, 1.5, 2])
          y_points = np.arctan(x_points)
          plt.figure()
          plt.plot(x, f(x), label="f(x)")
          plt.plot(x, T0(x), label="P4(x) ", linestyle="--")
          plt.title("Polinomio de Lagrange grado 4")
          plt.xlabel("x")
          plt.ylabel("Valor")
          plt.scatter(x_points, y_points, color='red', label='Puntos dados')
          plt.legend()
          plt.grid(True)
          plt.show()
```




$$P_5(x) = -0.0208397605651139x^5 + 0.126713715616397x^4 - 0.226503229907357x^3 -$$

```
In [17]: def f(x):
          return np.arctan(x)

          def T0(x):
              return 1.03274115386992 * x - 0.126713715616396 * x**2 - 0.226503229907357 *

          x = np.linspace(-2, 3, 400)
          x_points = np.array([-1, 0, 0.5, 1, 1.5, 2])
          y_points = np.arctan(x_points)

          plt.figure()
          plt.plot(x, f(x), label="f(x)")
          plt.plot(x, T0(x), label="P5(x) ", linestyle="--")
          plt.title("Polinomio de Lagrange grado 5")
          plt.xlabel("x")
          plt.ylabel("Valor")
          plt.scatter(x_points, y_points, color='red', label='Puntos dados')
          plt.legend()
          plt.grid(True)
          plt.show()
```



El mejor orden de la aproximación de 5