



PROBLEM SET 2

- 1 How many three-digit numbers contain no zeros or nines as digits?
- 2 How many three-digit numbers have a digit sum of 10?
- 3 Let $p_n(k)$ denote the number of permutations of n objects with k fixed points. Show that

$$\sum_{k=0}^{n} k p_n(k) = n!$$

4 A permutation $(x_1, x_2, \dots x_{2n})$ of the set $\{1, 2, \dots 2n\}$, where n is a positive integer, is said to be *good* if

$$\exists i < n : |x_i - x_{i+1}| = n,$$

and is otherwise said to be bad. Show that, for any particular n, there are more good permutations than bad permutations.

5 At Mario's Magnificent Pizza, there are n customers and n tables, and any customer can sit at any table.

At Luigi's Luxurious Pizza, there are n customers and (2n-1) tables such that customer k can sit at tables $1, 2, 3, \ldots, (2k-1)$, but not at any other table.

Let M(n, r) denote the number of different ways in which r customers at Mario's can be seated at r tables, forming r customer-table pairs.

Similarly, let L(n, r) denote the number of different ways in which r customers at Luigi's can be seated at r tables, forming r customer-table pairs.

Prove that M(n,r) = L(n,r), for $r = 1, 2, \dots, n$.

- 6 Prove the following from first principles, without invoking known combinatorial identities:
- (a) Pascal's Rule: Not counting permutations, the number of ways to choose k objects from a set of n is equal to the number of ways to choose (k-1) objects from a set of (n-1), plus the number of ways to choose k objects from a set of (n-1).
- (b) Not counting permutations, the number of ways to choose an odd number of objects from a set is the same as the number of ways to choose an even number of objects from that set (where zero objects can be chosen).
- (c) Binomial Theorem: For any two real numbers x and y,

$$\sum_{k=0}^{n} \binom{n}{k} x^k y^{n-k} = (x+y)^k.$$





- (d) n Choose k Formula: Not counting permutations, there are $\frac{n!}{k!(n-k)!}$ ways to choose k objects from a set of n.
- (e) Hockey Stick Identity: Not counting permutations, the number of ways to choose (k+1) objects from a set of (n+1) is the sum of the number of ways to choose k objects from a set of 1, a set of 2, and so on up to n.
- (f) The number of ways to choose k objects from a set of n is $\frac{n}{k}$ times the number of ways to choose (k-1) objects from a set of (n-1), not counting permutations (so long as both quantities are well-defined).