

COM303: Digital Signal Processing

Lecture 4: Introduction to Fourier Analysis


Overview

- ▶ Fourier analysis: concept and motivation
- ▶ the complex exponential
- ▶ the Fourier basis
- ▶ the DFT

The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:


$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

signals are often expressed as a linear combination of “atomic” time units:

$$x[n] = \sum_{k=0}^{N-1} x[k] \delta[n - k]$$



The time domain

in vector notation:

$$\mathbf{x} = \sum_{k=0}^{N-1} x_k \boldsymbol{\delta}^{(k)}$$

where $\{\boldsymbol{\delta}^{(k)}\}$ is the canonical basis for \mathbb{C}^N ; e.g.:

$$\boldsymbol{\delta}^{(2)} = [0 \quad 0 \quad 1 \quad 0 \quad \dots \quad 0]^T$$

The frequency domain

Fourier analysis: express a signal as a combination of periodic oscillations:

$$\mathbf{x} = \sum_{k=0}^{N-1} X_k \mathbf{w}^{(k)}$$

where $\{\mathbf{w}^{(k)}\}$ is the Fourier basis.

Fourier transform: a change of basis in the space of discrete-time signals

The frequency domain



Fourier analysis: express a signal as a combination of periodic oscillations:

$$\mathbf{x} = \sum_{k=0}^{N-1} X_k \mathbf{w}^{(k)}$$

where $\{\mathbf{w}^{(k)}\}$ is the Fourier basis.

Fourier transform: a change of basis in the space of discrete-time signals

Oscillations are everywhere!



Oscillations are everywhere

- ▶ sustainable dynamic systems exhibit oscillatory behavior
- ▶ intuitively: things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...

Oscillations are everywhere

- ▶ sustainable dynamic systems exhibit oscillatory behavior
- ▶ intuitively: things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...

Oscillations are everywhere

- ▶ sustainable dynamic systems exhibit oscillatory behavior
- ▶ intuitively: things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...

Oscillations are everywhere

- ▶ sustainable dynamic systems exhibit oscillatory behavior
- ▶ intuitively: things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...


Oscillations are everywhere

- ▶ sustainable dynamic systems exhibit oscillatory behavior
- ▶ intuitively: things that don't move in circles can't last:
 - bombs
 - rockets
 - human beings...


Oscillations are everywhere



Oscillations are everywhere




Oscillations are everywhere



You too can detect sinusoids!


the human body has two receptors for sinusoidal signals:

cochlea (inner ear)



- ▶ air pressure sinusoids
- ▶ frequencies from 20Hz to 20KHz

rods and cones (retina)



- ▶ electromagnetic sinusoids
- ▶ frequencies from 430THz to 790THz

The intuition

- ▶ humans analyze complex signals (audio, images) in terms of their sinusoidal components
- ▶ we can build instruments that “resonate” at one or multiple frequencies (tuning fork vs piano)
- ▶ the “frequency domain” seems to be as important as the time domain

The intuition

- ▶ humans analyze complex signals (audio, images) in terms of their sinusoidal components
- ▶ we can build instruments that “resonate” at one or multiple frequencies (tuning fork vs piano)
- ▶ the “frequency domain” seems to be as important as the time domain

The intuition

- ▶ humans analyze complex signals (audio, images) in terms of their sinusoidal components
- ▶ we can build instruments that “resonate” at one or multiple frequencies (tuning fork vs piano)
- ▶ the “frequency domain” seems to be as important as the time domain

Fundamental question

can we decompose any signal into sinusoidal elements?

yes, and Fourier showed us how to do it *exactly*!

analysis

- ▶ from time domain to frequency domain
- ▶ find the contribution of different frequencies
- ▶ discover “hidden” signal properties

synthesis

- ▶ from frequency domain to time domain
- ▶ create signals with known frequency content
- ▶ fit signals to specific frequency regions

Fundamental question

can we decompose any signal into sinusoidal elements?

yes, and Fourier showed us how to do it *exactly*!

analysis

- ▶ from time domain to frequency domain
- ▶ find the contribution of different frequencies
- ▶ discover “hidden” signal properties

synthesis

- ▶ from frequency domain to time domain
- ▶ create signals with known frequency content
- ▶ fit signals to specific frequency regions

Fundamental question

can we decompose any signal into sinusoidal elements?

yes, and Fourier showed us how to do it *exactly*!

analysis

- ▶ from time domain to frequency domain
- ▶ find the contribution of different frequencies
- ▶ discover “hidden” signal properties

synthesis

- ▶ from frequency domain to time domain
- ▶ create signals with known frequency content
- ▶ fit signals to specific frequency regions

Fundamental question

can we decompose any signal into sinusoidal elements?

yes, and Fourier showed us how to do it *exactly*!

analysis

- ▶ from time domain to frequency domain
- ▶ find the contribution of different frequencies
- ▶ discover “hidden” signal properties

synthesis

- ▶ from frequency domain to time domain
- ▶ create signals with known frequency content
- ▶ fit signals to specific frequency regions

The mathematical setup

- ▶ let's start with finite-length signals (i.e. vectors in \mathbb{C}^N)
- ▶ Fourier analysis is a simple change of basis
- ▶ a change of basis is a change of perspective
- ▶ a change of perspective can reveal things (if the basis is good)

The mathematical setup

- ▶ let's start with finite-length signals (i.e. vectors in \mathbb{C}^N)
- ▶ Fourier analysis is a simple change of basis
- ▶ a change of basis is a change of perspective
- ▶ a change of perspective can reveal things (if the basis is good)


The mathematical setup

- ▶ let's start with finite-length signals (i.e. vectors in \mathbb{C}^N)
- ▶ Fourier analysis is a simple change of basis
- ▶ a change of basis is a change of perspective
- ▶ a change of perspective can reveal things (if the basis is good)


The mathematical setup

- ▶ let's start with finite-length signals (i.e. vectors in \mathbb{C}^N)
- ▶ Fourier analysis is a simple change of basis
- ▶ a change of basis is a change of perspective
- ▶ a change of perspective can reveal things (if the basis is good)

Mystery signal



Mystery signal in the Fourier basis




the complex exponential

Prerequisite Warning!

$$e^{j\alpha}$$


The complex exponential

$$e^{j\alpha} = \cos \alpha + j \sin \alpha$$




The complex exponential

z : point on the complex plane



The complex exponential

rotation: $z' = z e^{j\alpha}$



The discrete-time oscillatory heartbeat

Ingredients:

- ▶ a frequency ω (units: radians)
- ▶ an initial phase ϕ (units: radians)
- ▶ an amplitude A

$$\begin{aligned}x[n] &= Ae^{j(\omega n + \phi)} \\&= A[\cos(\omega n + \phi) + j \sin(\omega n + \phi)]\end{aligned}$$

The discrete-time oscillatory heartbeat

Ingredients:

- ▶ a frequency ω (units: radians)
- ▶ an initial phase ϕ (units: radians)
- ▶ an amplitude A

$$\begin{aligned}x[n] &= Ae^{j(\omega n + \phi)} \\&= A[\cos(\omega n + \phi) + j \sin(\omega n + \phi)]\end{aligned}$$

Why complex exponentials?

- ▶ we can use complex numbers in digital systems, so why not?
- ▶ it makes sense: every sinusoid can always be written as a sum of sine and cosine
- ▶ math is simpler: trigonometry becomes algebra

Why complex exponentials?

- ▶ we can use complex numbers in digital systems, so why not?
- ▶ it makes sense: every sinusoid can always be written as a sum of sine and cosine
- ▶ math is simpler: trigonometry becomes algebra

Why complex exponentials?

- ▶ we can use complex numbers in digital systems, so why not?
- ▶ it makes sense: every sinusoid can always be written as a sum of sine and cosine
- ▶ math is simpler: trigonometry becomes algebra

The advantages of complex exponentials

Example: change the phase of a cosine the “old-school” way

$$\cos(\omega n + \phi) = a \cos(\omega n) - b \sin(\omega n), \quad a = \cos \phi, \quad b = \sin \phi$$

- ▶ we have to remember complex trigonometric formulas
- ▶ we have to carry more terms in our equations

The advantages of complex exponentials

Example: change the phase of a cosine the “old-school” way

$$\cos(\omega n + \phi) = a \cos(\omega n) - b \sin(\omega n), \quad a = \cos \phi, \quad b = \sin \phi$$

- ▶ we have to remember complex trigonometric formulas
- ▶ we have to carry more terms in our equations

The advantages of complex exponentials

Example: change the phase of a cosine the “old-school” way

$$\cos(\omega n + \phi) = a \cos(\omega n) - b \sin(\omega n), \quad a = \cos \phi, \quad b = \sin \phi$$

- ▶ we have to remember complex trigonometric formulas
- ▶ we have to carry more terms in our equations

The advantages of complex exponentials

Example: change the phase of a pure cosine with complex exponentials

$$\cos(\omega n + \phi) = \operatorname{Re}\{e^{j(\omega n + \phi)}\} = \operatorname{Re}\{e^{j\omega n} e^{j\phi}\}$$

- ▶ sine and cosine “live” together
- ▶ phase shift is simple multiplication
- ▶ notation is simpler

The advantages of complex exponentials

Example: change the phase of a pure cosine with complex exponentials

$$\cos(\omega n + \phi) = \operatorname{Re}\{e^{j(\omega n + \phi)}\} = \operatorname{Re}\{e^{j\omega n} e^{j\phi}\}$$

- ▶ sine and cosine “live” together
- ▶ phase shift is simple multiplication
- ▶ notation is simpler

The advantages of complex exponentials

Example: change the phase of a pure cosine with complex exponentials

$$\cos(\omega n + \phi) = \operatorname{Re}\{e^{j(\omega n + \phi)}\} = \operatorname{Re}\{e^{j\omega n} e^{j\phi}\}$$

- ▶ sine and cosine “live” together
- ▶ phase shift is simple multiplication
- ▶ notation is simpler

The advantages of complex exponentials

Example: change the phase of a pure cosine with complex exponentials

$$\cos(\omega n + \phi) = \operatorname{Re}\{e^{j(\omega n + \phi)}\} = \operatorname{Re}\{e^{j\omega n} e^{j\phi}\}$$


- ▶ sine and cosine “live” together
- ▶ phase shift is simple multiplication
- ▶ notation is simpler

The discrete-time oscillatory heartbeat

$$x[n] = Ae^{j(\omega n + \phi)}$$


The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




The complex exponential generating machine

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Initial phase

$$x[n] = e^{j(\omega n + \phi)}; \quad x[n+1] = e^{j\omega} x[n], \quad x[0] = e^{j\phi}$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$




Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$



Careful: not every discrete-time sinusoid is periodic!

$$x[n] = e^{j\omega n}; \quad x[n+1] = e^{j\omega} x[n]$$



Condition for periodicity in discrete time

$$e^{j\omega n} \text{ periodic in } n \iff \omega = \frac{M}{N}2\pi, \quad M, N \in \mathbb{Z}$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$

$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$

$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$

$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$

$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$

$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$

Condition for periodicity in discrete time

$$x[n] = x[n + N]$$

$$e^{j(\omega n + \phi)} = e^{j(\omega(n+N) + \phi)}$$

$$e^{j\omega n} e^{j\phi} = e^{j\omega n} e^{j\omega N} e^{j\phi}$$

$$e^{j\omega N} = 1$$


$$\omega N = 2M\pi, \quad M \in \mathbb{Z}$$

$$\omega = \frac{M}{N} 2\pi$$


2π phase periodicity of complex exponentials

$$e^{j\alpha} = e^{j(\alpha+2k\pi)} \quad \forall k \in \mathbb{Z}$$


2π -periodicity: one point, many names




2π -periodicity: one point, many names




2π -periodicity: one point, many names



One point, many names



One point, many names




How “fast” can we go?


How “fast” can we go?

$$0 \leq \omega < 2\pi$$

Remember the complex exponential generating machine




Remember the complex exponential generating machine




How “fast” can we go?

$$\omega = 2\pi/12$$




How “fast” can we go?

$$\omega = 2\pi/6$$




How “fast” can we go?

$$\omega = 2\pi/5$$




How “fast” can we go?

$$\omega = 2\pi/4$$




How “fast” can we go?

$$\omega = 2\pi/3$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




How “fast” can we go?

$$\omega = 2\pi/2 = \pi$$




What if we go “faster”?

$$\pi < \omega < 2\pi$$




What if we go “faster”?

$$\pi < \omega < 2\pi$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$




Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$



Let's go really too fast

$$\omega = 2\pi - \alpha, \quad \alpha \text{ small}$$



The wagonwheel effect

Summary

- ▶ $x[n] = e^{j(\omega n + \phi)}$ is the prototypical DSP oscillation
- ▶ discrete-time oscillations are periodic ONLY if frequency a rational multiple of π
- ▶ in discrete time, ω and $\omega + 2k\pi$ are indistinguishable frequencies

Digital vs physical frequency

► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats

► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

Digital vs physical frequency

► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats

► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

Digital vs physical frequency

► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats

► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

Digital vs physical frequency

► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats

► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

Digital vs physical frequency

► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats

► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

Digital vs physical frequency


► Discrete time:

- n : no physical dimension (just a counter)
- periodicity: how many samples before pattern repeats


► “Real world”:

- periodicity: how many *seconds* before pattern repeats
- frequency measured in Hz (s^{-1})

How your PC plays sounds



How your PC plays sounds



Digital vs physical frequency

- ▶ set T_s , time in seconds between samples
- ▶ periodicity of M samples \longrightarrow periodicity of MT_s seconds
- ▶ real world frequency:

$$f = \frac{1}{MT_s}$$

Digital vs physical frequency

- ▶ set T_s , time in seconds between samples
- ▶ periodicity of M samples \longrightarrow periodicity of MT_s seconds
- ▶ real world frequency:

$$f = \frac{1}{MT_s}$$

Digital vs physical frequency

- ▶ set T_s , time in seconds between samples
- ▶ periodicity of M samples \longrightarrow periodicity of MT_s seconds
- ▶ real world frequency:

$$f = \frac{1}{MT_s}$$

the Fourier basis

The Fourier Basis for \mathbb{C}^N

Claim: the set of N signals in \mathbb{C}^N

$$w_k[n] = e^{j\frac{2\pi}{N}nk}, \quad n, k = 0, 1, \dots, N-1$$

is an orthogonal basis in \mathbb{C}^N .

The Fourier Basis for \mathbb{C}^N

In vector notation:


$$\{\mathbf{w}^{(k)}\}_{k=0,1,\dots,N-1}$$

with


$$w_n^{(k)} = e^{j\frac{2\pi}{N}nk}$$

is an orthogonal basis in \mathbb{C}^N


Recall the complex exponential generating machine...




Recall the complex exponential generating machine...




Recall the complex exponential generating machine...




Recall the complex exponential generating machine...




Recall the complex exponential generating machine...




Recall the complex exponential generating machine...




Basis vector $\mathbf{w}^{(0)} \in \mathbb{C}^{64}$




$$\omega_0 = 0$$




Basis vector $\mathbf{w}^{(1)} \in \mathbb{C}^{64}$




$$\omega_1 = \frac{2\pi}{64}$$




Basis vector $\mathbf{w}^{(2)} \in \mathbb{C}^{64}$




$$\omega_2 = 2\pi \frac{2}{64}$$




Basis vector $\mathbf{w}^{(3)} \in \mathbb{C}^{64}$




$$\omega_3 = 2\pi \frac{3}{64}$$




Basis vector $\mathbf{w}^{(4)} \in \mathbb{C}^{64}$




$$\omega_4 = 2\pi \frac{4}{64}$$




Basis vector $\mathbf{w}^{(5)} \in \mathbb{C}^{64}$




$$\omega_5 = 2\pi \frac{5}{64}$$




Basis vector $\mathbf{w}^{(16)} \in \mathbb{C}^{64}$




$$\omega_{16} = 2\pi \frac{16}{64}$$




Basis vector $\mathbf{w}^{(16)} \in \mathbb{C}^{64}$




$$\omega_{16} = 2\pi \frac{16}{64}$$




Basis vector $\mathbf{w}^{(20)} \in \mathbb{C}^{64}$




$$\omega_{20} = 2\pi \frac{20}{64}$$




Basis vector $\mathbf{w}^{(20)} \in \mathbb{C}^{64}$




$$\omega_{20} = 2\pi \frac{20}{64}$$




Basis vector $\mathbf{w}^{(30)} \in \mathbb{C}^{64}$




$$\omega_{30} = 2\pi \frac{30}{64}$$




Basis vector $\mathbf{w}^{(30)} \in \mathbb{C}^{64}$




$$\omega_{30} = 2\pi \frac{30}{64}$$




Basis vector $\mathbf{w}^{(31)} \in \mathbb{C}^{64}$




$$\omega_{31} = 2\pi \frac{31}{64}$$




Basis vector $\mathbf{w}^{(31)} \in \mathbb{C}^{64}$




$$\omega_{31} = 2\pi \frac{31}{64}$$




Basis vector $\mathbf{w}^{(32)} \in \mathbb{C}^{64}$




$$\omega_{32} = 2\pi \frac{32}{64}$$




Basis vector $\mathbf{w}^{(32)} \in \mathbb{C}^{64}$




$$\omega_{32} = 2\pi \frac{32}{64}$$




Basis vector $\mathbf{w}^{(33)} \in \mathbb{C}^{64}$




$$\omega_{33} = 2\pi \frac{33}{64}$$




Basis vector $\mathbf{w}^{(34)} \in \mathbb{C}^{64}$




$$\omega_{34} = 2\pi \frac{34}{64}$$




Basis vector $\mathbf{w}^{(60)} \in \mathbb{C}^{64}$




$$\omega_{60} = 2\pi \frac{60}{64}$$




Basis vector $\mathbf{w}^{(61)} \in \mathbb{C}^{64}$




$$\omega_{61} = 2\pi \frac{61}{64}$$




Basis vector $\mathbf{w}^{(62)} \in \mathbb{C}^{64}$




$$\omega_{62} = 2\pi \frac{62}{64}$$



Basis vector $\mathbf{w}^{(63)} \in \mathbb{C}^{64}$



$$\omega_{63} = 2\pi \frac{63}{64}$$



Proof of orthogonality

$$\begin{aligned}\langle \mathbf{w}^{(k)}, \mathbf{w}^{(h)} \rangle &= \sum_{n=0}^{N-1} (e^{j\frac{2\pi}{N}nk})^* e^{j\frac{2\pi}{N}nh} \\ &= \sum_{n=0}^{N-1} e^{j\frac{2\pi}{N}(h-k)n} \\ &= \begin{cases} N & \text{for } h = k \\ \frac{1 - e^{j2\pi(h-k)}}{1 - e^{j\frac{2\pi}{N}(h-k)}} = 0 & \text{otherwise} \end{cases}\end{aligned}$$

Proof of orthogonality

$$\begin{aligned}\langle \mathbf{w}^{(k)}, \mathbf{w}^{(h)} \rangle &= \sum_{n=0}^{N-1} (e^{j\frac{2\pi}{N}nk})^* e^{j\frac{2\pi}{N}nh} \\ &= \sum_{n=0}^{N-1} e^{j\frac{2\pi}{N}(h-k)n} \\ &= \begin{cases} N & \text{for } h = k \\ \frac{1 - e^{j2\pi(h-k)}}{1 - e^{j\frac{2\pi}{N}(h-k)}} = 0 & \text{otherwise} \end{cases}\end{aligned}$$

Proof of orthogonality

$$\begin{aligned}\langle \mathbf{w}^{(k)}, \mathbf{w}^{(h)} \rangle &= \sum_{n=0}^{N-1} (e^{j\frac{2\pi}{N}nk})^* e^{j\frac{2\pi}{N}nh} \\ &= \sum_{n=0}^{N-1} e^{j\frac{2\pi}{N}(h-k)n} \\ &= \begin{cases} N & \text{for } h = k \\ \frac{1 - e^{j2\pi(h-k)}}{1 - e^{j\frac{2\pi}{N}(h-k)}} = 0 & \text{otherwise} \end{cases}\end{aligned}$$

Proof of orthogonality

$$\begin{aligned}\langle \mathbf{w}^{(k)}, \mathbf{w}^{(h)} \rangle &= \sum_{n=0}^{N-1} (e^{j\frac{2\pi}{N}nk})^* e^{j\frac{2\pi}{N}nh} \\ &= \sum_{n=0}^{N-1} e^{j\frac{2\pi}{N}(h-k)n} \\ &= \begin{cases} N & \text{for } h = k \\ \frac{1 - e^{j2\pi(h-k)}}{1 - e^{j\frac{2\pi}{N}(h-k)}} = 0 & \text{otherwise} \end{cases}\end{aligned}$$

Remarks

- ▶ N orthogonal vectors \longrightarrow basis for \mathbb{C}^N
- ▶ vectors are not *orthonormal*. Normalization factor would be $1/\sqrt{N}$

Remarks

- ▶ N orthogonal vectors \longrightarrow basis for \mathbb{C}^N
- ▶ vectors are not *orthonormal*. Normalization factor would be $1/\sqrt{N}$

the Discrete Fourier Transform

The Fourier Basis for \mathbb{C}^N

- ▶ in “signal” notation: $w_k[n] = e^{j\frac{2\pi}{N}nk}$, $n, k = 0, 1, \dots, N-1$
- ▶ in vector notation: $\{\mathbf{w}^{(k)}\}_{k=0,1,\dots,N-1}$ with $w_n^{(k)} = e^{j\frac{2\pi}{N}nk}$

The Fourier Basis for \mathbb{C}^N

- ▶ in “signal” notation: $w_k[n] = e^{j\frac{2\pi}{N}nk}$, $n, k = 0, 1, \dots, N-1$
- ▶ in vector notation: $\{\mathbf{w}^{(k)}\}_{k=0,1,\dots,N-1}$ with $w_n^{(k)} = e^{j\frac{2\pi}{N}nk}$

The Fourier Basis for \mathbb{C}^N

- ▶ N orthogonal vectors \longrightarrow basis for \mathbb{C}^N
- ▶ vectors are not *orthonormal*. Normalization factor would be $1/\sqrt{N}$
- ▶ will keep normalization factor explicit in DFT formulas

Basis expansion

Analysis formula:

$$X_k = \langle \mathbf{w}^{(k)}, \mathbf{x} \rangle$$

Synthesis formula:

$$\mathbf{x} = \frac{1}{N} \sum_{k=0}^{N-1} X_k \mathbf{w}^{(k)}$$

Basis expansion (signal notation)

Analysis formula:

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$

N -point signal in the *frequency domain*

Synthesis formula:

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n = 0, 1, \dots, N-1$$

N -point signal in the *“time” domain*

Basis expansion (signal notation)

Analysis formula:

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$

N -point signal in the *frequency domain*

Synthesis formula:

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n = 0, 1, \dots, N-1$$

N -point signal in the *“time” domain*

Change of basis in matrix form

Define $W_N = e^{-j\frac{2\pi}{N}}$
(or simply W when N is evident from the context)

Change of basis matrix \mathbf{W} with $\mathbf{W}[n, m] = W_N^{nm}$:

$$\mathbf{W} = \begin{bmatrix} 1 & 1 & 1 & 1 & \dots & 1 \\ 1 & W^1 & W^2 & W^3 & \dots & W^{N-1} \\ 1 & W^2 & W^4 & W^6 & \dots & W^{2(N-1)} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & W^{N-1} & W^{2(N-1)} & W^{3(N-1)} & \dots & W^{(N-1)^2} \end{bmatrix}$$

Change of basis in matrix form

Define $W_N = e^{-j\frac{2\pi}{N}}$
(or simply W when N is evident from the context)

Change of basis matrix \mathbf{W} with $\mathbf{W}[n, m] = W_N^{nm}$:

$$\mathbf{W} = \begin{bmatrix} 1 & 1 & 1 & 1 & \dots & 1 \\ 1 & W^1 & W^2 & W^3 & \dots & W^{N-1} \\ 1 & W^2 & W^4 & W^6 & \dots & W^{2(N-1)} \\ & & & \dots & & \\ 1 & W^{N-1} & W^{2(N-1)} & W^{3(N-1)} & \dots & W^{(N-1)^2} \end{bmatrix}$$

Change of basis in matrix form

Analysis formula:

$$\mathbf{X} = \mathbf{W}\mathbf{x}$$

Synthesis formula:

$$\mathbf{x} = \frac{1}{N} \mathbf{W}^H \mathbf{X}$$

$$W_N^m = W_N^{(m \bmod N)}$$

e.g. $W_8^{11} = W_8^3$

$$W_N^m = W_N^{(m \bmod N)}$$

$$\text{e.g. } W_8^{11} = W_8^3$$

Small DFT matrices: $N = 2, 3$

$$W_2 = e^{-j\frac{2\pi}{2}} = -1$$

$$\mathbf{W}_2 = \begin{bmatrix} 1 & 1 \\ 1 & W \end{bmatrix} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$$

$$W_3 = e^{-j\frac{2\pi}{3}} = -(1 + j\sqrt{3})/2$$

$$\mathbf{W}_3 = \begin{bmatrix} 1 & 1 & 1 \\ 1 & W & W^2 \\ 1 & W^2 & W^4 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & W & W^2 \\ 1 & W^2 & W \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 \\ 1 & -(1 + j\sqrt{3})/2 & -(1 - j\sqrt{3})/2 \\ 1 & -(1 - j\sqrt{3})/2 & (1 - j\sqrt{3})/2 \end{bmatrix}$$

Small DFT matrices: $N = 4$

$$W_4 = e^{-j\frac{2\pi}{4}} = e^{-j\frac{\pi}{2}} = -j$$

$$\mathbf{W}_4 = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 \\ 1 & W^2 & W^4 & W^6 \\ 1 & W^3 & W^6 & W^9 \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 \\ 1 & W^2 & 1 & W^2 \\ 1 & W^3 & W^2 & W \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -j & -1 & j \\ 1 & -1 & 1 & -1 \\ 1 & j & -1 & -j \end{bmatrix}$$

Small DFT matrices: $N = 5$

$$\mathbf{W}_5 = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 & W^4 \\ 1 & W^2 & W^4 & W^6 & W^8 \\ 1 & W^3 & W^6 & W^9 & W^{12} \\ 1 & W^4 & W^8 & W^{12} & W^{16} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 & W^4 \\ 1 & W^2 & W^4 & W & W^3 \\ 1 & W^3 & W & W^4 & W^2 \\ 1 & W^4 & W^3 & W^2 & W \end{bmatrix}$$

Small DFT matrices: $N = 6$

$$\mathbf{W}_6 = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 & W^4 & W^5 \\ 1 & W^2 & W^4 & W^6 & W^8 & W^{10} \\ 1 & W^3 & W^6 & W^9 & W^{12} & W^{15} \\ 1 & W^4 & W^8 & W^{12} & W^{16} & W^{20} \\ 1 & W^5 & W^{10} & W^{15} & W^{20} & W^{25} \end{bmatrix} = \begin{bmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & W & W^2 & W^3 & W^4 & W^5 \\ 1 & W^2 & W^4 & 1 & W^2 & W^4 \\ 1 & W^3 & 1 & W^3 & 1 & W^3 \\ 1 & W^4 & W^2 & 1 & W^4 & W^2 \\ 1 & W^5 & W^4 & W^3 & W^2 & W \end{bmatrix}$$