

Solutions 5

1. a) This chain is clearly ergodic. The transition matrix is

$$\begin{pmatrix} 1-p & p & 0 \\ 1/2 & 0 & 1/2 \\ 0 & p & 1-p \end{pmatrix}$$

Assume that the detailed balance equation is satisfied. Then

$$\pi_1^*/2 = \pi_2^*p = \pi_0^*p$$

We conclude that

$$\pi_0^* = \pi_2^* = \frac{1}{2(1+p)} \quad \pi_1^* = \frac{p}{1+p}$$

It is then easy to verify that $\pi^* = \pi^*P$, and so this is indeed a stationary distribution, which obviously satisfies the detailed balance equation.

b) We know that $\lambda_0 = 1$, and so, to compute the eigenvalues, we must solve the equations

$$\begin{aligned} 2 - 2p &= 1 + \lambda_1 + \lambda_2 \\ -p(1-p) &= \lambda_1\lambda_2 \end{aligned}$$

Solving this, we obtain that $\lambda_1 = 1-p$ and $\lambda_2 = -p$. So $\lambda_* = \max(p, 1-p)$ and the spectral gap is given by $\gamma = 1 - \lambda_* = \min(p, 1-p)$.

c) For $p = \frac{1}{N}$, the spectral gap is $\gamma = \frac{1}{N}$. From the theorem seen in class, we know that $\|P_i^n - \pi\|_{\text{TV}} \leq \frac{\exp(-\gamma n)}{2\sqrt{\pi_i}}$, so here,

$$\max_{i \in S} \|P_i^n - \pi\|_{\text{TV}} \leq \frac{1}{2} \sqrt{\frac{1+1/N}{1/N}} \exp(-n/N) \leq \sqrt{N} \exp(-n/N) = \exp\left(\frac{\log N}{2} - \frac{n}{N}\right)$$

Taking therefore $n \geq N\left(\frac{\log N}{2} + c\right)$ with $c > 0$ sufficiently large (more precisely, $c = \log(1/\varepsilon)$) ensures that the maximum total variation norm is below ε .

d) For $p = 1 - \frac{1}{N}$, the spectral gap is again $\gamma = \frac{1}{N}$. So

$$\max_{i \in S} \|P_i^n - \pi\|_{\text{TV}} \leq \frac{1}{2} \sqrt{2(2 - 1/N)} \exp(-n/N) \leq \exp(-n/N)$$

Taking therefore $n \geq cN$ with $c = \log(1/\varepsilon)$ ensures that the maximum total variation norm is below ε , so

$$T_\varepsilon \leq N \log(1/\varepsilon)$$

2. a) The transition matrix being doubly stochastic, the stationary distribution is uniform (i.e. $\pi_i = \frac{1}{2N}$ for every $i \in S$) and satisfies the detailed balance equation.

b) Solving the equation $P\phi^{(1)} = \lambda\phi^{(1)}$, we obtain

$$\begin{aligned}\frac{N-1}{N}a + \frac{1}{N}b &= \lambda a \\ \frac{N-1}{N}a - \frac{1}{N}b &= \lambda b\end{aligned}$$

which is saying that λ is an eigenvalue of the 2×2 matrix

$$\begin{pmatrix} \frac{N-1}{N} & \frac{1}{N} \\ \frac{N-1}{N} & -\frac{1}{N} \end{pmatrix} = \begin{pmatrix} 1-\delta & \delta \\ 1-\delta & -\delta \end{pmatrix}$$

where we have set $\delta = \frac{1}{N}$. These eigenvalues are given by

$$\lambda_{\pm} = \frac{1 - 2\delta \pm \sqrt{(1 - 2\delta)^2 + 8\delta(1 - \delta)}}{2} = \frac{1 - 2\delta \pm \sqrt{1 + 4\delta - 4\delta^2}}{2}$$

For δ small (i.e. N large), the largest of these 2 eigenvalues is λ_+ , which is approximately given by

$$\lambda_+ \simeq \frac{1 - 2\delta + (1 + 2\delta - 4\delta^2)}{2} = 1 - 2\delta^2 = 1 - \frac{2}{N^2}$$

so the spectral gap $\gamma \simeq \frac{2}{N^2}$.

c) By the theorem seen in class,

$$\max_{i \in S} \|P_i^n - \pi\|_{\text{TV}} \leq \frac{\sqrt{2N}}{2} \exp(-\gamma n) \leq \sqrt{2} \exp\left(\frac{\log N}{2} - \frac{2n}{N^2}\right)$$

is below ε for $n \geq \frac{N^2}{2} \left(\frac{\log N}{2} + \log(\sqrt{2}/\varepsilon) \right)$, so

$$T_{\varepsilon} \leq \frac{N^2}{2} \left(\frac{\log N}{2} + \log(\sqrt{2}/\varepsilon) \right)$$