

COM303: Digital Signal Processing

Lecture 6: DFS and DTFT

Overview

- ▶ periodicity in the DFT
- ▶ the DFS
- ▶ the DTFT

Overview

- ▶ periodicity in the DFT
- ▶ the DFS
- ▶ the DTFT

Overview

- ▶ periodicity in the DFT
- ▶ the DFS
- ▶ the DTFT

DFT formulas

Analysis formula:

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$

N -point signal in the *frequency domain*

Synthesis formula:

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n = 0, 1, \dots, N-1$$

N -point signal in the *"time" domain*

DFT formulas

Analysis formula:

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$


N -point signal in the *frequency domain*

Synthesis formula:

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n = 0, 1, \dots, N-1$$

N -point signal in the *“time” domain*

DFT synthesis formula



Running the machine too long...

$$x[n + N] = x[n]$$

output signal is N -periodic!

Inherent periodicities in the DFT

the synthesis formula:

$$x[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n = 0, 1, \dots, N-1$$

produces an N -point signal in the time domain

the analysis formula:

$$X[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$

produces an N -point signal in the frequency domain

Inherent periodicities in the DFT

the synthesis formula:

$$\tilde{x}[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n \in \mathbb{Z}$$

produces an ***N*-periodic** signal in the time domain

the analysis formula:

$$\tilde{X}[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k = 0, 1, \dots, N-1$$

produces an *N*-point signal in the frequency domain

Inherent periodicities in the DFT

the synthesis formula:

$$\tilde{x}[n] = \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{j\frac{2\pi}{N}nk}, \quad n \in \mathbb{Z}$$

produces an ***N*-periodic** signal in the time domain

the analysis formula:

$$\tilde{X}[k] = \sum_{n=0}^{N-1} x[n] e^{-j\frac{2\pi}{N}nk}, \quad k \in \mathbb{Z}$$

produces an ***N*-periodic** signal in the frequency domain

Discrete Fourier Series (DFS)

DFS = DFT with periodicity explicit


- ▶ the DFS maps an N -periodic signal onto an N -periodic sequence of Fourier coefficients
- ▶ the inverse DFS maps an N -periodic sequence of Fourier coefficients a set onto an N -periodic signal
- ▶ the DFS of an N -periodic signal is mathematically equivalent to the DFT of one period

the Fourier transform for periodic signals


Periodic sequences: a bridge to infinite-length signals

- ▶ N -periodic sequence: N degrees of freedom
- ▶ DFS: only N Fourier coefficients capture all the information


Example: 32-tap sawtooth wave




Example: DFT of 32-tap sawtooth wave




What if we take the DFT of two periods?




Example: 64-point DFT of two periods




DFT of two periods: intuition




DFT of two periods: intuition




DFT of two periods: intuition



DFT of two periods: intuition



DFT of two periods: intuition



DFT of L periods

ingredients:

- ▶ finite-length signal $x[n]$, $n = 0, 1, \dots, N - 1$
- ▶ periodic signal: $\tilde{x}[n] = x[n \bmod N]$
- ▶ obviously $\tilde{x}[n] = \tilde{x}[n + pM]$ for all $p \in \mathbb{Z}$

DFT of L periods

$$\begin{aligned}X_L[k] &= \sum_{n=0}^{LM-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} \quad k = 0, 1, 2, \dots, LM - 1 \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n + pM] e^{-j \frac{2\pi}{LM} (n+pM)k} \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} e^{-j \frac{2\pi}{L} pk} \\&= \left(\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} \right) \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk}\end{aligned}$$

DFT of L periods

$$\begin{aligned}X_L[k] &= \sum_{n=0}^{LM-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} \quad k = 0, 1, 2, \dots, LM - 1 \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n + pM] e^{-j \frac{2\pi}{LM} (n+pM)k} \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} e^{-j \frac{2\pi}{L} pk} \\&= \left(\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} \right) \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk}\end{aligned}$$

DFT of L periods

$$\begin{aligned}X_L[k] &= \sum_{n=0}^{LM-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} \quad k = 0, 1, 2, \dots, LM - 1 \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n + pM] e^{-j \frac{2\pi}{LM} (n+pM)k} \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} e^{-j \frac{2\pi}{L} pk} \\&= \left(\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} \right) \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk}\end{aligned}$$

DFT of L periods

$$\begin{aligned}X_L[k] &= \sum_{n=0}^{LM-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} \quad k = 0, 1, 2, \dots, LM - 1 \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n + pM] e^{-j \frac{2\pi}{LM} (n+pM)k} \\&= \sum_{p=0}^{L-1} \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk} e^{-j \frac{2\pi}{L} pk} \\&= \left(\sum_{p=0}^{L-1} e^{-j \frac{2\pi}{L} pk} \right) \sum_{n=0}^{M-1} \tilde{x}[n] e^{-j \frac{2\pi}{LM} nk}\end{aligned}$$

We've seen this before

$$\sum_{p=0}^{L-1} e^{-j\frac{2\pi}{L}pk} = \begin{cases} L & \text{if } k \text{ multiple of } L \\ 0 & \text{otherwise} \end{cases}$$

(remember the orthogonality proof for the DFT basis)

DFT of L periods

if k is a multiple of L then k/L is an integer, so:

$$\sum_{n=0}^{M-1} x[n] e^{-j\frac{2\pi}{M}n\frac{k}{L}} = X[k/L]$$

DFT of L periods

$$X_L[k] = \begin{cases} L X[k/L] & \text{if } k = 0, L, 2L, 3L, \dots \\ 0 & \text{otherwise} \end{cases}$$

DFT and DFS

- ▶ again, all the spectral information for a periodic signal is contained in the DFT coefficients of a single period
- ▶ to stress the periodicity of the underlying signal, we use the term DFS

DFT and DFS

- ▶ again, all the spectral information for a periodic signal is contained in the DFT coefficients of a single period
- ▶ to stress the periodicity of the underlying signal, we use the term DFS

Finite-length time shifts revisited

The DFS helps us understand how to define time shifts for finite-length signals.

For an N -periodic sequence $\tilde{x}[n]$:

- ▶ $\tilde{x}[n - M]$ is well-defined for all $M \in \mathbb{N}$
- ▶ DFS $\{\tilde{x}[n - M]\} = e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k]$ (easy derivation)
- ▶ IDFS $\{\tilde{X}[k]\} = \tilde{x}[n - M]$

Finite-length time shifts revisited

The DFS helps us understand how to define time shifts for finite-length signals.

For an N -periodic sequence $\tilde{x}[n]$:

- ▶ $\tilde{x}[n - M]$ is well-defined for all $M \in \mathbb{N}$
- ▶ DFS $\{\tilde{x}[n - M]\} = e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k]$ (easy derivation)
- ▶ IDFS $\{\tilde{X}[k]\} = \tilde{x}[n - M]$

Finite-length time shifts revisited

The DFS helps us understand how to define time shifts for finite-length signals.

For an N -periodic sequence $\tilde{x}[n]$:

- ▶ $\tilde{x}[n - M]$ is well-defined for all $M \in \mathbb{N}$
- ▶ DFS $\{\tilde{x}[n - M]\} = e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k]$ (easy derivation)
- ▶ IDFS $\{\tilde{X}[k]\} = \tilde{x}[n - M]$

Finite-length time shifts revisited

The DFS helps us understand how to define time shifts for finite-length signals.

For an N -periodic sequence $\tilde{x}[n]$:

- ▶ $\tilde{x}[n - M]$ is well-defined for all $M \in \mathbb{N}$
- ▶ DFS $\{\tilde{x}[n - M]\} = e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k]$ (easy derivation)
- ▶ IDFS $\left\{ e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k] \right\} = \tilde{x}[n - M]$

Finite-length time shifts revisited

The DFS helps us understand how to define time shifts for finite-length signals.

For an N -periodic sequence $\tilde{x}[n]$:

- ▶ $\tilde{x}[n - M]$ is well-defined for all $M \in \mathbb{N}$
- ▶ DFS $\{\tilde{x}[n - M]\} = e^{-j\frac{2\pi}{N}Mk} \tilde{X}[k]$ (easy derivation)
- ▶ IDFS $\left\{ \boxed{e^{-j\frac{2\pi}{N}Mk}} \tilde{X}[k] \right\} = \tilde{x}[n - M]$



a delay in time becomes a *linear phase* factor in frequency

Finite-length time shifts revisited

For an N -point signal $x[n]$:

- ▶ $x[n - M]$ is *not* well-defined
- ▶ what is IDFT $\left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\}$?

Finite-length time shifts revisited

For an N -point signal $x[n]$:

- ▶ $x[n - M]$ is *not* well-defined
- ▶ what is IDFT $\left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\}$?

Finite-length time shifts revisited

$$\begin{aligned}\text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\ &= \frac{1}{N} \sum_{k=0}^{N-1} \left(\sum_{m=0}^{N-1} x[m] e^{-j\frac{2\pi}{N}mk} \right) e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k}\end{aligned}$$

Finite-length time shifts revisited

$$\begin{aligned}\text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\&= \frac{1}{N} \sum_{k=0}^{N-1} \left(\sum_{m=0}^{N-1} x[m] e^{-j\frac{2\pi}{N}mk} \right) e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\&= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k}\end{aligned}$$

Finite-length time shifts revisited

$$\begin{aligned}\text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{k=0}^{N-1} X[k] e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\ &= \frac{1}{N} \sum_{k=0}^{N-1} \left(\sum_{m=0}^{N-1} x[m] e^{-j\frac{2\pi}{N}mk} \right) e^{-j\frac{2\pi}{N}Mk} e^{j\frac{2\pi}{N}nk} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k}\end{aligned}$$

We've seen something like this before...

$$\sum_{n=0}^{N-1} e^{-j\frac{2\pi}{N}nk} = \begin{cases} N & \text{if } k \text{ multiple of } N \\ 0 & \text{otherwise} \end{cases}$$

(remember the orthogonality proof for the DFT basis)

Finite-length time shifts revisited

$$\sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} = \begin{cases} N & \text{for } (n - M - m) \text{ multiple of } N \\ 0 & \text{otherwise} \end{cases}$$

$$\forall L, N \in \mathbb{N}, \exists p \in \mathbb{N} : \quad L = pN + (L \bmod N)$$

Finite-length time shifts revisited

$$\sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} = \begin{cases} N & \text{for } (n - M - m) \text{ multiple of } N \\ 0 & \text{otherwise} \end{cases}$$

$$\forall L, N \in \mathbb{N}, \exists p \in \mathbb{N} : \quad L = pN + (L \bmod N)$$

Finite-length time shifts revisited

$$\sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} = \begin{cases} N & \text{for } m = (n - M) \bmod N \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} \text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} \\ &= x[(n - M) \bmod N] \end{aligned}$$

shifts for finite-length signals are “naturally” circular

Finite-length time shifts revisited

$$\sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} = \begin{cases} N & \text{for } m = (n - M) \bmod N \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} \text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} \\ &= x[(n - M) \bmod N] \end{aligned}$$

shifts for finite-length signals are “naturally” circular

Finite-length time shifts revisited

$$\sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} = \begin{cases} N & \text{for } m = (n - M) \bmod N \\ 0 & \text{otherwise} \end{cases}$$

$$\begin{aligned} \text{IDFT} \left\{ e^{-j\frac{2\pi}{N}Mk} X[k] \right\} [n] &= \frac{1}{N} \sum_{m=0}^{N-1} x[m] \sum_{k=0}^{N-1} e^{j\frac{2\pi}{N}(n-M-m)k} \\ &= x[(n - M) \bmod N] \end{aligned}$$

shifts for finite-length signals are “naturally” circular

The situation so far

Fourier representation for signal classes:

- ▶ N -point finite-length: DFT
- ▶ N -point periodic: DFS
- ▶ infinite length: ?

The situation so far

Fourier representation for signal classes:

- ▶ N -point finite-length: DFT
- ▶ N -point periodic: DFS
- ▶ infinite length: ?

The situation so far

Fourier representation for signal classes:

- ▶ N -point finite-length: DFT
- ▶ N -point periodic: DFS
- ▶ infinite length: ?

the Fourier transform for infinite-length signals

DFT of increasingly long signals

► Start with the DFT. What happens when $N \rightarrow \infty$?

► $\frac{2\pi}{N}k$ becomes denser in $[0, 2\pi]$...

► In the limit $\frac{2\pi}{N}k \rightarrow \omega$:

$$\sum_n x[n] e^{-j\omega n} \quad \omega \in \mathbb{R}$$

DFT of increasingly long signals

► Start with the DFT. What happens when $N \rightarrow \infty$?

► $\frac{2\pi}{N}k$ becomes denser in $[0, 2\pi]$...

► In the limit $\frac{2\pi}{N}k \rightarrow \omega$:

$$\sum_n x[n] e^{-j\omega n} \quad \omega \in \mathbb{R}$$

DFT of increasingly long signals

- ▶ Start with the DFT. What happens when $N \rightarrow \infty$?
- ▶ $\frac{2\pi}{N}k$ becomes denser in $[0, 2\pi]$...
- ▶ In the limit $\frac{2\pi}{N}k \rightarrow \omega$:

$$\sum_n x[n] e^{-j\omega n} \quad \omega \in \mathbb{R}$$

Discrete-Time Fourier Transform (DTFT)

Formal definition:

- ▶ $x[n] \in \ell_2(\mathbb{Z})$
- ▶ define the *function* of $\omega \in \mathbb{R}$

$$F(\omega) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ inversion (when $F(\omega)$ exists):

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\omega) e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

Discrete-Time Fourier Transform (DTFT)

Formal definition:

- ▶ $x[n] \in \ell_2(\mathbb{Z})$
- ▶ define the *function* of $\omega \in \mathbb{R}$

$$F(\omega) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ inversion (when $F(\omega)$ exists):

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\omega) e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

Discrete-Time Fourier Transform (DTFT)

Formal definition:

- ▶ $x[n] \in \ell_2(\mathbb{Z})$
- ▶ define the *function* of $\omega \in \mathbb{R}$

$$F(\omega) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ inversion (when $F(\omega)$ exists):

$$x[n] = \frac{1}{2\pi} \int_{-\pi}^{\pi} F(\omega) e^{j\omega n} d\omega, \quad n \in \mathbb{Z}$$

DTFT periodicity and notation

- ▶ $e^{j\omega n} = e^{j(\omega+2k\pi)n} \quad \forall k \in \mathbb{N}$

- ▶ $F(\omega)$ is 2π -periodic

- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$

DTFT periodicity and notation

- ▶ $e^{j\omega n} = e^{j(\omega+2k\pi)n} \quad \forall k \in \mathbb{N}$

- ▶ $F(\omega)$ is 2π -periodic

- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$

DTFT periodicity and notation

- ▶ $e^{j\omega n} = e^{j(\omega+2k\pi)n} \quad \forall k \in \mathbb{N}$
- ▶ $F(\omega)$ is 2π -periodic
- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$


DTFT periodicity and notation

- ▶ $e^{j\omega n} = e^{j(\omega+2k\pi)n} \quad \forall k \in \mathbb{N}$
- ▶ $F(\omega)$ is 2π -periodic
- ▶ to stress periodicity (and for other reasons) we will write

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ by convention, $X(e^{j\omega})$ is represented over $[-\pi, \pi]$

$$x[n] = \alpha^n u[n], \quad |\alpha| < 1$$



DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$

DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$

DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$


DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$\begin{aligned} X(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n \\ &= \frac{1}{1 - \alpha e^{-j\omega}} \end{aligned}$$


DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$

$$|X(e^{j\omega})|^2 = \frac{1}{1 + \alpha^2 - 2\alpha \cos \omega}$$


Plotting the DTFT




Plotting the DTFT




Plotting the DTFT




Plotting the DTFT




DTFT of $x[n] = \alpha^n u[n]$, $|\alpha| < 1$




Remember the periodicity!




Remember the periodicity!



Remember the periodicity!



Remember the periodicity!



DTFT intuition and properties

Overview:

- ▶ DTFT Existence
- ▶ Properties
- ▶ DTFT as basis expansion

Discrete-Time Fourier Transform

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ when does it exist?
- ▶ is it a change of basis?

Discrete-Time Fourier Transform

$$X(e^{j\omega}) = \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n}$$

- ▶ when does it exist?
- ▶ is it a change of basis?

Existence easy for absolutely summable sequences

$$\begin{aligned}|X(e^{j\omega})| &= \left| \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \right| \\ &\leq \sum_{n=-\infty}^{\infty} |x[n] e^{-j\omega n}| \\ &= \sum_{n=-\infty}^{\infty} |x[n]| \\ &< \infty\end{aligned}$$

Existence easy for absolutely summable sequences

$$\begin{aligned}|X(e^{j\omega})| &= \left| \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \right| \\ &\leq \sum_{n=-\infty}^{\infty} |x[n] e^{-j\omega n}| \\ &= \sum_{n=-\infty}^{\infty} |x[n]| \\ &< \infty\end{aligned}$$

Existence easy for absolutely summable sequences

$$\begin{aligned}|X(e^{j\omega})| &= \left| \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \right| \\ &\leq \sum_{n=-\infty}^{\infty} |x[n] e^{-j\omega n}| \\ &= \sum_{n=-\infty}^{\infty} |x[n]| \\ &< \infty\end{aligned}$$

Existence easy for absolutely summable sequences

$$\begin{aligned}|X(e^{j\omega})| &= \left| \sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} \right| \\ &\leq \sum_{n=-\infty}^{\infty} |x[n] e^{-j\omega n}| \\ &= \sum_{n=-\infty}^{\infty} |x[n]| \\ &< \infty\end{aligned}$$

Inversion easy for absolutely summable sequences

$$\begin{aligned}\frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) e^{j\omega n} d\omega &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{k=-\infty}^{\infty} x[k] e^{-j\omega k} \right) e^{j\omega n} d\omega \\ &= \sum_{k=-\infty}^{\infty} x[k] \int_{-\pi}^{\pi} \frac{e^{j\omega(n-k)}}{2\pi} d\omega \\ &= x[n]\end{aligned}$$

Inversion easy for absolutely summable sequences

$$\begin{aligned}\frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) e^{j\omega n} d\omega &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{k=-\infty}^{\infty} x[k] e^{-j\omega k} \right) e^{j\omega n} d\omega \\ &= \sum_{k=-\infty}^{\infty} x[k] \int_{-\pi}^{\pi} \frac{e^{j\omega(n-k)}}{2\pi} d\omega \\ &= x[n]\end{aligned}$$

Inversion easy for absolutely summable sequences

$$\begin{aligned}\frac{1}{2\pi} \int_{-\pi}^{\pi} X(e^{j\omega}) e^{j\omega n} d\omega &= \frac{1}{2\pi} \int_{-\pi}^{\pi} \left(\sum_{k=-\infty}^{\infty} x[k] e^{-j\omega k} \right) e^{j\omega n} d\omega \\ &= \sum_{k=-\infty}^{\infty} x[k] \int_{-\pi}^{\pi} \frac{e^{j\omega(n-k)}}{2\pi} d\omega \\ &= x[n]\end{aligned}$$

the DTFT as the limit of the DFS

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Synopsis

- ▶ $x[n]$ absolutely summable $\Rightarrow X(e^{j\omega})$ exists formally
- ▶ $x[n]$ absolutely summable \Rightarrow we can *periodize* it into $\tilde{x}_N[n]$
- ▶ natural Fourier representation for $\tilde{x}_N[n]$ is DFS
- ▶ DFS of $\tilde{x}_N[n]$ turns out to be $X(e^{j\omega})$ at $\omega = (2\pi/N)k$
- ▶ as N grows to infinity $\tilde{x}_N[n]$ becomes $x[n]$
- ▶ as N grows to infinity natural Fourier representation becomes $X(e^{j\omega})$

Some intuition

With $x[n]$ absolutely summable we can build arbitrarily “periodized” sequences:

$$\tilde{x}_N[n] = \sum_{p=-\infty}^{\infty} x[n + pN]$$

clearly $\tilde{x}_N[n] = \tilde{x}_N[n + N]$


Some intuition

With $x[n]$ absolutely summable we can build arbitrarily “periodized” sequences:


$$\tilde{x}_N[n] = \sum_{p=-\infty}^{\infty} x[n + pN]$$

clearly $\tilde{x}_N[n] = \tilde{x}_N[n + N]$


Periodization




Periodization




Periodization




Periodization




Periodization



Periodization




Periodization



Periodization


Let N grow large...



$N = 10$

Periodization


Let N grow large...



$N = 15$

Periodization


Let N grow large...



$N = 20$

Periodization


Let N grow large...



$N = 40$

Periodization

... as N grows, $\tilde{x}_N[n] \rightarrow x[n]$



$N = 80$

From DFS to DTFT

Natural spectral representation for $\tilde{x}_N[n]$ is the DFS:

$$\begin{aligned}\tilde{X}[k] &= \sum_{n=0}^{N-1} \tilde{x}_N[n] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{n=0}^{N-1} \sum_{p=-\infty}^{\infty} x[n + pN] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k}\end{aligned}$$

(remember $e^{j\alpha} = e^{j(\alpha+2K\pi)} \quad \forall K$)

From DFS to DTFT

Natural spectral representation for $\tilde{x}_N[n]$ is the DFS:

$$\begin{aligned}\tilde{X}[k] &= \sum_{n=0}^{N-1} \tilde{x}_N[n] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{n=0}^{N-1} \sum_{p=-\infty}^{\infty} x[n + pN] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k}\end{aligned}$$

(remember $e^{j\alpha} = e^{j(\alpha+2K\pi)} \quad \forall K$)

From DFS to DTFT

Natural spectral representation for $\tilde{x}_N[n]$ is the DFS:

$$\begin{aligned}\tilde{X}[k] &= \sum_{n=0}^{N-1} \tilde{x}_N[n] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{n=0}^{N-1} \sum_{p=-\infty}^{\infty} x[n + pN] e^{-j\frac{2\pi}{N}nk} \\ &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k}\end{aligned}$$

(remember $e^{j\alpha} = e^{j(\alpha+2K\pi)} \quad \forall K$)

From double sum to single sum

we can always write

$$\sum_{m=-\infty}^{\infty} y[m] = \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} y[n + pN]$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0				
	1				
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0			
	1				
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1		
	1				
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1	2	
	1				
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1	2	3
	1				
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1	2	3
	1	4			
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1	2	3
	1	4	5		
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		n			
		0	1	2	3
p					
	...				
	-1				
	0	0	1	2	3
	1	4	5	6	
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		<i>n</i>			
		0	1	2	3
<i>p</i>					
	...				
	-1				
	0	0	1	2	3
	1	4	5	6	7
	2				
	...				

$$m = n + 4p$$

Example (N=4)

		m			
		0	1	2	3
p					
	...				
	-1	-4	-3	-2	-1
	0	0	1	2	3
	1	4	5	6	7
	2	8	9	10	11
	...				

$$n = 4p + m$$

From DFS to DTFT

$$\begin{aligned}\tilde{X}[k] &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k} \\ &= \sum_{m=-\infty}^{\infty} x[m] e^{-j\frac{2\pi}{N}km} \\ &= X(e^{j\omega})|_{\omega=\frac{2\pi}{N}k}\end{aligned}$$

From DFS to DTFT

$$\begin{aligned}\tilde{X}[k] &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k} \\ &= \sum_{m=-\infty}^{\infty} x[m] e^{-j\frac{2\pi}{N}km} \\ &= X(e^{j\omega})|_{\omega=\frac{2\pi}{N}k}\end{aligned}$$

From DFS to DTFT

$$\begin{aligned}\tilde{X}[k] &= \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} x[n + pN] e^{-j\frac{2\pi}{N}(n+pN)k} \\ &= \sum_{m=-\infty}^{\infty} x[m] e^{-j\frac{2\pi}{N}km} \\ &= X(e^{j\omega})|_{\omega=\frac{2\pi}{N}k}\end{aligned}$$

From DFS to DTFT

- ▶ we're comfortable with DFS: change of basis, energy conservation, etc.
- ▶ as N grows, $\tilde{x}_N[n] \rightarrow x[n]$ and the spectral representation “becomes” the DTFT
- ▶ we can retain the “change of basis” paradigm for the DTFT


From DFS to DTFT

- ▶ we're comfortable with DFS: change of basis, energy conservation, etc.
- ▶ as N grows, $\tilde{x}_N[n] \rightarrow x[n]$ and the spectral representation “becomes” the DTFT
- ▶ we can retain the “change of basis” paradigm for the DTFT

From DFS to DTFT


- ▶ we're comfortable with DFS: change of basis, energy conservation, etc.
- ▶ as N grows, $\tilde{x}_N[n] \rightarrow x[n]$ and the spectral representation “becomes” the DTFT
- ▶ we can retain the “change of basis” paradigm for the DTFT

Review: DFT




Review: DFT


$$X[k] = \langle e^{j\frac{2\pi}{N}nk}, x[n] \rangle$$




Review: DFT



Review: DFT



Review: DFS



What about the DTFT?

- ▶ formally DTFT is an inner product in \mathbb{C}^∞ :

$$\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} = \langle e^{j\omega n}, x[n] \rangle$$

- ▶ “basis” is an infinite, uncountable basis: $\{e^{j\omega n}\}_{\omega \in \mathbb{R}}$
- ▶ something “breaks down”: we start with sequences but the transform is a function
- ▶ we used absolutely summable sequences but DTFT exists for all square-summable sequences (proof is rather technical)

What about the DTFT?

- ▶ formally DTFT is an inner product in \mathbb{C}^∞ :

$$\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} = \langle e^{j\omega n}, x[n] \rangle$$

- ▶ “basis” is an infinite, uncountable basis: $\{e^{j\omega n}\}_{\omega \in \mathbb{R}}$
- ▶ something “breaks down”: we start with sequences but the transform is a function
- ▶ we used absolutely summable sequences but DTFT exists for all square-summable sequences (proof is rather technical)

What about the DTFT?

- ▶ formally DTFT is an inner product in \mathbb{C}^∞ :

$$\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} = \langle e^{j\omega n}, x[n] \rangle$$


- ▶ “basis” is an infinite, uncountable basis: $\{e^{j\omega n}\}_{\omega \in \mathbb{R}}$
- ▶ something “breaks down”: we start with sequences but the transform is a function
- ▶ we used absolutely summable sequences but DTFT exists for all square-summable sequences (proof is rather technical)

What about the DTFT?

- ▶ formally DTFT is an inner product in \mathbb{C}^∞ :

$$\sum_{n=-\infty}^{\infty} x[n] e^{-j\omega n} = \langle e^{j\omega n}, x[n] \rangle$$

- ▶ “basis” is an infinite, uncountable basis: $\{e^{j\omega n}\}_{\omega \in \mathbb{R}}$
- ▶ something “breaks down”: we start with sequences but the transform is a function
- ▶ we used absolutely summable sequences but DTFT exists for all square-summable sequences (proof is rather technical)



DTFT properties

- ▶ linearity

$$\text{DTFT}\{\alpha x[n] + \beta y[n]\} = \alpha X(e^{j\omega}) + \beta Y(e^{j\omega})$$

- ▶ time shift

$$\text{DTFT}\{x[n - M]\} = e^{-j\omega M} X(e^{j\omega})$$

- ▶ modulation (dual)

$$\text{DTFT}\{e^{j\omega_0 n} x[n]\} = X(e^{j(\omega - \omega_0)})$$

DTFT properties

- ▶ linearity

$$\text{DTFT}\{\alpha x[n] + \beta y[n]\} = \alpha X(e^{j\omega}) + \beta Y(e^{j\omega})$$

- ▶ time shift

$$\text{DTFT}\{x[n - M]\} = e^{-j\omega M} X(e^{j\omega})$$

- ▶ modulation (dual)

$$\text{DTFT}\{e^{j\omega_0 n} x[n]\} = X(e^{j(\omega - \omega_0)})$$

DTFT properties

- ▶ linearity

$$\text{DTFT}\{\alpha x[n] + \beta y[n]\} = \alpha X(e^{j\omega}) + \beta Y(e^{j\omega})$$

- ▶ time shift

$$\text{DTFT}\{x[n - M]\} = e^{-j\omega M} X(e^{j\omega})$$

- ▶ modulation (dual)

$$\text{DTFT}\{e^{j\omega_0 n} x[n]\} = X(e^{j(\omega - \omega_0)})$$

DTFT properties

- ▶ time reversal

$$\text{DTFT}\{x[-n]\} = X(e^{-j\omega})$$

- ▶ conjugation

$$\text{DTFT}\{x^*[n]\} = X^*(e^{-j\omega})$$

DTFT properties

- ▶ time reversal

$$\text{DTFT}\{x[-n]\} = X(e^{-j\omega})$$

- ▶ conjugation

$$\text{DTFT}\{x^*[n]\} = X^*(e^{-j\omega})$$

Some particular cases:

- ▶ if $x[n]$ is symmetric, the DTFT is symmetric:

$$x[n] = x[-n] \iff X(e^{j\omega}) = X(e^{-j\omega})$$

- ▶ if $x[n]$ is real, the DTFT is Hermitian-symmetric:

$$x[n] = x^*[n] \iff X(e^{j\omega}) = X^*(e^{-j\omega})$$

- ▶ in other words: if $x[n]$ is real, the magnitude of the DTFT is symmetric:

$$x[n] \in \mathbb{R} \iff |X(e^{j\omega})| = |X(e^{-j\omega})|$$

- ▶ finally, if $x[n]$ is real and symmetric, the DTFT is also real and symmetric!

Some particular cases:

- ▶ if $x[n]$ is symmetric, the DTFT is symmetric:

$$x[n] = x[-n] \iff X(e^{j\omega}) = X(e^{-j\omega})$$

- ▶ if $x[n]$ is real, the DTFT is Hermitian-symmetric:

$$x[n] = x^*[n] \iff X(e^{j\omega}) = X^*(e^{-j\omega})$$

- ▶ in other words: if $x[n]$ is real, the magnitude of the DTFT is symmetric:

$$x[n] \in \mathbb{R} \iff |X(e^{j\omega})| = |X(e^{-j\omega})|$$

- ▶ finally, if $x[n]$ is real and symmetric, the DTFT is also real and symmetric!

Some particular cases:

- ▶ if $x[n]$ is symmetric, the DTFT is symmetric:

$$x[n] = x[-n] \iff X(e^{j\omega}) = X(e^{-j\omega})$$

- ▶ if $x[n]$ is real, the DTFT is Hermitian-symmetric:

$$x[n] = x^*[n] \iff X(e^{j\omega}) = X^*(e^{-j\omega})$$

- ▶ in other words: if $x[n]$ is real, the magnitude of the DTFT is symmetric:

$$x[n] \in \mathbb{R} \iff |X(e^{j\omega})| = |X(e^{-j\omega})|$$

- ▶ finally, if $x[n]$ is real and symmetric, the DTFT is also real and symmetric!

Some particular cases:

- ▶ if $x[n]$ is symmetric, the DTFT is symmetric:

$$x[n] = x[-n] \iff X(e^{j\omega}) = X(e^{-j\omega})$$

- ▶ if $x[n]$ is real, the DTFT is Hermitian-symmetric:

$$x[n] = x^*[n] \iff X(e^{j\omega}) = X^*(e^{-j\omega})$$

- ▶ in other words: if $x[n]$ is real, the magnitude of the DTFT is symmetric:

$$x[n] \in \mathbb{R} \iff |X(e^{j\omega})| = |X(e^{-j\omega})|$$

- ▶ finally, if $x[n]$ is real and symmetric, the DTFT is also real and symmetric!

DTFT as basis expansion

Some things are OK:

- ▶ DFT $\{\delta[n]\} = 1$
- ▶ DTFT $\{\delta[n]\} = \langle e^{j\omega n}, \delta[n] \rangle = 1$

DTFT as basis expansion

Some things are OK:

- ▶ DFT $\{\delta[n]\} = 1$
- ▶ DTFT $\{\delta[n]\} = \langle e^{j\omega n}, \delta[n] \rangle = 1$

DTFT as basis expansion

Some things aren't:

- ▶ $\text{DFT}\{1\} = \delta[n]$
- ▶ $\text{DTFT}\{1\} = \sum_{n=-\infty}^{\infty} e^{-j\omega n} = ?$
- ▶ problem: too many interesting sequences are *not* square summable!

DTFT as basis expansion

Some things aren't:

- ▶ DFT $\{1\} = \delta[n]$
- ▶ DTFT $\{1\} = \sum_{n=-\infty}^{\infty} e^{-j\omega n} = ?$
- ▶ problem: too many interesting sequences are *not* square summable!


DTFT as basis expansion

Some things aren't:

- ▶ DFT $\{1\} = \delta[n]$
- ▶ DTFT $\{1\} = \sum_{n=-\infty}^{\infty} e^{-j\omega n} = ?$
- ▶ problem: too many interesting sequences are *not* square summable!

The DTFT of the Karplus-Strong output

Karplus-Strong revisited



$$y[n] = \alpha y[n - M] + x[n]$$

Karplus-Strong revisited

- ▶ choose a signal $\bar{x}[n]$ that is nonzero only for $0 \leq n < M$
- ▶ generated signal is infinite-length but not periodic:

$$y[n] = \bar{x}[0], \bar{x}[1], \dots, \bar{x}[M-1], \alpha\bar{x}[0], \alpha\bar{x}[1], \dots, \alpha\bar{x}[M-1], \alpha^2\bar{x}[0], \alpha^2\bar{x}[1], \dots$$

- ▶ what is the DTFT of this signal?

Karplus-Strong revisited

- ▶ choose a signal $\bar{x}[n]$ that is nonzero only for $0 \leq n < M$
- ▶ generated signal is infinite-length but not periodic:

$$y[n] = \underbrace{\bar{x}[0], \bar{x}[1], \dots, \bar{x}[M-1]}_{1^{\text{st}} \text{ period}}, \underbrace{\alpha\bar{x}[0], \alpha\bar{x}[1], \dots, \alpha\bar{x}[M-1]}_{2^{\text{nd}} \text{ period}}, \underbrace{\alpha^2\bar{x}[0], \alpha^2\bar{x}[1], \dots}_{\dots}$$

- ▶ what is the DTFT of this signal?

Karplus-Strong revisited


- ▶ choose a signal $\bar{x}[n]$ that is nonzero only for $0 \leq n < M$
- ▶ generated signal is infinite-length but not periodic:

$$y[n] = \underbrace{\bar{x}[0], \bar{x}[1], \dots, \bar{x}[M-1]}_{1^{\text{st}} \text{ period}}, \underbrace{\alpha\bar{x}[0], \alpha\bar{x}[1], \dots, \alpha\bar{x}[M-1]}_{2^{\text{nd}} \text{ period}}, \underbrace{\alpha^2\bar{x}[0], \alpha^2\bar{x}[1], \dots}_{\dots}$$

- ▶ what is the DTFT of this signal?


KS revisited: 32-tap sawtooth wave

$$x[n] = 2n/(M - 1) - 1, \quad n = 0, 1, \dots, M - 1$$



KS revisited: decay $\alpha = 0.9$

$$y[n] = \alpha^{\lfloor n/M \rfloor} \bar{x}[n \bmod M] u[n]$$



DTFT of KS signal

$$Y(e^{j\omega}) = \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n}$$

Same trick we used before:

$$\sum_{m=-\infty}^{\infty} y[m] = \sum_{p=-\infty}^{\infty} \sum_{n=0}^{N-1} y[n + pN]$$

DTFT of KS signal

$$\begin{aligned} Y(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n} \\ &= \sum_{p=0}^{\infty} \sum_{m=0}^{M-1} \alpha^p \bar{x}[m] e^{-j\omega(pM+m)} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=0}^{M-1} \bar{x}[m] e^{-j\omega m} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=-\infty}^{\infty} \bar{x}[m] e^{-j\omega m} \\ &= A(e^{j\omega M}) \bar{X}(e^{j\omega}) \end{aligned}$$

DTFT of KS signal

$$\begin{aligned} Y(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n} \\ &= \sum_{p=0}^{\infty} \sum_{m=0}^{M-1} \alpha^p \bar{x}[m] e^{-j\omega(pM+m)} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=0}^{M-1} \bar{x}[m] e^{-j\omega m} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=-\infty}^{\infty} \bar{x}[m] e^{-j\omega m} \\ &= A(e^{j\omega M}) \bar{X}(e^{j\omega}) \end{aligned}$$

DTFT of KS signal

$$\begin{aligned} Y(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n} \\ &= \sum_{p=0}^{\infty} \sum_{m=0}^{M-1} \alpha^p \bar{x}[m] e^{-j\omega(pM+m)} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=0}^{M-1} \bar{x}[m] e^{-j\omega m} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=-\infty}^{\infty} \bar{x}[m] e^{-j\omega m} \\ &= A(e^{j\omega M}) \bar{X}(e^{j\omega}) \end{aligned}$$

DTFT of KS signal


$$\begin{aligned} Y(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n} \\ &= \sum_{p=0}^{\infty} \sum_{m=0}^{M-1} \alpha^p \bar{x}[m] e^{-j\omega(pM+m)} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=0}^{M-1} \bar{x}[m] e^{-j\omega m} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=-\infty}^{\infty} \bar{x}[m] e^{-j\omega m} \\ &= A(e^{j\omega M}) \bar{X}(e^{j\omega}) \end{aligned}$$

DTFT of KS signal

$$\begin{aligned} Y(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} y[n] e^{-j\omega n} \\ &= \sum_{p=0}^{\infty} \sum_{m=0}^{M-1} \alpha^p \bar{x}[m] e^{-j\omega(pM+m)} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=0}^{M-1} \bar{x}[m] e^{-j\omega m} \\ &= \sum_{p=0}^{\infty} \alpha^p e^{-j\omega Mp} \sum_{m=-\infty}^{\infty} \bar{x}[m] e^{-j\omega m} \\ &= A(e^{j\omega M}) \bar{X}(e^{j\omega}) \end{aligned}$$


We know the first term

$$A(e^{j\omega}) = \text{DTFT} \{ \alpha^n u[n] \} = \frac{1}{1 - \alpha e^{-j\omega}}$$




We know the first term

$A(e^{j\omega M})$ rescales the frequency axis: **periodicity!**




We know the first term

$A(e^{j\omega M})$ rescales the frequency axis: **periodicity!**




We know the first term

$A(e^{j\omega M})$ rescales the frequency axis: **periodicity!**




We know the first term

$A(e^{j\omega M})$ rescales the frequency axis: **periodicity!**



Second term is left as an exercise

$$\bar{X}(e^{j\omega}) = e^{-j\omega} \left(\frac{M+1}{M-1} \right) \frac{1 - e^{-j(M-1)\omega}}{(1 - e^{-j\omega})^2} - \frac{1 - e^{-j(M+1)\omega}}{(1 - e^{-j\omega})^2}$$



DTFT of KS with decay

$$Y(e^{j\omega}) = A(e^{j\omega M})\bar{X}(e^{j\omega})$$

