

---

## Review Session

---

### Problem 1: Review of Random Variables

Let  $X$  and  $Y$  be discrete random variables defined on some probability space with a joint pmf  $p_{XY}(x, y)$ . Let  $a, b \in \mathbb{R}$  be fixed.

(a) Prove that  $\mathbb{E}[aX + bY] = a\mathbb{E}[X] + b\mathbb{E}[Y]$ . Do not assume independence.

(b) Prove that if  $X$  and  $Y$  are independent random variables, then  $\mathbb{E}[X \cdot Y] = \mathbb{E}[X] \cdot \mathbb{E}[Y]$ .

(c) Assume that  $X$  and  $Y$  are not independent. Find an example where  $\mathbb{E}[X \cdot Y] \neq \mathbb{E}[X] \cdot \mathbb{E}[Y]$ , and another example where  $\mathbb{E}[X \cdot Y] = \mathbb{E}[X] \cdot \mathbb{E}[Y]$ .

(d) Prove that if  $X$  and  $Y$  are independent, then they are also uncorrelated, i.e.,

$$\text{Cov}(X, Y) := \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])] = 0. \quad (1)$$

(e) Find an example where  $X$  and  $Y$  are uncorrelated but dependent.

(f) Assume that  $X$  and  $Y$  are uncorrelated and let  $\sigma_X^2$  and  $\sigma_Y^2$  be the variances of  $X$  and  $Y$ , respectively. Find the variance of  $aX + bY$  and express it in terms of  $\sigma_X^2, \sigma_Y^2, a, b$ .

**Hint:** First show that  $\text{Cov}(X, Y) = \mathbb{E}[X \cdot Y] - \mathbb{E}[X] \cdot \mathbb{E}[Y]$ .

### Problem 2: Review of Gaussian Random Variables

A random variable  $X$  with probability density function

$$p_X(x) = \frac{1}{\sqrt{2\pi}\sigma^2} e^{-\frac{(x-m)^2}{2\sigma^2}} \quad (2)$$

is called a *Gaussian* random variable.

(a) Explicitly calculate the mean  $\mathbb{E}[X]$ , the second moment  $\mathbb{E}[X^2]$ , and the variance  $\text{Var}[X]$  of the random variable  $X$ .

(b) Let us now consider events of the following kind:

$$\mathbb{P}(X < \alpha). \quad (3)$$

Unfortunately for Gaussian random variables this cannot be calculated in closed form. Instead, we will rewrite it in terms of the standard Q-function:

$$Q(x) = \int_x^\infty \frac{1}{\sqrt{2\pi}} e^{-\frac{u^2}{2}} du \quad (4)$$

Express  $\mathbb{P}(X < \alpha)$  in terms of the Q-function and the parameters  $m$  and  $\sigma^2$  of the Gaussian pdf.

Like we said, the Q-function cannot be calculated in closed form. Therefore, it is important to have *bounds* on the Q-function. In the next 3 subproblems, you derive the most important of these bounds, learning some very general and powerful tools along the way:

(c) Derive the Markov inequality, which says that for any non-negative random variable  $X$  and positive  $a$ , we have

$$\mathbb{P}(X \geq a) \leq \frac{\mathbb{E}[X]}{a}. \quad (5)$$

(d) Use the Markov inequality to derive the Chernoff bound: the probability that a real random variable  $Z$  exceeds  $b$  is given by

$$\mathbb{P}(Z \geq b) \leq \mathbb{E}[e^{s(Z-b)}], \quad s \geq 0. \quad (6)$$

(e) Use the Chernoff bound to show that

$$Q(x) \leq e^{-\frac{x^2}{2}} \quad \text{for } x \geq 0. \quad (7)$$

### Problem 3: Moment Generating Function

Let  $X$  be a real-valued random variable taking values on a finite set. The logarithmic moment generating function is defined as follows.

$$\phi(s) := \ln E[\exp(sX)] = \ln \sum_x p(x) \exp(sx)$$

(a) Show that  $p_s(x) := p(x) \exp(sx) \exp(-\phi(s))$  is a probability mass function.

(b) Let  $X_s$  be a random variable taking the same value as  $X$  but with probabilities  $p_s(x)$ , show that  $\phi'(s) = E[X_s]$ .

(c) Show that

$$\phi''(s) = \text{Var}(X_s) := E[X_s^2] - E[X_s]^2$$

and conclude that  $\phi''(s) \geq 0$  and the inequality is strict except when  $X$  is deterministic.

(d) Let  $x_{\min} := \min\{x : p(x) > 0\}$  and  $x_{\max} := \max\{x : p(x) > 0\}$  be the smallest and largest values  $X$  takes. Show that

$$\lim_{s \rightarrow -\infty} \phi'(s) = x_{\min}, \quad \text{and} \quad \lim_{s \rightarrow \infty} \phi'(s) = x_{\max}.$$