

## COM303: Digital Signal Processing

### Lecture 18: Multirate signal processing

- ▶ ideal and practical sampling and interpolation
- ▶ bandpass sampling
- ▶ multirate signal processing

advanced topics in sampling

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

ideally

in practice

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

ideally

$$x[n] = \langle x(t), \text{sinc}\left(\frac{t - nT_s}{T_s}\right) \rangle$$

in practice

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$


ideally

$$x[n] = \langle x(t), \text{sinc} \left( \frac{t - nT_s}{T_s} \right) \rangle$$

$$X(e^{j\omega}) = X \left( j\Omega_N \frac{\omega \bmod 2\pi}{\pi} \right)$$


in practice

## Ideal case







## Ideal case




## Ideal case




## Ideal case



## Ideal case



## Ideal case



# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

ideally

$$x[n] = \langle x(t), \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right) \rangle$$

$$X(e^{j\omega}) = X\left(j\Omega_N \frac{\omega \bmod 2\pi}{\pi}\right)$$

in practice

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

ideally

$$x[n] = \langle x(t), \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right) \rangle$$

$$X(e^{j\omega}) = X\left(j\Omega_N \frac{\omega \bmod 2\pi}{\pi}\right)$$

in practice

$$x[n] = x(nT_s)$$

# From Continuous to Discrete Time

$$x(t) \longrightarrow x[n]$$

ideally

$$x[n] = \langle x(t), \text{sinc}\left(\frac{t - nT_s}{T_s}\right) \rangle$$

$$X(e^{j\omega}) = X\left(j\Omega_N \frac{\omega \bmod 2\pi}{\pi}\right)$$


in practice

$$x[n] = x(nT_s)$$


$$X(e^{j\omega}) = \frac{\Omega_N}{\pi} \sum_{k=-\infty}^{\infty} X_c\left(j\Omega_N \frac{\omega}{\pi} - 2jk\Omega_N\right)$$




## In practice




## In practice



## In practice



## In practice



# From Discrete to Continuous Time

$$x[n] \longrightarrow x(t)$$

# From Discrete to Continuous Time

$$x[n] \longrightarrow x(t)$$

ideally

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right)$$

# From Discrete to Continuous Time


$$x[n] \longrightarrow x(t)$$

ideally

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right)$$


$$X(j\Omega) = \frac{\pi}{\Omega_N} X(e^{j\pi\Omega/\Omega_N}) \operatorname{rect}\left(\frac{\Omega}{2\Omega_N}\right)$$

## sinc interpolation







## sinc interpolation



## sinc interpolation



## sinc interpolation



# From Discrete to Continuous Time

$$x[n] \longrightarrow x(t)$$

ideally

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right)$$

$$X(j\Omega) = \frac{\pi}{\Omega_N} X(e^{j\pi\Omega/\Omega_N}) \operatorname{rect}\left(\frac{\Omega}{2\Omega_N}\right)$$

in practice

# From Discrete to Continuous Time

$$x[n] \longrightarrow x(t)$$

ideally

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right)$$

$$X(j\Omega) = \frac{\pi}{\Omega_N} X(e^{j\pi\Omega/\Omega_N}) \operatorname{rect}\left(\frac{\Omega}{2\Omega_N}\right)$$

in practice

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] i\left(\frac{t - nT_s}{T_s}\right)$$

# From Discrete to Continuous Time

$$x[n] \longrightarrow x(t)$$

ideally

$$x(t) = \sum_{n=-\infty}^{\infty} x[n] \operatorname{sinc}\left(\frac{t - nT_s}{T_s}\right)$$

$$X(j\Omega) = \frac{\pi}{\Omega_N} X(e^{j\pi\Omega/\Omega_N}) \operatorname{rect}\left(\frac{\Omega}{2\Omega_N}\right)$$

in practice


$$x(t) = \sum_{n=-\infty}^{\infty} x[n] i\left(\frac{t - nT_s}{T_s}\right)$$

$$X(j\Omega) = ?$$

## Practical interpolation


$$x(t) = \sum_{n=-\infty}^{\infty} x[n] i\left(\frac{t - nT_s}{T_s}\right)$$

## Local interpolators






## Local interpolators



## Local interpolators



## Spectral representation (I)

$$\begin{aligned}X(j\Omega) &= \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt \\&= \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n] i\left(\frac{t - nT_s}{T_s}\right) e^{-j\Omega t} dt \\&= \sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} i\left(\frac{t - nT_s}{T_s}\right) e^{-j\Omega t} dt \\&= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega}\end{aligned}$$

## Spectral representation (I)

$$\begin{aligned}X(j\Omega) &= \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt \\&= \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n] i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= \sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega}\end{aligned}$$

## Spectral representation (I)

$$\begin{aligned}X(j\Omega) &= \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt \\&= \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n] i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= \sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega}\end{aligned}$$

## Spectral representation (I)

$$\begin{aligned}X(j\Omega) &= \int_{-\infty}^{\infty} x(t) e^{-j\Omega t} dt \\&= \int_{-\infty}^{\infty} \sum_{n=-\infty}^{\infty} x[n] i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= \sum_{n=-\infty}^{\infty} x[n] \int_{-\infty}^{\infty} i \left( \frac{t - nT_s}{T_s} \right) e^{-j\Omega t} dt \\&= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega}\end{aligned}$$

## Spectral representation (II)

$$\begin{aligned}X(j\Omega) &= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) \sum_{n=-\infty}^{\infty} x[n] e^{-j\pi \frac{\Omega}{\Omega_N} n} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) X(e^{j\pi \frac{\Omega}{\Omega_N}})\end{aligned}$$

## Spectral representation (II)

$$\begin{aligned}X(j\Omega) &= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) \sum_{n=-\infty}^{\infty} x[n] e^{-j\pi \frac{\Omega}{\Omega_N} n} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) X(e^{j\pi \frac{\Omega}{\Omega_N}})\end{aligned}$$



## Spectral representation (II)

$$\begin{aligned}X(j\Omega) &= T_s \sum_{n=-\infty}^{\infty} x[n] I(jT_s\Omega) e^{-jnT_s\Omega} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) \sum_{n=-\infty}^{\infty} x[n] e^{-j\pi \frac{\Omega}{\Omega_N} n} \\&= \frac{\pi}{\Omega_N} I\left(j\pi \frac{\Omega}{\Omega_N}\right) X(e^{j\pi \frac{\Omega}{\Omega_N}})\end{aligned}$$


## sinc interpolation

$$i(t) = \text{sinc}(t)$$


$$I(j\Omega) = \text{rect}\left(\frac{\Omega}{2\pi}\right)$$

$$\begin{aligned} X(j\Omega) &= \frac{\pi}{\Omega_N} I(j\pi\Omega/\Omega_N) X(e^{j\pi\Omega/\Omega_N}) \\ &= \frac{\pi}{\Omega_N} \text{rect}\left(\frac{\Omega}{2\Omega_N}\right) X(e^{j\pi\Omega/\Omega_N}) \end{aligned}$$


## sinc interpolation




## sinc interpolation




## sinc interpolation



## sinc interpolation



## sinc interpolation



## zero-order hold


$$i(t) = \text{rect}(t)$$

$$I(j\Omega) = \text{sinc}\left(\frac{\Omega}{2\pi}\right)$$


$$\begin{aligned} X(j\Omega) &= \frac{\pi}{\Omega_N} I(j\pi\Omega/\Omega_N) X(e^{j\pi\Omega/\Omega_N}) \\ &= \frac{\pi}{\Omega_N} \text{sinc}\left(\frac{\Omega}{2\Omega_N}\right) X(e^{j\pi\Omega/\Omega_N}) \end{aligned}$$




## zero-order hold




## Zero-order hold




## Zero-order hold





## Zero-order hold




## Zero-order hold




## first-order interpolator



## first-order interpolator




## first-order interpolator






## first-order interpolator



## first-order interpolator



## Bandpass Sampling


## sampling theorem gives a *sufficient* condition

- ▶ in theory,  $\Omega_N > \Omega_{\max}$
- ▶ what if signal is bandpass?


## sampling theorem gives a *sufficient* condition

- ▶ in theory,  $\Omega_N > \Omega_{\max}$
- ▶ what if signal is bandpass?


## Bandpass sampling



## Bandpass sampling




# Bandpass sampling







# Bandpass sampling




# Bandpass sampling



# Bandpass sampling



# Bandpass sampling



## Bandpass sampling conditions

- ▶ bandpass signal:  $X(j\Omega) = 0$  for  $|\Omega - \Omega_c| > \Omega_0$
- ▶ no alias requires at least:  $\Omega_N \geq \Omega_0$
- ▶ baseband condition:  $\Omega_N = \Omega_c/k$  for some  $k \in \mathbb{N}$
- ▶ for  $T_s = \pi/\Omega_N$ ,  $x[n] = x(nT_s)$  is an alias-free, baseband DT version of  $x(t)$

## Bandpass sampling conditions

- ▶ bandpass signal:  $X(j\Omega) = 0$  for  $|\Omega - \Omega_c| > \Omega_0$
- ▶ no alias requires at least:  $\Omega_N \geq \Omega_0$
- ▶ baseband condition:  $\Omega_N = \Omega_c/k$  for some  $k \in \mathbb{N}$
- ▶ for  $T_s = \pi/\Omega_N$ ,  $x[n] = x(nT_s)$  is an alias-free, baseband DT version of  $x(t)$

## Bandpass sampling conditions

- ▶ bandpass signal:  $X(j\Omega) = 0$  for  $|\Omega - \Omega_c| > \Omega_0$
- ▶ no alias requires at least:  $\Omega_N \geq \Omega_0$
- ▶ baseband condition:  $\Omega_N = \Omega_c/k$  for some  $k \in \mathbb{N}$
- ▶ for  $T_s = \pi/\Omega_N$ ,  $x[n] = x(nT_s)$  is an alias-free, baseband DT version of  $x(t)$

## Bandpass sampling conditions

- ▶ bandpass signal:  $X(j\Omega) = 0$  for  $|\Omega - \Omega_c| > \Omega_0$
- ▶ no alias requires at least:  $\Omega_N \geq \Omega_0$
- ▶ baseband condition:  $\Omega_N = \Omega_c/k$  for some  $k \in \mathbb{N}$
- ▶ for  $T_s = \pi/\Omega_N$ ,  $x[n] = x(nT_s)$  is an alias-free, baseband DT version of  $x(t)$



## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$

## Example: AM Channel

- ▶ AM radio band: 500kHz to 1.6MHz
- ▶ channel width is 9kHz, i.e.  $f_0 = 4.5\text{KHz}$
- ▶ take a channel at  $f_c = 1.5\text{MHz}$
- ▶ in theory:  $F_s \geq 2 * 1,504,500\text{Hz}$ ,  $T_s < 10^{-6}$  seconds!
- ▶ antialias:  $F_s \geq 2f_0 \Rightarrow F_s \geq 9\text{KHz}$
- ▶ baseband:  $kF_s/2 = 1500\text{kHz}$
- ▶ pick  $k = 300$
- ▶  $F_s = 10\text{KHz}$ ,  $T_s = 10^{-4}$



multirate signal processing

# Overview

- ▶ why multirate?
- ▶ upsampling
- ▶ downsampling
- ▶ applications

# Overview

- ▶ why multirate?
- ▶ upsampling
- ▶ downsampling
- ▶ applications

# Overview


- ▶ why multirate?
- ▶ upsampling
- ▶ downsampling
- ▶ applications

# Overview


- ▶ why multirate?
- ▶ upsampling
- ▶ downsampling
- ▶ applications

# Upsampling


$$x_{NU}[n] = \begin{cases} x[k] & \text{for } n = kN, \quad k \in \mathbb{Z} \\ 0 & \text{otherwise.} \end{cases}$$



## Example: upsampling by 4



## Example: upsampling by 4





## Spectral representation

$$X_{NU}(z) = \sum_{k=-\infty}^{\infty} x_{NU}[k]z^{-k}$$

$$= \sum_{k=-\infty}^{\infty} x[k]z^{-Nk}$$

$$= X(z^N)$$

$$X_{NU}(e^{j\omega}) = X(e^{j\omega N})$$

# Spectral representation

$$\begin{aligned}X_{NU}(z) &= \sum_{k=-\infty}^{\infty} x_{NU}[k]z^{-k} \\&= \sum_{k=-\infty}^{\infty} x[k]z^{-Nk} \\&= X(z^N)\end{aligned}$$

$$X_{NU}(e^{j\omega}) = X(e^{j\omega N})$$

# Spectral representation

$$\begin{aligned}X_{NU}(z) &= \sum_{k=-\infty}^{\infty} x_{NU}[k]z^{-k} \\&= \sum_{k=-\infty}^{\infty} x[k]z^{-Nk} \\&= X(z^N)\end{aligned}$$




$$X_{NU}(e^{j\omega}) = X(e^{j\omega N})$$

# Spectral representation


$$\begin{aligned}X_{NU}(z) &= \sum_{k=-\infty}^{\infty} x_{NU}[k]z^{-k} \\&= \sum_{k=-\infty}^{\infty} x[k]z^{-Nk} \\&= X(z^N)\end{aligned}$$

$$X_{NU}(e^{j\omega}) = X(e^{j\omega N})$$


## Upsampling by 2




## Upsampling by 2



## Upsampling by 2




## Upsampling by 3









## Upsampling by 3






## Upsampling by 3




## Upsampling by 4



## Upsampling by 4



## Upsampling by 4



## Upsampling: what we don't like

- ▶ in the time domain: zeros between nonzero samples are not “natural”
- ▶ in the frequency domain: extra replicas of the spectrum; can we get rid of them?


the two problems are the same!

## Upsampling: what we don't like

- ▶ in the time domain: zeros between nonzero samples are not “natural”
- ▶ in the frequency domain: extra replicas of the spectrum; can we get rid of them?


the two problems are the same!

## Upsampling in the time domain, revisited







## Upsampling in the time domain, revisited




## Upsampling in the time domain, revisited




## Upsampling in the time domain, revisited



## Upsampling in the time domain, revisited




## Zero-order interpolator




## Zero-order interpolator for 4-upsampling

$$i_0[n] = u[n] - u[n - 4]$$




## Zero-order interpolator for 4-upsampling

$$|l_0(e^{j\omega})| = \left| \frac{\sin(\frac{\omega}{2}N)}{\sin(\frac{\omega}{2})} \right| \quad N = 4$$




## Upsampling in the time domain, revisited






## Upsampling in the time domain, revisited




## first-order interpolator




## first-order interpolator for 4-upsampling

$$i_1[n] = (i_0[n] * i_0[n])/N$$




## first-order interpolator for 4-upsampling


$$|I_1(e^{j\omega})| = |I_0(e^{j\omega})|^2 / N \quad N = 4$$




in the frequency domain...




in the frequency domain...



in the frequency domain...




in the frequency domain...







in the frequency domain...




in the frequency domain...




in the frequency domain...



in the frequency domain...




## ideal digital interpolator



$$H(e^{j\omega}) = \text{rect}(\omega N / 2\pi)$$


$$h[n] = (1/N) \text{sinc}(n/N)$$

## ideal digital interpolator




# Downsampling

$$x_{ND}[n] = x[nN]$$




## Example: downsampling by 4






## Example: downsampling by 4



## Example: downsampling by 4



# Spectral representation

$$X_{ND}(z) = \sum_{k=-\infty}^{\infty} x[kN]z^{-k} = ?$$

if we can compute

$$A(z) = \sum_{k=-\infty}^{\infty} x[kN]z^{-kN}$$

then

$$X_{ND}(z) = A(z^{1/N})$$

# Spectral representation

$$X_{ND}(z) = \sum_{k=-\infty}^{\infty} x[kN]z^{-k} = ?$$

if we can compute

$$A(z) = \sum_{k=-\infty}^{\infty} x[kN]z^{-kN}$$

then

$$X_{ND}(z) = A(z^{1/N})$$

# Spectral representation

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} x[kN]z^{-kN} \\ &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \end{aligned}$$

$$\xi[n] = \begin{cases} 1 & \text{for } n = kN \\ 0 & \text{otherwise} \end{cases}$$

# Spectral representation

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} x[kN]z^{-kN} \\ &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \end{aligned}$$


$$\xi[n] = \begin{cases} 1 & \text{for } n = kN \\ 0 & \text{otherwise} \end{cases}$$

# Spectral representation

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} x[kN]z^{-kN} \\ &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \end{aligned}$$

$$\xi[n] = \begin{cases} 1 & \text{for } n = kN \\ 0 & \text{otherwise} \end{cases}$$

$\xi[n]$  for  $N = 4$







Blast from the past: DFT of  $x[n] = \delta[n]$ ,  $x[n] \in \mathbb{C}^N$

$$X[k] = \sum_{n=0}^{N-1} \delta[n] e^{-j\frac{2\pi}{N}nk} = 1$$

$x[n]$





$X[k]$





From the other side:

$$\text{IDFT} \{1\} = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn} = \delta[n]$$



From the other side:


$$\text{IDFT}\{1\} = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn} = \begin{cases} 1 & \text{for } n = 0 \\ 0 & \text{otherwise} \end{cases} \quad n = 0, \dots, N-1$$




From the other side:

$$\frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn} = \begin{cases} 1 & \text{for } n \bmod N = 0 \\ 0 & \text{otherwise} \end{cases}$$

$\tilde{X}[k]$



$\tilde{x}[n]$



# Spectral representation

$$\xi[n] = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn}$$

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} \sum_{k=-\infty}^{\infty} x[k]e^{j\frac{2\pi}{N}mk}z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m}z) \end{aligned}$$

# Spectral representation

$$\xi[n] = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn}$$

$$A(z) = \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k}$$

$$= \frac{1}{N} \sum_{m=0}^{N-1} \sum_{k=-\infty}^{\infty} x[k] e^{j\frac{2\pi}{N}mk} z^{-k}$$

$$= \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m} z)$$

# Spectral representation

$$\xi[n] = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn}$$

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} \sum_{k=-\infty}^{\infty} x[k]e^{j\frac{2\pi}{N}mk}z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m}z) \end{aligned}$$

# Spectral representation

$$\xi[n] = \frac{1}{N} \sum_{m=0}^{N-1} e^{j\frac{2\pi}{N}mn}$$

$$\begin{aligned} A(z) &= \sum_{k=-\infty}^{\infty} \xi[k]x[k]z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} \sum_{k=-\infty}^{\infty} x[k]e^{j\frac{2\pi}{N}mk}z^{-k} \\ &= \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m}z) \end{aligned}$$



## Spectral representation

$$A(e^{j\omega}) = \frac{1}{N} \sum_{m=0}^{N-1} X(e^{j(\omega - \frac{2\pi}{N}m)})$$

## Spectral representation

$$X_{ND}(z) = A(z^{1/N}) = \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m} z^{\frac{1}{N}})$$




$$X_{ND}(e^{j\omega}) = \frac{1}{N} \sum_{m=0}^{N-1} X(e^{j(\frac{\omega - 2\pi m}{N})})$$

## Spectral representation


$$X_{ND}(z) = A(z^{1/N}) = \frac{1}{N} \sum_{m=0}^{N-1} X(e^{-j\frac{2\pi}{N}m} z^{\frac{1}{N}})$$

$$X_{ND}(e^{j\omega}) = \frac{1}{N} \sum_{m=0}^{N-1} X(e^{j(\frac{\omega - 2\pi m}{N})})$$


## Downsampling by 2




## Downsampling by 2




## Downsampling by 2



## Downsampling by 2




## Downsampling by 2 with aliasing







## Downsampling by 2 with aliasing




## Downsampling by 2 with aliasing




## Downsampling by 2 with aliasing




## Downsampling by 2 with aliasing




## Downsampling by 2 with antialiasing filter




## Downsampling by 2 with antialiasing filter



## Downsampling by 2 with antialiasing filter




## Downsampling by 2 with antialiasing filter







## Downsampling by 2 with antialiasing filter




## Downsampling by 2 with antialiasing filter




# Downsampling




## Downsampling by 2, careful with highpass signals




## Downsampling by 2, careful with highpass signals




## Downsampling by 2, careful with highpass signals



## Downsampling by 2, careful with highpass signals



## Rational Sampling Rate Change






# Rational Sampling Rate Change


Example CD to DVD:

- ▶ CD:  $F_s = 44100\text{Hz}$
- ▶ DVD:  $F_s = 48000\text{Hz}$
- ▶  $\frac{N}{M} = \frac{160}{147}$
- ▶ in practice, we use time-varying local interpolation

## Subsample Interpolation



## Subsample Interpolation



# Subsample Interpolation

- ▶ we want to compute  $x(n + \tau)$ , with  $|\tau| < 1/2$
- ▶ local Lagrange approximation around  $n$

$$x_L(n; t) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(t)$$

$$L_k^{(N)}(t) = \prod_{\substack{i=-N \\ i \neq n}}^N \frac{t - i}{k - i} \quad k = -N, \dots, N$$

- ▶  $x(n + \tau) \approx x_L(n; \tau)$

# Subsample Interpolation

- ▶ we want to compute  $x(n + \tau)$ , with  $|\tau| < 1/2$
- ▶ local Lagrange approximation around  $n$

$$x_L(n; t) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(t)$$

$$L_k^{(N)}(t) = \prod_{\substack{i=-N \\ i \neq n}}^N \frac{t - i}{k - i} \quad k = -N, \dots, N$$

▶  $x(n + \tau) \approx x_L(n; \tau)$

# Subsample Interpolation


- ▶ we want to compute  $x(n + \tau)$ , with  $|\tau| < 1/2$
- ▶ local Lagrange approximation around  $n$

$$x_L(n; t) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(t)$$


$$L_k^{(N)}(t) = \prod_{\substack{i=-N \\ i \neq n}}^N \frac{t - i}{k - i} \quad k = -N, \dots, N$$

- ▶  $x(n + \tau) \approx x_L(n; \tau)$

## 2nd-order Lagrange interpolation polynomials ( $N = 1$ )




## 2nd-order Lagrange interpolation polynomials ( $N = 1$ )







## 2nd-order Lagrange interpolation polynomials ( $N = 1$ )




## Lagrange interpolation ( $N = 1$ )




## Lagrange interpolation ( $N = 1$ )




## Lagrange interpolation ( $N = 1$ )




## Lagrange interpolation ( $N = 1$ )




## Lagrange interpolation ( $N = 1$ )



## Lagrange interpolation ( $N = 1$ )



## Lagrange interpolation ( $N = 1$ )





# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

# Lagrange interpolation as an FIR

- ▶  $x(n + \tau) \approx x_L(n; \tau)$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] L_k^{(N)}(\tau)$
- ▶ define  $d_\tau[k] = L_k^{(N)}(\tau)$ ,  $k = -N, \dots, N$
- ▶  $x_L(n; \tau) = \sum_{k=-N}^N x[n - k] d_\tau[k]$
- ▶  $x_L(n; \tau) = (x * d_\tau)[n]$
- ▶  $d_\tau[k]$  is a  $(2N + 1)$ -tap FIR (dependent on  $\tau$ )

## Example ( $N = 1$ , second order approximation)

$$L_{-1}^{(1)}(t) = t \frac{t-1}{2}$$

$$L_0^{(1)}(t) = (1-t)(1+t)$$

$$L_1^{(1)}(t) = t \frac{t+1}{2}$$

## Example ( $N = 1$ , second order approximation)


$$d_{0.2}[n] = \begin{cases} -0.08 & n = -1 \\ 0.96 & n = 0 \\ 0.12 & n = 1 \\ 0 & \text{otherwise} \end{cases}$$




## CD to DVD, revisited

for every 147 CD samples, generate 160 DVD samples


## CD to DVD, revisited




CD to DVD, revisited,  $\tau[1] = 0.06875$




## CD to DVD, revisited



## CD to DVD, revisited




## CD to DVD, revisited




but  $-7\tau < -0.5$

## CD to DVD, revisited: repeat a sample



## CD to DVD, revisited





## CD to DVD, revisited

efficient local interpolation with 160 3-tap filters, used in sequence