

ISOMORPHIC DATA TRANSFORMATION

Alessandro Coglio

Kestrel Institute

© 2020

Assumptions

given isomorphic domains (see separate 'Isomorphism' notes):

$$A \xleftrightarrow[\alpha']{\alpha} A' \quad \text{and optionally} \quad G[A \xleftrightarrow[\alpha']{\alpha} A']$$

$$B \xleftrightarrow[\beta']{\beta} B' \quad \text{and optionally} \quad G[B \xleftrightarrow[\beta']{\beta} B']$$

Non-Recursive Function

old function: $f(x) \triangleq e(x)$ $f: \mathcal{U} \rightarrow \mathcal{U}$

condition: \boxed{fAB} $x \in A \Rightarrow f(x) \in B$ — $f(A) \subseteq B$ — $f: A \rightarrow B$

new function: $f'(x') \triangleq \beta(e(\alpha'(x')))$

$\vdash \boxed{f'f}$ $f'(x') = \beta(f(\alpha'(x')))$

$$\underset{\delta_{f'}}{f'(x')} = \underset{\delta_f}{\beta(e(\alpha'(x')))} = \underset{\delta_f}{\beta(f(\alpha'(x')))}$$

QED

$\vdash \boxed{f'A'B'}$ $x' \in A' \Rightarrow f'(x') \in B'$ — $f'(A') \subseteq B'$ — $f': A' \rightarrow B'$


$$x' \in A' \xrightarrow{\alpha'A'} \alpha'(x') \in A \xrightarrow{fAB} f(\alpha'(x')) \in B \xrightarrow{\beta B} \beta(f(\alpha'(x'))) \in B' \xrightarrow{f'f} f'(x') \in B'$$

QED

$\vdash \boxed{ff'}$ $x \in A \Rightarrow f(x) = \beta'(f'(\alpha(x)))$

$$\begin{aligned} x \in A &\xrightarrow{\alpha'\alpha} \alpha'(\alpha(x)) = x \\ &\xrightarrow[f'f]{x' := \alpha(x)} f'(\alpha(x)) = \beta(f(\alpha'(\alpha(x)))) = \beta(f(x)) \\ &\xrightarrow{fAB} f(x) \in B \\ &\xrightarrow{\beta'\beta} \beta'(f'(\alpha(x))) = \beta'(\beta(f(x))) = f(x) \end{aligned}$$

QED



Guards for Non-Recursive Function

$$\boxed{\forall f} \quad \gamma_{\gamma_f}(x) \wedge [\gamma_f(x) \Rightarrow \gamma_e(x)]$$

condition: $\boxed{Gf} \quad \gamma_f(x) \Rightarrow x \in A$

$$\gamma_{f'}(x') \triangleq [x' \in A' \wedge \gamma_f(\alpha'(x'))]$$

$$\vdash \gamma_f(x) \Rightarrow \gamma_{f'}(\alpha(x))$$

$$\begin{array}{l} \gamma_f(x) \xrightarrow{Gf} x \in A \xrightarrow{\alpha' \alpha} \alpha'(\alpha(x)) = x \\ \delta_{\gamma_{f'}} \xrightarrow{x' = \alpha(x)} \gamma_{f'}(\alpha(x)) = \gamma_f(\alpha'(\alpha(x))) = \gamma_f(x) \end{array}$$

QED

$$\vdash \gamma_{f'}(x') \Rightarrow \gamma_f(\alpha'(x'))$$


$$\gamma_{f'}(x') \xrightarrow{\delta_{\gamma_{f'}}} x' \in A' \wedge \gamma_f(\alpha'(x'))$$

QED

$$\vdash \boxed{\forall f'}$$

$$\omega_{f'}(x') = \cancel{\gamma_{A'}(x')} \wedge [x' \in A' \Rightarrow \cancel{\gamma_{\alpha'}(x')} \wedge \cancel{\gamma_{\gamma_f}(\alpha'(x'))}] \wedge [x' \in A' \wedge \gamma_f(\alpha'(x')) \Rightarrow \cancel{\gamma_{\alpha'}(x')} \wedge \cancel{\gamma_e(\alpha'(x'))} \wedge \cancel{\gamma_{\beta}(e(\alpha'(x')))}]$$

$$\begin{array}{l} \alpha' A' \rightarrow \alpha'(x') \in A \xrightarrow{fAB} f(\alpha'(x')) \in B \xrightarrow{\delta_F} e(\alpha'(x')) \in B \end{array}$$




Recursive Function

old function: $f(x) \triangleq \underline{\text{if } a(x) \text{ then } b(x) \text{ else } c(x, f(d(x)))}$ $f: U \rightarrow U$

$\boxed{\tau_f} \neg a(x) \Rightarrow \mu_f(d(x)) <_f \mu_f(x)$

conditions $\begin{cases} \boxed{fAB} & x \in A \Rightarrow f(x) \in B \\ \boxed{Ad} & x \in A, \neg a(x) \Rightarrow d(x) \in A \end{cases}$ — as in non-recursive case
— recursive call preserves A




new function: $f'(x') \triangleq \underline{\text{if } x' \in A' \text{ then } [\text{if } a(\alpha'(x')) \text{ then } \beta(b(\alpha'(x'))) \text{ else } \beta(c(\alpha'(x'), \beta'(f'(\alpha(d(\alpha'(x'))))))]} \text{ else } \dots$ any value (irrelevant)
 $\mu_{f'}(x') \triangleq \mu_f(\alpha'(x')) <_{f'} \triangleq <_f$

$\vdash \boxed{\tau_{f'}} x' \in A' \wedge \neg a(\alpha'(x')) \Rightarrow \mu_{f'}(\alpha(d(\alpha'(x')))) <_{f'} \mu_{f'}(x')$ — f' terminates

$x' \in A' \xrightarrow{\alpha'A'} \alpha'(x') \in A \xrightarrow{Ad} d(\alpha'(x')) \in A \xrightarrow{\alpha'\alpha} \alpha'(\alpha(d(\alpha'(x')))) = d(\alpha'(x'))$
 $\neg a(\alpha'(x'))$
 $\mu_{f'}(\alpha(d(\alpha'(x')))) \stackrel{\delta_{\mu_{f'}}}{=} \mu_f(\alpha'(\alpha(d(\alpha'(x'))))) = \mu_f(d(\alpha'(x'))) <_f \mu_f(\alpha'(x')) \stackrel{\delta_{\mu_{f'}}}{=} \mu_{f'}(x')$
 $\parallel \delta_{<_{f'}} <_{f'}$
QED

$\vdash \boxed{f'f} \quad x' \in A' \Rightarrow f'(x') = \beta(f(\alpha'(x')))$ — as in non-recursive case, with additional hypothesis




$\vdash \boxed{f'A'B'} \quad x' \in A' \Rightarrow f'(x') \in B'$ — as in non-recursive case; same proof but using $x' \in A'$ hypothesis

$$x' \in A' \xrightarrow{\alpha'A'} \alpha'(x') \in A \xrightarrow{fAB} f(\alpha'(x')) \in B \xrightarrow{\beta B} \beta(f(\alpha'(x'))) \in B' \xrightarrow{f'f} f'(x') \in B'$$

QED

$\vdash \boxed{ff'} \quad x \in A \Rightarrow f(x) = \beta'(f'(\alpha(x)))$ — as in non-recursive case; same proof but using αA for applying $f'f$



Guards for Recursive Function


$$\boxed{\sqrt{f}} \quad \gamma_{g_f}(x) \wedge [\gamma_f(x) \Rightarrow \gamma_a(x) \wedge [a(x) \Rightarrow \gamma_b(x)] \wedge [\neg a(x) \Rightarrow \gamma_d(x) \wedge \gamma_f(d(x)) \wedge \gamma_c(x, f(d(x)))]]]$$

condition: $\boxed{Gf} \quad \gamma_f(x) \Rightarrow x \in A$

$$\gamma_{f'}(x') \triangleq [x' \in A' \wedge \gamma_f(\alpha'(x'))]$$
 — as in non-recursive case

$\vdash \gamma_f(x) \Rightarrow \gamma_{f'}(\alpha(x))$ — as in non-recursive case; same proof

$\vdash \gamma_{f'}(x') \Rightarrow \gamma_f(\alpha'(x))$ — as in non-recursive case; same proof


$$\vdash \sqrt{f'}$$

$$\omega_{f'}(x') = \cancel{\gamma_{A'}(x')} \wedge [x' \in A' \Rightarrow \cancel{\gamma_{A'}(x')} \wedge \gamma_{\gamma_{f'}(A'(x'))}] \wedge$$

$$[x' \in A' \wedge \gamma_f(\alpha'(x')) \Rightarrow \gamma_{A'}(x')]^{GA'}$$

$$[x' \in A' \Rightarrow \cancel{\gamma_{a'}(x')} \wedge \gamma_a(a'(x')) \wedge$$

$$[a(a'(x')) \Rightarrow \gamma_{a'}(x') \wedge \gamma_b(a'(x')) \wedge \gamma_\beta(b(a'(x')))] \wedge$$

$$[\neg a(d'(x')) \Rightarrow \overset{Ga'}{\cancel{\neg a(d'(x'))}} \wedge \overset{rf}{\cancel{\neg a(d'(x'))}} \wedge \overset{Ga}{\cancel{\neg a(d'(x'))}}]$$

$$\alpha'(\alpha(d(\alpha'(x'))) = d(\alpha'(x'))$$

$$\alpha(d(\alpha'(\kappa'))) \in A'$$

[illegible]

$$\frac{\delta_c(\alpha(x), \beta(\alpha(d(\alpha(x)))))) \wedge G\beta}{\dots}$$

$$\Rightarrow \gamma_\beta(c(\alpha'(x'), \underbrace{f(d(\alpha'(x'))))}_{= f(d(\alpha'(x')))) \rceil \rceil \wedge$$

$$[x \notin A' \Rightarrow \sqrt{\dots}]$$

QED

Non-Recursive Predicate

old predicate: $p(x) \triangleq e(x)$ $p \subseteq \mathcal{U}$

condition: \boxed{pA} $p(x) \Rightarrow x \in A$ — $p \subseteq A$

new predicate: $p'(x') \triangleq [x' \in A' \wedge e(\alpha'(x'))]$

$\vdash \boxed{p'p}$ $x' \in A' \Rightarrow p'(x') = p(\alpha'(x'))$

$$x' \in A' \quad p'(x') \stackrel{\delta_{p'}}{=} [x' \in A' \wedge e(\alpha'(x'))] \quad \begin{matrix} \parallel \delta_p \\ p(\alpha'(x')) \end{matrix}$$

QED

$\vdash \boxed{p'A'}$ $p'(x') \Rightarrow x' \in A'$ — $p' \subseteq A'$


$$p'(x') \stackrel{\delta_{p'}}{=} x' \in A' \wedge \dots \rightarrow x' \in A'$$

QED

$\vdash \boxed{pp'}$ $x \in A \Rightarrow p(x) = p'(\alpha(x))$

$$\begin{array}{lcl} x \in A & \xrightarrow{\alpha' \alpha} & \alpha'(\alpha(x)) = x \\ & \xrightarrow{\alpha} & \alpha(x) \in A' \\ & \xrightarrow{p'p} & p'(\alpha(x)) = p(\alpha'(\alpha(x))) = p(x) \end{array}$$

QED



Guards for Non-Recursive Predicate

$$\boxed{\forall p} \quad \gamma_{\gamma_p}(x) \wedge [\gamma_p(x) \Rightarrow \gamma_e(x)]$$

$$\text{condition: } \boxed{G_p} \quad x \in A \Rightarrow \gamma_p(x)$$


$$\gamma_{p'}(x') \triangleq x' \in A'$$

$$\vdash \boxed{\forall p'}$$

$$\omega_{p'}(x') = \left[\cancel{\gamma_{A'}(x')}^{GA'} \wedge \left[x' \in A' \Rightarrow \cancel{\gamma_{A'}(x')}^{GA'} \wedge \cancel{\gamma_{\alpha'}(x')}^{G\alpha'} \wedge \cancel{\gamma_e(\alpha'(x'))}^{\forall p} \right] \right]$$

$$\begin{array}{c} \alpha'A' \searrow \\ \alpha'(x') \in A \xrightarrow{G_p} \gamma_p(\alpha'(x')) \nearrow \end{array}$$

QED



Recursive Predicate

old predicate : $p(x) \triangleq \text{if } a(x) \text{ then } b(x) \text{ else } c(x, p(d(x)))$ $p \subseteq \mathcal{U}$

$$\boxed{\tau_p} \quad \neg a(x) \Rightarrow \mu_p(d(x)) <_p \mu_p(x)$$

conditions $\begin{cases} \boxed{PA} & p(x) \Rightarrow x \in A \\ \boxed{Ad} & x \in A \wedge \neg a(x) \Rightarrow d(x) \in A \end{cases}$ — as in non-recursive predicate case
— as in recursive function case

new predicate : $p'(x') \triangleq x' \in A' \wedge \left[\text{if } a(\alpha'(x')) \text{ then } b(\alpha'(x')) \text{ else } c(\alpha'(x'), p'(\alpha(d(\alpha'(x'))))) \right]$
 $\mu_{p'}(x') \triangleq \mu_p(\alpha'(x')) \quad <_{p'} \triangleq <_p$

$\vdash \boxed{\tau_{p'}} \quad x' \in A' \wedge \neg a(\alpha'(x')) \Rightarrow \mu_{p'}(\alpha(d(\alpha'(x')))) <_{p'} \mu_{p'}(x') \quad \text{— } p' \text{ terminates — same proof as recursive function case}$

$\vdash \boxed{p'p} \quad x' \in A' \Rightarrow p'(x') = p(\alpha'(x')) \quad \text{— as in non-recursive predicate case}$

induct p'

- base) $a(\alpha'(x')) \xrightarrow{\delta_{p'}} p'(x') = b(\alpha'(x'))$
 $\xrightarrow{\delta_p} p(\alpha'(x')) = b(\alpha'(x')) \rightarrow p'(x') = p(\alpha'(x'))$
- step) $\neg a(\alpha'(x')) \xrightarrow{\delta_{p'}} p'(x') = c(\alpha'(x'), p'(\alpha(d(\alpha'(x'))))) = c(\alpha'(x'), p(\alpha'(\alpha(d(\alpha'(x'))))))$
 $\xrightarrow{\delta_p} p(\alpha'(x')) = c(\alpha'(x'), p(d(\alpha'(x'))))$
 $\xrightarrow{\alpha' \alpha} \alpha'(\alpha(d(\alpha'(x')))) = d(\alpha'(x')) \xrightarrow{\alpha A} \alpha(d(\alpha'(x'))) \in A' \xrightarrow{IH} p(\alpha'(\alpha(d(\alpha'(x'))))) = c(\alpha'(x'), p(d(\alpha'(x'))))$
 $\parallel p(\alpha'(x'))$


QED

$\vdash \boxed{p'A'} \quad p'(x') \Rightarrow x' \in A' \quad \text{— } p' \subseteq A' \quad \text{— as in non-recursive predicate case}$

$p'(x') \quad x' \notin A' \xrightarrow{\delta_{p'}} \neg p'(x') \rightarrow \text{impossible} \rightarrow x' \in A$

QED

$\vdash \boxed{pp'} \quad x \in A \Rightarrow p(x) = p'(\alpha(x)) \quad \text{— as in non-recursive predicate case ; same proof}$




same picture as non-recursive case

Guards for Recursive Predicate

$$\boxed{\forall p} \quad \gamma_{\gamma_p}(x) \wedge [\gamma_p(x) \Rightarrow \gamma_a(x) \wedge [a(x) \Rightarrow \gamma_b(x)] \wedge [\neg a(x) \Rightarrow \gamma_d(x) \wedge \gamma_p(d(x)) \wedge \gamma_c(x, p(d(x)))]]]$$

condition: $\boxed{Gp} \quad x \in A \Rightarrow \gamma_p(x)$ — as in non-recursive predicate case

$\gamma_{p'}(x') \triangleq x' \in A'$ — as in recursive predicate case



$\vdash \boxed{\forall p'}$

$$\omega_{p'}(x') = \cancel{\gamma_{A'}(x')} \wedge$$

$$[x' \in A' \Rightarrow \cancel{\gamma_{A'}(x')} \wedge$$

$$\xrightarrow{\alpha' A'} \alpha'(x') \in A \xrightarrow{Gp} \gamma_p(\alpha'(x'))$$

$$[x' \in A' \Rightarrow \cancel{\gamma_{A'}(x')} \wedge \gamma_a(\alpha'(x')) \wedge$$

$$[a(\alpha'(x')) \Rightarrow \cancel{\gamma_{A'}(x')} \wedge \gamma_b(\alpha'(x'))] \wedge$$

$$[\neg a(\alpha'(x')) \Rightarrow \cancel{\gamma_{A'}(x')} \wedge$$

$$\gamma_d(\alpha'(x')) \wedge \gamma_c(\alpha'(x'), p(d(\alpha'(x')))) \wedge$$

$$\xrightarrow{\alpha A} \alpha(d(\alpha'(x'))) \in A' \xrightarrow{p' P} p'(\alpha(d(\alpha'(x')))) = p(\alpha'(d(d(\alpha'(x')))))$$

$$\parallel \leftarrow p(d(\alpha'(x'))) \rightarrow \gamma_c(\alpha'(x'), p'(\alpha(d(\alpha'(x'))))) \right]]]$$

QED

Generalization to Tuples

$$\begin{array}{cccccc} f: \mathcal{U}^n \rightarrow \mathcal{U}^m & p \in \mathcal{U}^n & A \subseteq \mathcal{U}^n & B \subseteq \mathcal{U}^m & \alpha: \mathcal{U}^n \rightarrow \mathcal{U}^{n'} & \beta: \mathcal{U}^m \rightarrow \mathcal{U}^{m'} \\ f': \mathcal{U}^{n'} \rightarrow \mathcal{U}^{m'} & p' \in \mathcal{U}^{n'} & A' \subseteq \mathcal{U}^{n'} & B' \subseteq \mathcal{U}^{m'} & \alpha': \mathcal{U}^{n'} \rightarrow \mathcal{U}^n & \beta': \mathcal{U}^{m'} \rightarrow \mathcal{U}^m \end{array}$$

straightforward, similar to 'Isomorphisms' notes

Compositional Establishment of Isomorphic Mappings on Tuples

partition old and new inputs into equal numbers of disjoint non-empty subsets:

$$\left. \begin{aligned} \{1, \dots, n\} &= \{i_{1,1}, \dots, i_{1,n_1}\} \uplus \dots \uplus \{i_{k,1}, \dots, i_{k,n_k}\} \\ \{1, \dots, n'\} &= \{i'_{1,1}, \dots, i'_{1,n'_1}\} \uplus \dots \uplus \{i'_{k,1}, \dots, i'_{k,n'_k}\} \end{aligned} \right\} \text{same } k$$

$n_1 > 0, \dots, n_k > 0, \quad n_1 + \dots + n_k = n \geq 1$
 $n'_1 > 0, \dots, n'_k > 0, \quad n'_1 + \dots + n'_k = n' \geq 1$

establish isomorphic mappings between each pair of partitions:

$$A_1 \xleftrightarrow[\alpha'_1]{\alpha_1} A'_1, \dots, A_k \xleftrightarrow[\alpha'_k]{\alpha_k} A'_k, \quad A_1 \subseteq \mathcal{U}^{n_1}, \dots, A_k \subseteq \mathcal{U}^{n_k}, \quad A'_1 \subseteq \mathcal{U}^{n'_1}, \dots, A'_k \subseteq \mathcal{U}^{n'_k}$$

combine the isomorphic mappings:

$$\begin{aligned} A &\triangleq \{ \langle x_1, \dots, x_n \rangle \in \mathcal{U}^n \mid \langle x_{i_{1,1}}, \dots, x_{i_{1,n_1}} \rangle \in A_1 \wedge \dots \wedge \langle x_{i_{k,1}}, \dots, x_{i_{k,n_k}} \rangle \in A_k \} \\ A' &\triangleq \{ \langle x'_1, \dots, x'_{n'} \rangle \in \mathcal{U}^{n'} \mid \langle x'_{i'_{1,1}}, \dots, x'_{i'_{1,n'_1}} \rangle \in A'_1 \wedge \dots \wedge \langle x'_{i'_{k,1}}, \dots, x'_{i'_{k,n'_k}} \rangle \in A'_k \} \\ \alpha(x_1, \dots, x_n) &\triangleq \langle \alpha_1(x_{i_{1,1}}, \dots, x_{i_{1,n_1}}), \dots, \alpha_k(x_{i_{k,1}}, \dots, x_{i_{k,n_k}}) \rangle \\ \alpha'(x'_1, \dots, x'_{n'}) &\triangleq \langle \alpha'_1(x'_{i'_{1,1}}, \dots, x'_{i'_{1,n'_1}}), \dots, \alpha'_k(x'_{i'_{k,1}}, \dots, x'_{i'_{k,n'_k}}) \rangle \end{aligned} \left. \vphantom{\begin{aligned} A \\ A' \\ \alpha \\ \alpha' \end{aligned}} \right\} \text{flatten nested tuples}$$

do analogously for old and new outputs:

$$\left. \begin{aligned} \{1, \dots, m\} &= \{j_{1,1}, \dots, j_{1,m_1}\} \uplus \dots \uplus \{j_{h,1}, \dots, j_{h,m_h}\} \\ \{1, \dots, m'\} &= \{j'_{1,1}, \dots, j'_{1,m'_1}\} \uplus \dots \uplus \{j'_{h,1}, \dots, j'_{h,m'_h}\} \end{aligned} \right\} \text{same } k$$

$m_1 > 0, \dots, m_h > 0, \quad m_1 + \dots + m_h = m \geq 1$
 $m'_1 > 0, \dots, m'_h > 0, \quad m'_1 + \dots + m'_h = m' \geq 1$

$$B_1 \xleftrightarrow[\beta'_1]{\beta_1} B'_1, \dots, B_h \xleftrightarrow[\beta'_h]{\beta_h} B'_h, \quad B_1 \subseteq \mathcal{U}^{m_1}, \dots, B_h \subseteq \mathcal{U}^{m_h}, \quad B'_1 \subseteq \mathcal{U}^{m'_1}, \dots, B'_h \subseteq \mathcal{U}^{m'_h}$$

$$\begin{aligned} B &\triangleq \{ \langle y_1, \dots, y_m \rangle \in \mathcal{U}^m \mid \langle y_{j_{1,1}}, \dots, y_{j_{1,m_1}} \rangle \in B_1 \wedge \dots \wedge \langle y_{j_{h,1}}, \dots, y_{j_{h,m_h}} \rangle \in B_h \} \\ B' &\triangleq \{ \langle y'_1, \dots, y'_{m'} \rangle \in \mathcal{U}^{m'} \mid \langle y'_{j'_{1,1}}, \dots, y'_{j'_{1,m'_1}} \rangle \in B'_1 \wedge \dots \wedge \langle y'_{j'_{h,1}}, \dots, y'_{j'_{h,m'_h}} \rangle \in B'_h \} \end{aligned}$$

$$\begin{aligned} \beta(y_1, \dots, y_m) &\triangleq \langle \beta_1(y_{j_{1,1}}, \dots, y_{j_{1,m_1}}), \dots, \beta_h(y_{j_{h,1}}, \dots, y_{j_{h,m_h}}) \rangle \\ \beta'(y'_1, \dots, y'_{m'}) &\triangleq \langle \beta'_1(y'_{j'_{1,1}}, \dots, y'_{j'_{1,m'_1}}), \dots, \beta'_h(y'_{j'_{h,1}}, \dots, y'_{j'_{h,m'_h}}) \rangle \end{aligned} \left. \vphantom{\begin{aligned} \beta \\ \beta' \end{aligned}} \right\} \text{flatten nested tuples}$$