

# Analyzing the Average Complexity of QuickSelect

Karl Stroetmann

May 12, 2021

## 1 The Algorithm QuickSelect

The goal of QuickSelect is to compute the  $k$  smallest elements of a given list  $L$ , i.e. the function `quickSelect` takes two arguments:

- (a)  $L$  is a list of numbers,
- (b)  $k$  is a natural number such that  $2 \leq k \leq \text{len}(L)$ .

The function call `quickSelect(L, k)` returns a list of length  $k$  that contains the  $k$  smallest elements of  $L$ , i.e. it satisfies the following specification:

- $\text{len}(\text{quickSelect}(L, k)) = k$ ,
- $\text{set}(\text{quickSelect}(L, k)) = \text{set}(\text{sorted}(L)[:k])$ .

Note that the list returned by `quickSelect(L, k)` is not required to be sorted. It can contain the  $k$  smallest elements of  $L$  in any order.

The function can be implemented recursively via the following equations:

1.  $\text{len}(L) < k \rightarrow \text{quickSelect}(L, k) = \Omega$ ,  
because if the length of  $L$  is less than  $k$ , then there is no way to select the  $k$  smallest elements from  $L$ .
2.  $\text{len}(L) = k \rightarrow \text{quickSelect}(L, k) = L$ ,  
because if the list  $L$  has exactly  $k$  elements, then  $L$  itself is a list containing the  $k$  smallest elements of  $L$ .
3. Otherwise we assume that  $L = [x] + R$  and partition  $L$  as in [QuickSort](#), i.e. we define

$$S := [y \in L \mid y \leq x] \quad \text{and} \quad B := [y \in L \mid y > x].$$

Then there are three cases:

- (a)  $k \leq \text{len}(S) \rightarrow \text{quickSelect}(L, k) = \text{quickSelect}(S, k)$ ,
- (b)  $k = \text{len}(S) + 1 \rightarrow \text{quickSelect}(L, k) = S + [x]$ ,
- (c)  $k > \text{len}(S) + 1 \rightarrow \text{quickSelect}(L, k) = S + [x] + \text{quickSelect}(B, k - \text{len}(S) - 1)$ .

## 2 Analysis of the Average Complexity of quickSelect

Let us define  $d_{n,k}$  as the average number of comparisons of list elements that are performed when `quickSelect(L, k)` is evaluated with a list  $L$  of length  $k$ . We proceed to construct a recurrence equation for  $d_{n,k}$ .

- (a)  $d_{k,k} = 0$ ,

$$(b) \ d_{n+1,k} = n + \frac{1}{n+1} \cdot \left( \sum_{i=k}^n d_{i,k} + \sum_{i=0}^{k-2} d_{n-i,k-i-1} \right).$$

We will not be able to solve this recurrence equation exactly, but we will be able to establish an upper bound for  $d_{n,k}$  that is independent of  $k$ . To this end, we need the following Lemma.

**Lemma 1** Assume that  $n \in \mathbb{N}$  and the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is defined as

$$f(x) = 4 \cdot (x-1) \cdot (n+1-x)$$

Then we have

$$\forall x \in \mathbb{R} : f(x) \leq n^2.$$

**Proof:** The function  $f$  is quadratic in  $x$  and the coefficient of  $x^2$  is negative. By *completing the square* we find the following chain of equations and inequations:

$$\begin{aligned} f(x) &= 4 \cdot (x-1) \cdot (n+1-x) \\ &= 4 \cdot (n \cdot x + x - x^2 - n - 1 + x) \\ &= -4 \cdot (x^2 - (n+2) \cdot x + n+1) \\ &= -4 \cdot \left( x^2 - (n+2) \cdot x + \left(\frac{n+2}{2}\right)^2 + n+1 - \left(\frac{n+2}{2}\right)^2 \right) \\ &= -4 \cdot \left( x - \left(\frac{n+2}{2}\right) \right)^2 - 4 \cdot \left( n+1 - \left(\frac{n+2}{2}\right)^2 \right) \\ &\leq -4 \cdot \left( n+1 - \left(\frac{n+2}{2}\right)^2 \right) \\ &= -4 \cdot n - 4 + (n+2)^2 \\ &= -4 \cdot n - 4 + n^2 + 4 \cdot n + 4 \\ &= n^2 \end{aligned}$$

Hence we have shown that  $f(x) \leq n^2$ . □

**Theorem 2** For all  $k \in \mathbb{N}$  such  $k \geq 2$  and all  $n \in \mathbb{N}$  such that  $k \leq n$  we have

$$d_{n,k} \leq 4 \cdot n.$$

Hence the complexity of QuickSelect is linear in the length of the list and the constant of linearity is independent of  $k$ .

**Proof:** We prove the claim by induction on  $n$ .

**Base case:**  $n = k$ . In this case we have

$$d_{n,k} = d_{k,k} = 0 \leq 4 \cdot n. \quad \checkmark$$

**Induction step:**  $0, 1, \dots, n \mapsto n+1$ . Then we have the following chain of equations and inequations:

$$\begin{aligned} d_{n+1,k} &= n + \frac{1}{n+1} \cdot \left( \sum_{i=k}^n d_{i,k} + \sum_{i=0}^{k-2} d_{n-i,k-i-1} \right) \\ &\stackrel{\text{ih}}{\leq} n + \frac{1}{n+1} \cdot \left( \sum_{i=k}^n 4 \cdot i + \sum_{i=0}^{k-2} 4 \cdot (n-i) \right) \end{aligned}$$

In order to prove our claim we proceed to show that

$$\sum_{i=k}^n 4 \cdot i + \sum_{i=0}^{k-2} 4 \cdot (n-i) \leq 3 \cdot (n+1)^2$$

This claim is shown as follows:

$$\begin{aligned} \sum_{i=k}^n 4 \cdot i + \sum_{i=0}^{k-2} 4 \cdot (n-i) &= 4 \cdot \left( \sum_{i=0}^n i - \sum_{i=0}^{k-1} i + (k-1) \cdot n - \sum_{i=0}^{k-2} i \right) \\ &= 4 \cdot \left( \sum_{i=0}^n i - 2 \cdot \sum_{i=0}^{k-2} i - (k-1) + (k-1) \cdot n \right) \\ &= 2 \cdot n \cdot (n+1) - 4 \cdot (k-2) \cdot (k-1) + 4 \cdot (k-1) \cdot (n-1) \end{aligned}$$

In order to show that this expression is less or equal than  $3 \cdot (n+1)^2$  we use Lemma 1 where we have shown that for  $f(x) = 4 \cdot (x-1) \cdot (n+1-x)$  the inequation  $f(x) \leq n^2$  holds for all  $x \in \mathbb{R}$ . We have

$$\begin{aligned} f(k) &\leq n^2 \\ \Rightarrow 4 \cdot (k-1) \cdot (n+1-k) &\leq n^2 \\ \Rightarrow -4 \cdot (k-1) \cdot (k-n-1) &\leq n^2 \\ \Rightarrow -4 \cdot (k-1) \cdot ((k-2)-(n-1)) &\leq n^2 \\ \Rightarrow -4 \cdot (k-2) \cdot (k-1) + 4 \cdot (k-1) \cdot (n-1) &\leq n^2 \\ \Rightarrow -4 \cdot (k-2) \cdot (k-1) + 4 \cdot (k-1) \cdot (n-1) &\leq (n+1)^2 \\ \Rightarrow 2 \cdot n \cdot (n+1) - 4 \cdot (k-2) \cdot (k-1) + 4 \cdot (k-1) \cdot (n-1) &\leq 3 \cdot (n+1)^2 \end{aligned}$$

Combining this with our previous result we now have

$$\begin{aligned} d_{n+1,k} &= n + \frac{1}{n+1} \cdot \left( \sum_{i=k}^n d_{i,k} + \sum_{i=0}^{k-2} d_{n-i,k-i-1} \right) \\ &\stackrel{\text{ih}}{\leq} n + \frac{1}{n+1} \cdot \left( \sum_{i=k}^n 4 \cdot i + \sum_{i=0}^{k-2} 4 \cdot (n-i) \right) \\ &\leq n + \frac{1}{n+1} \cdot 3 \cdot (n+1)^2 \\ &\leq n + 3 \cdot (n+1) \\ &< (n+1) + 3 \cdot (n+1) \\ &= 4 \cdot (n+1) \quad \checkmark \end{aligned}$$

Hence we have shown that  $d_{n+1,k} \leq 4 \cdot (n+1)$  and the claim is proven by induction.  $\square$