

# Ria Health



# Experienced mission driven team.

## Clinical experts

Seasoned group of healthcare operators with experience in all facets of scaling a national practice.

**JOHN MENDELSON, MD**

20 year NIH funded Addiction Researcher, Professor of Medicine, UCSF



**CMO**

**TOM NIX**

Seasoned operator with multiple startup and turn around and exit exp.  
Former CEO of Scala



**CEO**

**PAUL LINDE, MD**

25 years' experience in emergency psychiatric care. Professor Emeritus in Psychiatry at UCSF



**MEDICAL  
DIRECTOR**

**CARE COLLABORATION  
ALISON MANKOWSKI, LCSW**

Decade of behavioral health marketing, operations and sales.  
Most recently at MDUHS



**VP CLIENT  
DEVELOPMENT**

**JC MUYL**

Multiple healthcare startup and operation leader former COO of Abilita



**Contract  
COO**

**ALEX LEE, PhD**

Multiple BIH startup and leadership and clinical leadership roles. Former coaching lead at Abilita



**VP OF Care  
Innovation**


**BOB NIX, PhD**

Serial startup CTO, Led Abilita health care development for a decade



**CTO**

# The Next Pandemic



**Mental Health Crisis**

**Economic Crisis**

**Physical Health Crisis**

## Alcohol Misuse Epidemic

- Hospital admissions for Alcohol Related Liver Disease Increased 30% Since June 2020 <sup>1</sup>
- 2009-2015 Study Reports Cirrhosis Increased 300% for people under 45 <sup>2</sup>
- Alcohol Cirrhosis rates for Women Increased 50% since 2009-2015 <sup>2</sup>

1 in 10 Adults with AUD

7%

Treated

50% Comorbidity Anxiety &  
Depression <sup>3</sup>

\$131B Lost Productivity

\$249 Billion


3rd Leading Cause of Preventable  
Death <sup>4</sup>

**Low Treatment Rates**

***High Burden***

- Stigma and shame hold people back from choosing to get help<sup>5</sup>
- Out of pocket cost for comprehensive care, create financial barriers
- Residential and IOP programs don't fit into life, too much time and absence from work, family and community

# Tele-Mental Health Adoption



# About

## Ria Health is:

Telemental health practice that leverages technology and evidence-based care to deliver a suite of convenient, accessible, effective treatment of Alcohol Use Disorder:

- Medication-Assisted Treatment
- Digital health therapeutics
- Coaching
- Peer/family support
- Daily feedback on achieving personal goals
- Hyper focused on providing high quality clinical outcomes for Alcohol Use Disorder treatment

## We:

Harness technology and modern, integrated, behavioral health treatment approaches to:

- Reduce stigma
- Remove barriers to care
- Engage & empower consumers
- Improve outcomes
- Lower the cost of care



# Individuals prefer specialized alcohol treatment

- A recent survey shows that 68% of people seeking alcohol treatment prefer an alcohol specialist over a one-size-fits-all addiction generalist.
- NSDUH Survey shows 14% of treatment seekers did not find a program that met the treatment they wanted.
- Support that's at home and private and will not disrupt their lives.
- A coach or community that understands that treatment goals may include reduction or abstinence.



**67%**

**Choose Reduction Goal**

**68%**

**Prefer an Alcohol Specialist for Treatment vs. One Size Fits All addictions generalist.**



# Member Journey

1

24/7 Phone Consult or direct online scheduled with enrollment counselor



2

Download the App



3

Meet with the Medical Team



4

Track BAC with Breathalyzer



5

Start Medication



6

Work with your Coach via video and attend groups throughout the week



7

Stay Motivated and connected to your Family & PCP through Inner Circle



8


Watch your drinking decrease



# Ria Helps Individuals Reduce Drinking

## Key Points


- Members achieve 50% average reduction in first month
- Over 12 months average reduction of more than 75%




# Ria Health's Treatment System

## Key Points

- Breakthrough Persistence more than **55% of members stick to program over 6 months**
- Ria Health's members stay on **MAT 270 Days vs. the 40-70 day average** of the current Ambulatory Medical Management treatment system<sup>1</sup>



# Ria's Value Flywheel



# Defensible Quality Outcomes

# Alcohol Risk Scores

A metric of alcohol harm reduction, the World Health Organization (WHO) risk drinking level, has validity in real-world settings and as a viable alcohol use disorder (AUD) outcomes endpoint.

Alcohol risk drinking reduction endpoint appears to maintain its efficacy in predicting improved function after extended periods of time (at least 1 year for function and up to 3 years for health costs)

| Alcohol Risk Level | Alcohol Risk Level |         |
|--------------------|--------------------|---------|
|                    | Me                 | Women   |
| 4                  | 7+/day             | 4+/day  |
| 3                  | 5-7/day            | 3-4/day |
| 2                  | 3-4/day            | 2-3/day |
| 1                  | 0-3/day            | 0-1/day |

(drinks per day)

**70% of patients reduced Alcohol risk scores at 30 Days 85% of patients reduced Alcohol risk scores at one year**



# Health Care Savings

## One level Risk Shift

3 Year Cost Savings

30%

Lower Health Care Costs

## Two Level Risk Shift

3 Year Cost Savings

50%

Lower Health Care Costs

- Improved Mental Health
- Improved Quality of Life
- Reduced Societal Burdens





# Ria Health Bundled Payment Model

• Redacted

| Phase Code               | Price Per Month |
|--------------------------|-----------------|
| Phase I (<br>Month 1-3)  | \$800/month     |
| Phase II<br>(Month 4-12) | \$200/month     |

# Alcohol Treatment TAM



# Ria Health Licensed Providers State Coverage



- National Provider Coverage
- Supports the needs of multi-state employers

## Scaling Clinical Team



- Efficient utilization with mix of Full- time, Part-time, and 1099 contracts with flexible hours
- Addiction specialists trained in Ria Health protocols
- Team scaled for Patient availability, responsiveness, and growth across 50 states
- State Licensed MD, NP (many multi- state)
- Resilient coverage with multiple providers per payer/state

AUD Treatment  
Outcomes

Join  
Monument

Groups  
Eleanor  
Health

Workit  
Health

Bicycle  
Health

Boulder Health

Quit  
Genius

Geographic Reach

- Majority of Multi Modality treatment providers focused on OUD
- OUD treatment programs require in- person in most states

# Resources

# Resources

For more specific information on Ria's program, please see the list of resources linked below:

- [Watch: A Day in the Life of a Ria Member](#)
- [Read: How the Ria Program Works](#)
- [Read: How to Stop or Reduce Drinking the Ria Health Way](#)
- [Read: Our Treatment Plan](#)
- [Read: Meet Your Medical Team](#)
- [Read: Recovery Coaching with Ria Health](#)
- [Read: The Ria Health App](#)
- [Read: Medications for Alcoholism](#)
- [Read: FAQ](#)
- [Read: Ria Health Reviews](#)

# References



# References

1. [Mellinger JL, Shelden K, Winder GS, Tappel E, Adams M, Fontana RJ, Volk ML, Blow FC, Lok ASF. The high burden of alcoholic cirrhosis in privately insured persons in the United States. Hepatology. 2018 Sep;68\(3\):872-882. doi: 10.1002/hep.29887. Epub 2018 May 20. PMID: 29579356](#)
2. [Mowbray, O., Glass, J. E., & Grinnell-Davis, C. L. \(2015\). Latent class analysis of alcohol treatment utilization patterns and 3-year alcohol-related outcomes. Journal of substance abuse treatment, 54, 21-28. <https://doi.org/10.1016/j.jsat.2015.01.012>](#)
3. [Centers for Disease Control and Prevention. Alcohol Related Disease Impact \(ARDI\) application, 2021. Available at <https://drinks.cdc.gov/>](#)
4. [Knox, J., Hasle, D. S., Larson, P., & Khandker, H. R. \(2019\). Prevention, screening, and treatment for heavy drinking and alcohol use disorder. The lancet. Psychiatry, 6\(12\), 1054-1062. \[https://doi.org/10.1016/S2053-2554\\(19\\)30211-5\]\(https://doi.org/10.1016/S2053-2554\(19\)30211-5\)](#)
5. <https://www.cdc.gov/od/ohrt/ohrt/about-us/newsroom/innovative-covid-19-trends-fact-sheet-virtual-and-telehealth.html>
6. [Substance Abuse and Mental Health Services Administration. \(2019\) Key substance use and mental health indicators in the United States: Results from the 2018 National Survey on Drug Use and Health \(NSDUH\) Publication No. PE19-5068, NSDUH Series H-54. Rockville, MD: Center for Behavioral Health Statistics and Quality, Substance Abuse and Mental Health Services Administration. Retrieved from <https://www.samhsa.gov/data>](#)
7. [Bryson, W. C., McConnell, J., Korhonen, P. T., & McCarty, D. \(2018\). Extended-release naltrexone for alcohol dependence: persistence and healthcare costs and utilization. The American journal of managed care, 17 Suppl 8\(Suppl 8\), S222-S234. \[accessed 4/1/21\]\(#\)](#)
8. [Gough, G., Heather, L., Puckett, D., Westerhold, C., Ren, X., Yu, Z., Cobb, D. W., & Liangpongkiet, S. \(2015\). The Utility of Commonly Used Laboratory Tests to Screen for Excessive Alcohol Use in Clinical Practice. Alcoholism, clinical and experimental research, 39\(8\), 1493-1500. <https://doi.org/10.1111/acer.12781>](#)

# References

- 10Mittleman, B. S., Kessler, H. R., Halperin, K. A., O'Malley, S. S., Falk, D. E., Litten, R. Z., Hahn, D. S., Menn, K. F., & Aron, R. F. (2018). Drinking Risk Level Reductions Associated with Improvements in Physical Health and Quality of Life Among Individuals with Alcohol Use Disorder: Alcoholism, (clinical and experimental research, 42(2), 2453–2465. <https://doi.org/10.1111/acer.13667> "Table is computed from results presented in the paper. Computing the results for a Level 1 reduction using some simple math and uniformly presenting "to improvements" rather than requiring the reader to understand which measures improve when they get smaller vs which measures improve when they get larger.
- 11Gruet, B. P., Goldstein, R. B., Saha, T. D., Chou, S. P., Jung, J., Zhang, H., Pelton, R. P., Ruan, W. J., Smith, S. M., Huang, B., & Hahn, D. S. (2015). Epidemiology of DSM-5 Alcohol Use Disorder: Results From the National Epidemiologic Survey on Alcohol and Related Conditions II. *JAMA psychiatry*, 72(8), 757–766. <https://doi.org/10.1093/psypsy/2015.0584>
- 12 <https://nces.ed/ipeds/data/ipedsreports/ipedsreport.php?report=8&year=2015&table=1&id=1>
- 13 Substance Abuse and Mental Health Services Administration. Treatment Episode Data Set Discharges (TEDS-D)
- 14Stuart, E. A., Barty, C. L., Dardano, J. M., Greenfield, S. F., Dudworth, K., Song, Z., Modiano, R., Kouri, E. M., Ebnesajjad, C., Chernow, M. E., & Hyskamp, H. A. (2017). Effects of accountable care and payment reform on substance use disorder treatment: evidence from the initial 3 years of the alternative quality contract. *Addiction (Abingdon, England)*, 112(5), 614–623. <https://doi.org/10.1111/add.13555>
- 15 <https://nces.ed/ipeds/data/ipedsreports/ipedsreport.php>
- 16Mazumbar, L., Chan, M., Fava, R. P., Rossi, A., & Surobello, G. (2015). Heroin Reduction in "Continuous Care" in Alcohol Abuse Disorder. *International journal of environmental research and public health*, 12(7), 9425–9434. <https://doi.org/10.3390/ijerph120709425>
- 17Zanis, G. A., Bink, J. W., Aldridge, A., Mita, D., Maki, M. J., Cooper, D. J., O'Brien, R. A., & COMBINE Core Effectiveness Research Group (2006). Cost and cost-effectiveness of the COMBINE study in alcohol-dependent patients. *Archives of general psychiatry*, 63(9), 1214–1221. <https://doi.org/10.1093/archpsyc.63.9.1214>
- 18O'Neil, K. C., de la Cruz, E., Miles, J. N., Zeffman, S., Watkins, K., Lerner, M. E., & Metcal, G. A. (2010). Exploring productivity outcomes from a brief intervention for at-risk drinking in an employee assistance program. *Addictive behaviors*, 35(3), 394–200. <https://doi.org/10.1016/j.addbeh.2009.09.007>
- 19Aron, R. F., O'Malley, S. S., Chou, S. P., Coker, R. A., Cooper, D., Donovan, D. M., Gauffand, D. R., Hoising, J. D., Johnson, B. A., LeClerc, J. S., Longenecker, R., Mason, B. J., Mathew, M. F., Miles, W. R., Petroski, H. M., Randall, C. L., Smith, R., Weiss, R. D., Wilkins, L. D., Zuckerman, A., ... COMBINE Study Research Group. (2006). Combined pharmacotherapy and behavioral interventions for alcohol dependence: the COMBINE study: a randomized controlled trial. *JAMA*, 295(15), 2003–2013. <https://doi.org/10.1001/jama.295.15.2003>
- 20Mittleman, B. S., Kessler, H. R., Halperin, K. A., O'Malley, S. S., Falk, D. E., Litten, R. Z., Hahn, D. S., Menn, K. F., & Aron, R. F. (2018). Drinking Risk Level Reductions Associated with Improvements in Physical Health and Quality of Life Among Individuals with Alcohol Use Disorder. *Alcoholism, (clinical and experimental research*, 42(2), 2453–2465. <https://doi.org/10.1111/acer.13667>

# Thank you!

Q&A

[marketing@riahealth.com](mailto:marketing@riahealth.com)

- Safe, effective evidence based care
- Accessible, affordable and personalized
- Center of excellence for AUD treatment.