

Table 14-3.12
Mean NPI-X Total Score from Week 4 through Week 24 - Windowed

	Placebo (N=79)	Xanomeline Low Dose (N=81)	Xanomeline High Dose (N=74)
Baseline			
n	79	81	74
Mean (SD)	9.5 (12.10)	8.7 (9.82)	11.9 (13.70)
Median (Range)	5.0 (0;66)	4.0 (0;32)	8.0 (0;61)
Mean of Weeks 4-24			
n	76	69	65
Mean (SD)	9.1 (11.82)	7.7 (10.09)	7.8 (10.54)
Median (Range)	4.2 (0;67)	3.2 (0;49)	2.8 (0;42)
p-value(Dose Response) [1][2]			0.142
p-value(Xan - Placebo) [1][3]		0.594	0.119
Diff of LS Means (SE)		-0.6 (1.14)	-1.8 (1.16)
95% CI		(-2.9;1.6)	(-4.1;0.5)

[1] Based on Analysis of covariance (ANCOVA) model with treatment and site group as factors and baseline value as a covariate.

[2] Test for a non-zero coefficient for treatment (dose) as a continuous variable.

[3] Pairwise comparison with treatment as a categorical variable: p-values without adjustment for multiple comparisons.

Source: programs/t-14-3-12/t-14-3-12.R16:25 Thursday, September 03, 2026

Table 14-3.12
Mean NPI-X Total Score from Week 4 through Week 24 - Windowed

	Placebo (N=79)	Xanomeline Low Dose (N=81)	Xanomeline High Dose (N=74)
p-value(Xan High - Xan Low) [1][3]			0.311
Diff of LS Means (SE)			-1.2 (1.19)
95% CI			(-3.5;1.1)

[1] Based on Analysis of covariance (ANCOVA) model with treatment and site group as factors and baseline value as a covariate.

[2] Test for a non-zero coefficient for treatment (dose) as a continuous variable.

[3] Pairwise comparison with treatment as a categorical variable: p-values without adjustment for multiple comparisons.

Source: programs/t-14-3-12/t-14-3-12.R16:25 Thursday, September 03, 2026