

Table 1. Treatment-Emergent Adverse Events with Fisher Exact Test
Analysis Population

Adverse Event	Treatment 1 n (%) [E]	Treatment 2 n (%) [E]	Fisher p-value
Overall	30 (75.0%) [152]	38 (95.0%) [188]	0.0252
ABDOMINAL PAIN	0 [0]	2 (5.0%) [2]	0.4937
AGITATION	4 (10.0%) [4]	2 (5.0%) [2]	0.6752
APPLICATION SITE DERMATITIS	6 (15.0%) [8]	2 (5.0%) [4]	0.2633
APPLICATION SITE ERYTHEMA	2 (5.0%) [2]	6 (15.0%) [12]	0.2633
APPLICATION SITE INDURATION	2 (5.0%) [2]	0 [0]	0.4937
APPLICATION SITE IRRITATION	2 (5.0%) [4]	8 (20.0%) [14]	0.0872
APPLICATION SITE PRURITUS	4 (10.0%) [10]	6 (15.0%) [10]	0.7370
APPLICATION SITE URTICARIA	0 [0]	2 (5.0%) [2]	0.4937
ASTHENIA	2 (5.0%) [4]	0 [0]	0.4937
ATRIOVENTRICULAR BLOCK FIRST DEGREE	2 (5.0%) [2]	0 [0]	0.4937
BLISTER	0 [0]	4 (10.0%) [6]	0.1156
BLOOD CREATINE PHOSPHOKINASE INCREASED	2 (5.0%) [4]	0 [0]	0.4937
BODY TEMPERATURE INCREASED	0 [0]	2 (5.0%) [2]	0.4937
CATARACT OPERATION	0 [0]	2 (5.0%) [2]	0.4937
CERVICITIS	2 (5.0%) [4]	0 [0]	0.4937
COMPLETED SUICIDE	2 (5.0%) [2]	0 [0]	0.4937
CONFUSIONAL STATE	0 [0]	2 (5.0%) [2]	0.4937
CONJUNCTIVITIS	2 (5.0%) [4]	0 [0]	0.4937
COUGH	0 [0]	2 (5.0%) [2]	0.4937
DIARRHOEA	2 (5.0%) [2]	0 [0]	0.4937
DIZZINESS	2 (5.0%) [2]	6 (15.0%) [8]	0.2633
DRUG ERUPTION	2 (5.0%) [2]	0 [0]	0.4937
DYSPEPSIA	2 (5.0%) [4]	0 [0]	0.4937
DYSURIA	2 (5.0%) [2]	0 [0]	0.4937
EAR PAIN	2 (5.0%) [4]	0 [0]	0.4937
ELECTROCARDIOGRAM ST SEGMENT DEPRESSION	2 (5.0%) [2]	0 [0]	0.4937
ERYTHEMA	4 (10.0%) [6]	10 (25.0%) [18]	0.1395
EXCORIATION	2 (5.0%) [4]	0 [0]	0.4937
FALL	2 (5.0%) [4]	0 [0]	0.4937
GASTROENTERITIS VIRAL	2 (5.0%) [2]	0 [0]	0.4937
GASTROESOPHAGEAL REFLUX DISEASE	2 (5.0%) [2]	0 [0]	0.4937
HEADACHE	4 (10.0%) [4]	2 (5.0%) [2]	0.6752
HEART RATE IRREGULAR	2 (5.0%) [8]	0 [0]	0.4937
HEMIANOPIA HOMONYMOUS	0 [0]	2 (5.0%) [2]	0.4937
HYPERHIDROSIS	0 [0]	2 (5.0%) [2]	0.4937
HYPERSENSITIVITY	0 [0]	2 (5.0%) [4]	0.4937
HYPERTENSION	0 [0]	2 (5.0%) [2]	0.4937
HYPOTENSION	0 [0]	2 (5.0%) [2]	0.4937
INSOMNIA	2 (5.0%) [4]	0 [0]	0.4937
IRRITABILITY	2 (5.0%) [4]	0 [0]	0.4937
NASAL CONGESTION	2 (5.0%) [2]	0 [0]	0.4937
NASOPHARYNGITIS	0 [0]	2 (5.0%) [4]	0.4937
NAUSEA	0 [0]	4 (10.0%) [8]	0.1156
PARKINSON'S DISEASE	2 (5.0%) [2]	0 [0]	0.4937
PELVIC PAIN	2 (5.0%) [4]	0 [0]	0.4937
POSTNASAL DRIP	2 (5.0%) [4]	0 [0]	0.4937
PRURITUS	4 (10.0%) [8]	14 (35.0%) [24]	0.0145
RASH	2 (5.0%) [6]	6 (15.0%) [12]	0.2633
SHOULDER PAIN	0 [0]	2 (5.0%) [2]	0.4937
SINUS ARRHYTHMIA	2 (5.0%) [4]	0 [0]	0.4937
SINUS BRADYCARDIA	2 (5.0%) [2]	2 (5.0%) [2]	1.0000
SKIN EXFOLIATION	0 [0]	2 (5.0%) [4]	0.4937
SKIN IRRITATION	4 (10.0%) [6]	2 (5.0%) [4]	0.6752
SOMNOLENCE	0 [0]	2 (5.0%) [4]	0.4937
STUPOR	0 [0]	2 (5.0%) [2]	0.4937
SWELLING	0 [0]	2 (5.0%) [2]	0.4937
SYNCOPE	0 [0]	2 (5.0%) [2]	0.4937
TRANSIENT ISCHAEMIC ATTACK	0 [0]	2 (5.0%) [2]	0.4937
ULCER	0 [0]	2 (5.0%) [2]	0.4937
URINARY TRACT INFECTION	2 (5.0%) [4]	0 [0]	0.4937
URTICARIA	0 [0]	2 (5.0%) [6]	0.4937
VAGINAL MYCOSIS	2 (5.0%) [4]	0 [0]	0.4937

Adverse Event	Treatment 1 n (%) [E]	Treatment 2 n (%) [E]	Fisher p-value
VISION BLURRED	0 [0]	2 (5.0%) [2]	0.4937
VOMITING	0 [0]	4 (10.0%) [6]	0.1156

n = subjects with event; E = number of events; percentages use treatment-group safety denominators.
Source: Simulated data generated by this standalone program. Generated: 27Aug2026