

Table 1. Concordance of Independent Review Committee and Investigator Response

Analysis Population

Treatment	Independent Review Committee Response	Investigator CR	Investigator PR	Investigator SD	Investigator PD	Investigator NE	Overall Concordance n (%)
Treatment 1 (N=166)	CR	2 (1.1)	10 (5.4)	1 (0.5)	4 (2.2)	8 (4.3)	33(17.9)
	PR	5 (2.7)	5 (2.7)	11 (6.0)	7 (3.8)	2 (1.1)	
	SD	9 (4.9)	5 (2.7)	6 (3.3)	4 (2.2)	13 (7.1)	
	PD	11 (6.0)	6 (3.3)	8 (4.3)	9 (4.9)	8 (4.3)	
	NE	8 (4.3)	8 (4.3)	15 (8.2)	8 (4.3)	11 (6.0)	
	Overall Concordance Rate						
Treatment 2 (N=172)	CR	2 (1.3)	3 (1.9)	10 (6.5)	6 (3.9)	2 (1.3)	28(18.2)
	PR	7 (4.5)	14 (9.1)	15 (9.7)	3 (1.9)	8 (5.2)	
	SD	8 (5.2)	9 (5.8)	4 (2.6)	2 (1.3)	4 (2.6)	
	PD	6 (3.9)	13 (8.4)	4 (2.6)	4 (2.6)	2 (1.3)	
	NE	7 (4.5)	8 (5.2)	8 (5.2)	1 (0.6)	4 (2.6)	
	Overall Concordance Rate						
Total (N=338)	CR	4 (1.2)	13 (3.8)	11 (3.3)	10 (3.0)	10 (3.0)	61(18.0)
	PR	12 (3.6)	19 (5.6)	26 (7.7)	10 (3.0)	10 (3.0)	
	SD	17 (5.0)	14 (4.1)	10 (3.0)	6 (1.8)	17 (5.0)	
	PD	17 (5.0)	19 (5.6)	12 (3.6)	13 (3.8)	10 (3.0)	
	NE	15 (4.4)	16 (4.7)	23 (6.8)	9 (2.7)	15 (4.4)	
	Overall Concordance Rate						

Cells show n (percent) among subjects with both IRC and investigator assessments. CR = complete response; PR = partial response; SD = stable disease; PD = progressive disease; NE = not evaluable.

Source: Simulated data generated by this standalone program. Generated: 30Aug2026