

Table 1. Summary of Vital Signs and Change from Baseline
Safety Population

Parameter	Visit	Type	Treatment	n	Mean	SD	Min	Median	Max
Systolic Blood Pressure (mmHg)	Baseline	Observed Value	Placebo (N=38)	48	141.91	13.47	101.6	142.50	164.5
			Active (N=44)	59	142.81	10.74	115.8	144.70	163.9
			Total (N=82)	107	142.40	11.99	101.6	142.90	164.5
	Week 4	Change from Baseline	Placebo (N=38)	39	8.65	7.35	-0.7	7.90	30.7
			Active (N=44)	52	1.13	14.48	-32.4	0.65	33.3
			Total (N=82)	91	4.36	12.47	-32.4	5.20	33.3
		Observed Value	Placebo (N=38)	39	145.78	12.48	120.8	147.10	167.3
			Active (N=44)	52	144.29	17.27	111.0	144.85	174.8
			Total (N=82)	91	144.93	15.34	111.0	145.80	174.8
	Week 8	Change from Baseline	Placebo (N=38)	33	9.26	15.51	-18.7	7.20	52.7
			Active (N=44)	46	-3.63	11.29	-40.4	-4.20	12.0
			Total (N=82)	79	1.76	14.60	-40.4	2.20	52.7
		Observed Value	Placebo (N=38)	33	147.41	16.38	107.3	153.40	178.0
			Active (N=44)	46	138.53	14.63	111.1	144.05	154.6
			Total (N=82)	79	142.24	15.91	107.3	145.80	178.0
	Week 12	Change from Baseline	Placebo (N=38)	34	9.15	11.01	-7.4	10.75	24.7
			Active (N=44)	43	2.38	15.04	-35.1	5.10	22.2
			Total (N=82)	77	5.37	13.75	-35.1	6.70	24.7
		Observed Value	Placebo (N=38)	34	147.47	13.51	108.0	154.00	162.2
			Active (N=44)	43	146.33	15.03	118.4	149.00	170.4
			Total (N=82)	77	146.84	14.29	108.0	149.70	170.4
	Week 16	Change from Baseline	Placebo (N=38)	33	8.69	12.92	-16.1	8.10	34.4
			Active (N=44)	35	1.98	13.17	-35.8	1.40	24.5
			Total (N=82)	68	5.24	13.38	-35.8	6.85	34.4
		Observed Value	Placebo (N=38)	33	147.74	10.57	128.5	150.30	175.6
			Active (N=44)	35	146.92	11.24	126.9	147.30	163.9
			Total (N=82)	68	147.32	10.84	126.9	147.85	175.6
Diastolic Blood Pressure (mmHg)	Baseline	Observed Value	Placebo (N=38)	48	74.01	10.86	56.2	74.35	95.4
			Active (N=44)	59	76.73	9.00	62.5	74.70	97.0
			Total (N=82)	107	75.51	9.92	56.2	74.70	97.0
	Week 4	Change from Baseline	Placebo (N=38)	39	5.18	8.43	-6.2	2.10	22.8
			Active (N=44)	52	2.11	7.97	-17.3	2.95	13.0
			Total (N=82)	91	3.43	8.27	-17.3	2.80	22.8
		Observed Value	Placebo (N=38)	39	71.48	7.27	61.0	69.60	85.9
			Active (N=44)	52	75.86	7.84	62.4	78.90	87.7
			Total (N=82)	91	73.99	7.87	61.0	73.30	87.7
	Week 8	Change from Baseline	Placebo (N=38)	33	6.10	10.40	-6.7	4.90	26.3
			Active (N=44)	46	-1.39	7.84	-15.9	-1.25	18.4
			Total (N=82)	79	1.74	9.68	-15.9	-0.40	26.3
		Observed Value	Placebo (N=38)	33	70.42	8.60	58.6	71.10	86.1
			Active (N=44)	46	72.35	6.50	63.7	70.00	82.8
			Total (N=82)	79	71.54	7.46	58.6	71.10	86.1
	Week 12	Change from Baseline	Placebo (N=38)	34	5.40	12.98	-16.9	1.55	30.5
			Active (N=44)	43	0.28	5.56	-10.4	-2.10	9.4
			Total (N=82)	77	2.54	9.84	-16.9	1.10	30.5
		Observed Value	Placebo (N=38)	34	70.51	10.39	57.7	69.95	89.8
			Active (N=44)	43	74.33	9.04	64.3	72.70	92.1
			Total (N=82)	77	72.65	9.78	57.7	70.50	92.1
	Week 16	Change from Baseline	Placebo (N=38)	33	8.05	13.90	-11.7	4.20	35.3
			Active (N=44)	35	0.73	7.31	-13.3	-0.10	14.0
			Total (N=82)	68	4.29	11.53	-13.3	2.25	35.3
		Observed Value	Placebo (N=38)	33	73.07	10.95	60.8	68.70	91.8
			Active (N=44)	35	74.25	10.29	60.5	70.60	89.0
			Total (N=82)	68	73.68	10.55	60.5	70.50	91.8
Pulse Rate(BEATS/MIN)	Baseline	Observed Value	Placebo (N=38)	48	69.31	10.74	52.1	64.70	96.9
			Active (N=44)	59	68.02	9.57	51.3	69.10	87.8
			Total (N=82)	107	68.60	10.08	51.3	68.10	96.9
	Week 4	Change from Baseline	Placebo (N=38)	39	5.58	12.74	-10.4	2.10	31.1
			Active (N=44)	52	-1.75	11.15	-17.0	-6.80	19.8

Parameter	Visit	Type	Treatment	n	Mean	SD	Min	Median	Max	
Weight(kg)	Week 8	Observed Value	Total (N=82)	91	1.39	12.34	-17.0	-0.10	31.1	
			Placebo (N=38)	39	72.22	9.06	59.2	68.90	94.7	
			Active (N=44)	52	67.29	10.32	57.0	63.55	102.7	
			Total (N=82)	91	69.41	10.05	57.0	67.50	102.7	
		Change from Baseline	Placebo (N=38)	33	6.26	13.95	-18.7	4.10	23.0	
			Active (N=44)	50	-4.52	10.85	-18.4	-4.70	22.6	
			Total (N=82)	83	-0.23	13.21	-18.7	-0.20	23.0	
			Placebo (N=38)	33	72.33	8.03	53.7	72.90	87.2	
		Observed Value	Active (N=44)	50	64.65	10.03	55.6	60.70	94.4	
			Total (N=82)	83	67.70	9.98	53.7	65.40	94.4	
			Placebo (N=38)	34	8.72	14.35	-18.4	10.45	29.3	
			Active (N=44)	43	-2.78	8.28	-19.4	-3.60	9.8	
	Week 12	Observed Value	Total (N=82)	77	2.30	12.66	-19.4	-0.20	29.3	
			Placebo (N=38)	34	74.07	8.97	54.1	76.25	85.3	
			Active (N=44)	43	66.23	7.62	54.5	65.50	95.1	
			Total (N=82)	77	69.69	9.07	54.1	69.00	95.1	
		Change from Baseline	Placebo (N=38)	33	5.48	7.06	-13.1	4.60	14.0	
			Active (N=44)	35	-1.82	9.39	-17.7	-2.10	12.0	
			Total (N=82)	68	1.72	9.05	-17.7	3.95	14.0	
			Placebo (N=38)	33	69.87	5.30	59.9	69.50	81.3	
		Week 16	Observed Value	Active (N=44)	35	66.03	7.50	56.5	65.70	92.8
				Total (N=82)	68	67.89	6.76	56.5	67.90	92.8
				Placebo (N=38)	48	93.78	23.99	52.8	91.85	141.9
				Active (N=44)	59	91.57	12.60	72.0	92.60	118.8
	Change from Baseline		Total (N=82)	107	92.56	18.53	52.8	92.40	141.9	
			Placebo (N=38)	48	9.16	21.19	-6.5	0.20	60.5	
			Active (N=44)	48	-0.70	10.25	-32.2	0.75	19.2	
			Total (N=82)	96	4.23	17.28	-32.2	0.45	60.5	
	Baseline		Observed Value	Placebo (N=38)	48	94.16	24.23	53.1	91.20	145.1
				Active (N=44)	48	90.83	13.13	71.1	90.75	118.9
				Total (N=82)	96	92.49	19.45	53.1	91.20	145.1
				Placebo (N=38)	39	11.23	23.17	-7.2	0.20	60.9
		Week 2	Observed Value	Active (N=44)	52	-1.10	9.98	-32.8	0.10	19.7
				Total (N=82)	91	4.18	17.91	-32.8	0.20	60.9
				Placebo (N=38)	39	90.77	25.24	51.6	87.60	145.2
				Active (N=44)	52	91.89	11.83	72.1	93.60	118.7
			Change from Baseline	Total (N=82)	91	91.41	18.67	51.6	92.90	145.2
				Placebo (N=38)	39	10.62	21.55	-6.7	0.70	56.7
				Active (N=44)	52	-0.54	9.90	-33.4	0.30	20.1
				Total (N=82)	91	4.24	16.81	-33.4	0.50	56.7
	Week 4		Observed Value	Placebo (N=38)	39	90.16	24.39	54.4	88.40	144.0
				Active (N=44)	52	92.45	11.76	72.2	94.00	118.8
				Total (N=82)	91	91.47	18.19	54.4	93.00	144.0
				Placebo (N=38)	33	13.01	23.06	-6.6	0.90	56.8
		Week 6	Observed Value	Active (N=44)	50	-1.22	10.24	-34.2	-0.25	19.7
				Total (N=82)	83	4.44	17.87	-34.2	0.20	56.8
				Placebo (N=38)	33	88.02	25.58	52.0	80.50	144.8
				Active (N=44)	50	91.79	12.46	71.4	93.25	119.0
Change from Baseline			Total (N=82)	83	90.29	18.75	52.0	89.80	144.8	
			Placebo (N=38)	34	13.16	23.09	-6.9	1.35	60.1	
			Active (N=44)	43	-2.43	8.93	-31.3	0.10	7.1	
			Total (N=82)	77	4.46	18.34	-31.3	0.40	60.1	
Week 8	Observed Value		Placebo (N=38)	34	91.53	26.83	51.4	89.35	143.2	
			Active (N=44)	43	91.42	12.94	73.7	92.50	119.1	
			Total (N=82)	77	91.47	20.13	51.4	89.70	143.2	
			Placebo (N=38)	33	13.52	23.51	-7.4	1.70	58.5	
	Week 12	Observed Value	Active (N=44)	35	-2.59	9.63	-32.0	0.30	6.8	
			Total (N=82)	68	5.23	19.41	-32.0	1.05	58.5	
			Placebo (N=38)	33	90.19	26.28	49.9	88.80	146.3	
			Active (N=44)	35	89.50	10.77	72.9	89.80	112.2	
		Change from Baseline	Total (N=82)	68	89.84	19.72	49.9	89.25	146.3	
			Active (N=44)	35	-2.59	9.63	-32.0	0.30	6.8	
			Total (N=82)	68	5.23	19.41	-32.0	1.05	58.5	
			Placebo (N=38)	33	90.19	26.28	49.9	88.80	146.3	

Mean, standard deviation, median, minimum and maximum are calculated from non-missing values.
Source: simulated data generated by this standalone program. Generated: 30Aug2026