

Table 2. Concomitant Medications by Preferred Term
Safety Population

Concomitant.Medication	Placebo (N=12)	Active (N=12)	Total (N=24)
Overall	12 (100.0%)	12 (100.0%)	24 (100.0%)
ALLOPURINOL	3 (25.0%)	5 (41.7%)	8 (33.3%)
BISOPROLOL	1 (8.3%)	5 (41.7%)	6 (25.0%)
COLECALCIFEROL	0	5 (41.7%)	5 (20.8%)
FUROSEMIDE	3 (25.0%)	5 (41.7%)	8 (33.3%)
INSULIN HUMAN	6 (50.0%)	5 (41.7%)	11 (45.8%)
NOVOMIX	4 (33.3%)	5 (41.7%)	9 (37.5%)
SIMVASTATIN	0	5 (41.7%)	5 (20.8%)
THIOCTIC ACID	0	5 (41.7%)	5 (20.8%)
CO-DIOVAN	0	4 (33.3%)	4 (16.7%)
FORMOTEROL	0	4 (33.3%)	4 (16.7%)
HIERROQUICK	0	4 (33.3%)	4 (16.7%)
LATANOPROST	0	4 (33.3%)	4 (16.7%)
LEVOTHYROXINE	4 (33.3%)	4 (33.3%)	8 (33.3%)
LOSARTAN	2 (16.7%)	4 (33.3%)	6 (25.0%)
METOPROLOL	0	4 (33.3%)	4 (16.7%)
ROSUVASTATIN	0	4 (33.3%)	4 (16.7%)
SPIRONOLACTONE	0	4 (33.3%)	4 (16.7%)
WARFARIN	3 (25.0%)	4 (33.3%)	7 (29.2%)
ATORVASTATIN	7 (58.3%)	2 (16.7%)	9 (37.5%)
GLIMEPIRIDE	0	2 (16.7%)	2 (8.3%)
GLIQUIDONE	0	2 (16.7%)	2 (8.3%)
IRBESARTAN	0	2 (16.7%)	2 (8.3%)
LINAGLIPTIN	0	2 (16.7%)	2 (8.3%)
ACETYLSALICYLIC ACID	4 (33.3%)	1 (8.3%)	5 (20.8%)
AMLODIPINE	8 (66.7%)	1 (8.3%)	9 (37.5%)
BENAZEPRIL	0	1 (8.3%)	1 (4.2%)
BETAXOLOL	2 (16.7%)	1 (8.3%)	3 (12.5%)
CAPIVEN	0	1 (8.3%)	1 (4.2%)
GLICLAZIDE	4 (33.3%)	1 (8.3%)	5 (20.8%)
INDAPAMIDE	0	1 (8.3%)	1 (4.2%)
INSULIN ASPART	0	1 (8.3%)	1 (4.2%)
METFORMIN	6 (50.0%)	1 (8.3%)	7 (29.2%)
PERINDOPRIL	2 (16.7%)	1 (8.3%)	3 (12.5%)
TRIMETAZIDINE	0	1 (8.3%)	1 (4.2%)
URAPIDIL	0	1 (8.3%)	1 (4.2%)
AKRITON	3 (25.0%)	0	3 (12.5%)
ALPRAZOLAM	2 (16.7%)	0	2 (8.3%)
BLOPRESS PLUS	1 (8.3%)	0	1 (4.2%)
ENALAPRIL	3 (25.0%)	0	3 (12.5%)
FENOFIBRATE	1 (8.3%)	0	1 (4.2%)
HYDROMOL	1 (8.3%)	0	1 (4.2%)
HYPROMELLOSE	1 (8.3%)	0	1 (4.2%)
HYZAAR	2 (16.7%)	0	2 (8.3%)
IBUPROFEN	1 (8.3%)	0	1 (4.2%)
INSULIN GLARGINE	3 (25.0%)	0	3 (12.5%)
JENTADUETO	2 (16.7%)	0	2 (8.3%)
OMEGA-3-ACID ETHYL ESTER	1 (8.3%)	0	1 (4.2%)
OMEPRAZOLE	4 (33.3%)	0	4 (16.7%)
PARACETAMOL	1 (8.3%)	0	1 (4.2%)
POTASSIUM	3 (25.0%)	0	3 (12.5%)
PROPAFENONE	2 (16.7%)	0	2 (8.3%)
QUINAPRIL	3 (25.0%)	0	3 (12.5%)
RAMIPRIL	1 (8.3%)	0	1 (4.2%)
SENNOSIDE A+B	1 (8.3%)	0	1 (4.2%)
TRAMADOL	1 (8.3%)	0	1 (4.2%)

Values are number and percentage of subjects. Preferred terms are ordered by active-treatment frequency.
Source: simulated data generated by this standalone program. Generated: 28Aug2026