

Table 2. Vital Signs by Analysis Category and Visit

Safety Population

Parameter	Visit	Category	Placebo (N=146)	Active (N=154)	Total (N=300)
Systolic Blood Pressure (mmHg)	Baseline	<120 mmHg	59 (40.4%)	33 (21.4%)	92 (30.7%)
		120-139 mmHg	31 (21.2%)	29 (18.8%)	60 (20.0%)
		140-159 mmHg	21 (14.4%)	36 (23.4%)	57 (19.0%)
		>=160 mmHg	35 (24.0%)	56 (36.4%)	91 (30.3%)
	Week 2	<120 mmHg	55 (37.7%)	60 (39.0%)	115 (38.3%)
		120-139 mmHg	35 (24.0%)	16 (10.4%)	51 (17.0%)
		140-159 mmHg	19 (13.0%)	35 (22.7%)	54 (18.0%)
		>=160 mmHg	37 (25.3%)	43 (27.9%)	80 (26.7%)
	Week 4	<120 mmHg	46 (31.5%)	35 (22.7%)	81 (27.0%)
		120-139 mmHg	32 (21.9%)	43 (27.9%)	75 (25.0%)
		140-159 mmHg	31 (21.2%)	27 (17.5%)	58 (19.3%)
		>=160 mmHg	37 (25.3%)	49 (31.8%)	86 (28.7%)
	Week 6	<120 mmHg	34 (23.3%)	41 (26.6%)	75 (25.0%)
		120-139 mmHg	28 (19.2%)	41 (26.6%)	69 (23.0%)
		140-159 mmHg	20 (13.7%)	23 (14.9%)	43 (14.3%)
		>=160 mmHg	64 (43.8%)	49 (31.8%)	113 (37.7%)
	Week 8	<120 mmHg	43 (29.5%)	27 (17.5%)	70 (23.3%)
		120-139 mmHg	39 (26.7%)	35 (22.7%)	74 (24.7%)
		140-159 mmHg	17 (11.6%)	20 (13.0%)	37 (12.3%)
		>=160 mmHg	47 (32.2%)	72 (46.8%)	119 (39.7%)
	Week 10	<120 mmHg	52 (35.6%)	31 (20.1%)	83 (27.7%)
		120-139 mmHg	18 (12.3%)	31 (20.1%)	49 (16.3%)
		140-159 mmHg	19 (13.0%)	40 (26.0%)	59 (19.7%)
		>=160 mmHg	57 (39.0%)	52 (33.8%)	109 (36.3%)
	Week 12	<120 mmHg	42 (28.8%)	29 (18.8%)	71 (23.7%)
		120-139 mmHg	25 (17.1%)	30 (19.5%)	55 (18.3%)
		140-159 mmHg	19 (13.0%)	45 (29.2%)	64 (21.3%)
		>=160 mmHg	60 (41.1%)	50 (32.5%)	110 (36.7%)
Diastolic Blood Pressure (mmHg)	Baseline	<80 mmHg	47 (32.2%)	55 (35.7%)	102 (34.0%)
		80-89 mmHg	29 (19.9%)	33 (21.4%)	62 (20.7%)
		90-99 mmHg	41 (28.1%)	28 (18.2%)	69 (23.0%)
		>=100 mmHg	29 (19.9%)	38 (24.7%)	67 (22.3%)
	Week 2	<80 mmHg	80 (54.8%)	64 (41.6%)	144 (48.0%)
		80-89 mmHg	24 (16.4%)	23 (14.9%)	47 (15.7%)
		90-99 mmHg	30 (20.5%)	44 (28.6%)	74 (24.7%)
		>=100 mmHg	12 (8.2%)	23 (14.9%)	35 (11.7%)
	Week 4	<80 mmHg	42 (28.8%)	53 (34.4%)	95 (31.7%)
		80-89 mmHg	31 (21.2%)	41 (26.6%)	72 (24.0%)
		90-99 mmHg	24 (16.4%)	19 (12.3%)	43 (14.3%)
		>=100 mmHg	49 (33.6%)	41 (26.6%)	90 (30.0%)
	Week 6	<80 mmHg	52 (35.6%)	67 (43.5%)	119 (39.7%)
		80-89 mmHg	28 (19.2%)	26 (16.9%)	54 (18.0%)
		90-99 mmHg	28 (19.2%)	22 (14.3%)	50 (16.7%)
		>=100 mmHg	38 (26.0%)	39 (25.3%)	77 (25.7%)
	Week 8	<80 mmHg	64 (43.8%)	42 (27.3%)	106 (35.3%)
		80-89 mmHg	22 (15.1%)	36 (23.4%)	58 (19.3%)
		90-99 mmHg	28 (19.2%)	50 (32.5%)	78 (26.0%)
		>=100 mmHg	32 (21.9%)	26 (16.9%)	58 (19.3%)
	Week 10	<80 mmHg	58 (39.7%)	82 (53.2%)	140 (46.7%)
		80-89 mmHg	30 (20.5%)	15 (9.7%)	45 (15.0%)
		90-99 mmHg	34 (23.3%)	32 (20.8%)	66 (22.0%)
		>=100 mmHg	24 (16.4%)	25 (16.2%)	49 (16.3%)
	Week 12	<80 mmHg	59 (40.4%)	75 (48.7%)	134 (44.7%)
		80-89 mmHg	33 (22.6%)	29 (18.8%)	62 (20.7%)
		90-99 mmHg	21 (14.4%)	24 (15.6%)	45 (15.0%)
		>=100 mmHg	33 (22.6%)	26 (16.9%)	59 (19.7%)

Values are number and percentage of subjects with a non-missing assessment in the corresponding parameter and visit.

Source: simulated data generated by this standalone program. Generated: 30Aug2026