

Table 3. Subjects Meeting Potentially Clinically Significant Vital Sign Criteria

Safety Population

Parameter	Criteria	Placebo (N=10)	Active (N=10)	Total (N=20)
Systolic Blood Pressure (mmHg)	>= 170 (mmHg)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)
	<= 90 (mmHg)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)
	Increase of >= 20 (mmHg)	10/10 (100.0%)	10/10 (100.0%)	20/20 (100.0%)
Diastolic Blood Pressure (mmHg)	>= 110 (mmHg)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)
	<= 45 (mmHg)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)
	Increase of >= 15 (mmHg)	10/10 (100.0%)	10/10 (100.0%)	20/20 (100.0%)
Respiratory Rate (breaths/min)	>= 20 (bpm)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)
	<= 10 (bpm)	0/10 (0.0%)	10/10 (100.0%)	10/20 (50.0%)
	Increase of >= 5 (bpm)	0/10 (0.0%)	0/10 (0.0%)	0/20 (0.0%)

Values are n/m (%), where n is the number of subjects meeting the criterion and m is the number with an evaluable result.
Source: simulated data generated by this standalone program. Generated: 30Aug2026