

INDEX

Note: Page numbers followed by *f* indicate figures and *t* indicate tables.

A

ACF. *See* Autocorrelation function (ACF)
Actual data, 29
Aggregate function, 57, 58
Alcohol, 473–474, 473*f*
Analysis of covariance (ANCOVA), 554, 571–572
Analysis of variance (ANOVA), 554, 584
 between-group variance, 556
 equality of groups, 556–557
 hierarchical Bayesian approach, 557–568
 null hypothesis assumption, 556–557
 within-group variance, 556
ANOVA-like analysis, 7
ANOVA-like model, 642*f*, 643–644
Anything's-possible model, 290–291, 293, 294
Apply function, 59
Argument function, 64
Array function, 50
as.vector function, 54
Autocorrelation, 282–283, 285, 466,
 506, 520
Autocorrelation function (ACF), 182, 183,
 183*f*, 184

B

Baseball batting abilities
 CSV data file, 254–255, 294
 data analysis script, 255
 hierarchy data, 222
 JAGS model specification, 256–258
 marginal posterior distributions, 256*f*, 257*f*, 260*f*
 opportunities and hits, 253–254
 position-level batting abilities, 256*f*, 258
 restricted model *vs.* full model, 291–292
 selected individuals batting abilities, 257*f*, 258
BattingAverage.csv, 294
Bayes, Thomas, 100
Bayes' factor (BF), 268, 618
 Haldane prior, 344, 345*f*, 347, 347*f*
 null hypothesis, 344, 346–347, 348, 354
 posterior distribution, 346
 ratio of proportions, 353
 ROPE, 348

 Savage-Dickey method, 354, 355
 sequential testing, 385, 389–390
Bayesian analysis, 711, 712*f*
 benefits, 519
 data structure, 723
 false alarm mitigation, 328–329
 MCMC sample posting, 723, 724
 non-NHST analysis, 722
 posterior interpretation, 723
 posterior predictive check, 724
 posterior robustness, 724
 power analysis, 724
 prior justification and description, 723
 prior knowledge, 315–317
 p values, 314–315
 raw data posting, 724
Bayesian approaches
 null value (*see* Region of practical equivalence (ROPE))
 parameter-estimation approach (*see* Model-comparison method)
Bayesian estimation, 677–682, 695
Bayesian HDI, 324–325
Bayesian inference, 152, 675
 ball diameters, 20–21, 21*f*
 bias of a coin, 112
 credibility reallocation, 16
 degree of noise, 20
 fictional Sherlock Holmes, 19
 Holmes' reasoning, 16, 17*f*
 judicial *exoneration*, 18–19, 18*f*
 manufacturing process, 20
 non-trivial probability test, 21–22
 posterior distribution, 16–18
 random measurement, 19–20
Bayesian inference using Gibbs sampling (BUGS),
 193–194
Bayesian model, 504
Bayesian model averaging (BMA), 547
Bayesian *p* value, 502
Bayesian reasoning, 3
Bayesian software, 502–503, 649, 671
Bayesian statistics, 1–2

- Bayesian variable selection, 537
 - Bayes' rule, 3, 124, 222–223, 266, 267–268, 275–276
 - binomial likelihood function, 249
 - data and parameter values, 105–108
 - definition of conditional probability, 100–101
 - MCMC methods, 115–116
 - multiple coins flipping scenario, 235
 - numerical approximation, 115
 - R code, 116–117
 - single coin flipping scenario, 224–225
 - variational approximation, 115
 - Bayes, Thomas, 100
 - BernBeta function, 138–139, 355
 - Bernoulli-beta model, 413–414
 - Bernoulli distribution, 124–125, 276, 348, 408, 623, 631
 - Bernoulli likelihood and beta prior
 - histogram, 159–160, 159f
 - normal distribution, 158
 - preliminary phase, 160–161
 - proposal distribution, 159–161
 - proposal SD, 160
 - relative posterior probability, 157–158
 - Bernoulli likelihood function, 125–126, 249
 - Bernoulli ones trick, 214
 - Bernoulli *versus* binomial, 126, 249, 253, 303
 - Beta distribution
 - central tendency, 129
 - component names, 131
 - concentration, 129
 - definition, 127
 - examples of, 127, 128f
 - mean and standard deviation, 130–131
 - mode and mean, 129–130, 130f
 - parameterization, 131–132
 - posterior beta, 132–133
 - prior knowledge, 134–138
 - R code, 138–139
 - shape parameters, 127
 - Beta function, 127
 - Beta prior distribution, 315–316
 - Beta-prior program, 639–641
 - Bias of a coin
 - Bayesian inference, 112
 - data collection, 110–111
 - data type identification, 108–109
 - departure from fairness, 108
 - descriptive model, 109
 - likelihood function, 109, 110–111, 111f
 - numerical values, 108–109
 - posterior distribution, 112–113, 136–138
 - prior distribution, 110
 - probability of heads, 108
 - Bimodal distribution, 273
 - Binomial data, 124
 - Binomial likelihood function, 249, 250–251
 - Binomial probability distribution, 303–304, 304f
 - Binomial *versus* Bernoulli, 126, 249, 253, 303
 - Bootstrapping method. *See* Resampling method
 - Broad shouldered distribution, 675
 - Brooks-Gelman-Rubin statistic, 181–182
 - Burn-in period, 179–181
- ## C
- camelBack notation, 62
 - Central tendency, 87
 - CIs. *See* Confidence intervals (CIs)
 - coda.samples function, 203
 - Coin flip, 73, 223
 - multiple coin (*see* Multiple coins flipping scenario)
 - single coin (*see* Single coin flipping scenario)
 - Colon operator, 43
 - Combine function, 42
 - Comma separated values (CSV) format, 53
 - Component-by-component vector operations, 42–43
 - Computer program, 2–3
 - Conditional logistic regression
 - binary division hierarchy, 655–656, 655f
 - data generation, 657, 657f
 - hierarchical diagrams, 660f, 661
 - ilogit function, 661–662
 - linear division, 659
 - linear separations, 659
 - mnemonic subscripts, 658
 - outcome-partition hierarchy, 661, 662
 - partition structures, 667
 - posterior distributions, 664
 - posterior parameter estimates, 664, 665f, 666f
 - regression coefficients, 664
 - response probability, 655
 - Conditional probability, 91–92, 91t
 - Confidence intervals (CIs)
 - Bayesian intuition, 324
 - confidence distributions, 323–324
 - distribution (*see* Sampling distribution)

- hypothetical population, 319
 - misconception, 322–323
 - nonrejectable parameter values, 318
 - plausibility, 323
 - point estimates, 317–318
 - stopping and testing intentions, 322–323
 - Conjugate prior, 126
 - Continuous probability distribution, 80–84
 - Corn production, 611–612, 611*f*
 - Correlated parameters, 176, 340–342, 489
 - Correlated predictors, 632–633, 633*f*
 - Count predicted variable
 - Bayesian analysis, 717–719
 - crime and drinking data, 715, 716*f*
 - eye and hair colors, 704*t*, 711–713, 712*f*
 - interaction contrasts, 713, 714*f*, 717, 718*f*
 - log-linear models, 715
 - omnibus test, 714–715
 - shrinkage, 713, 717, 718*f*
 - Covariate, 554, 569
 - Credible threshold lines, 693, 694*f*
 - Crime, 715
 - Crossover interaction, 600, 600*f*, 601
 - Cumulative normal function, 439–440, 440*f*
 - Cumulative-normal model, 700
 - Cumulative normal probabilities, 676
 - Cumulative *t* function, 700
- D**
- Data fitting, 503–504, 505*f*
 - Data frame, 51
 - Data mimicking, 490
 - Data structures
 - ANCOVA, 554
 - experiments, 553
 - JAGS, 208
 - observational studies, 553
 - single-factor ANOVA, 554
 - Degree of belief, 72–73
 - Degree of noise, 20
 - Density plots, 180–181
 - Descartes, Rene, 34
 - Dichotomous data, 124
 - Dichotomous predicted variable
 - multiple group nominal predictors, 641–646
 - multiplemetric predictors, 622–629
 - regression coefficients, 629–635
 - robust logistic regression, 635–636
 - single group nominal predictors, 638–646
 - Discrete probability distribution, 78–80
 - Double exponential distribution, 532
 - Drosophila*, 562, 566
 - Dynamic shared object (DSO), 408
- E**
- Effective sample size (ESS), 170, 184
 - Entropy, 365
 - Equally informed priors, 294–295
 - Equal-tailed interval (ETI), 342
 - Equivalent prior sample method, 366
 - Error messages, 68
 - Ethanol, 473–474
 - Expected value, 84–85
 - Exponential link function, 704*t*, 705–707, 706*t*
 - Extended hierarchical model
 - hierarchical dependencies, 251–252, 252*f*
 - JAGS model specification, 253
 - over-arching distribution, 252
 - Extended model, 502
 - Eye color, 93, 95–96, 102–103, 719
- F**
- Family income, 490–491, 501–502, 506, 508
 - Family size, 490–491, 500
 - F* distribution, 556–557
 - Fertilizer, 611
 - Fisher, Ronald, 100, 556–557
 - Fisherian approach, 100
 - Fixed-duration stopping rule, 309, 309*f*
 - Formal analysis, 268–273
 - Bayesian updating, independent beta priors, 166, 167*f*
 - beta distribution, 165
 - contour plot, 167*f*, 168
 - probability density function, 165, 166
 - recapitulation, 166
 - two-parameter space, 167*f*, 168
 - F*-ratio, 100, 556–557
 - Frequentist approach, 100
 - Fruit fly, 473–474, 563*f*, 570*f*, 571
 - Funniness rating, 685, 686*f*
 - F* test, 685
- G**
- Gamma distribution, 236–238, 237*f*
 - Gaussian distribution. *See* Normal probability density function
 - Gelman-Rubin convergence statistic, 201

- Gelman–Rubin statistic, 181–182
- Generalized linear model (GLM), 4, 449–450. *See also* Predictor variables
- count data, 422
 - metric scales, 422
 - nominal scales, 422–423
 - non-manipulated variables, 421
 - ordinal scales, 422
- genMCMC function, 372
- Gibbs sampling
- applied to posterior, 174, 174*f*, 175
 - beta distributions, 173
 - component parameter selection, 171–172
 - conditional posterior distribution, 171, 173–174
 - conditional probability, 173–174
 - disadvantages, 175, 176
 - vs.* Metropolis algorithm, 170, 175
 - Metropolis–Hastings algorithm, 170–172
 - random walk type procedure, 171
 - two-parameter example, 171, 172*f*
 - working procedure, 175
- GLM. *See* Generalized linear model (GLM)
- Grid approximations, 144, 271, 273
- multiple coins flipping scenario, 231, 232*f*, 234*f*, 235
 - single coin flipping scenario, 226, 228
- Grouped metric data
- central tendency, 554
 - deflections, 554–555
 - description, 554, 555*f*, 556
 - predicted value, 554–555
 - random variation, 554
 - sum-to-zero constraint, 555–556
- ## H
- Hair color, 54, 58, 90, 91, 93, 95–96, 102, 704–705, 711
- Haldane, J. B. S., 308
- Haldane prior, 293, 344, 345*f*
- Hamilton, William Rowan, 399, 405–406
- Hamiltonian Monte Carlo (HMC), 484
- definition, 400
 - initial random momentum, 404, 406*f*
 - kinetic energy, 401
 - leapfrog procedure, 404
 - mathematical theory, 405–406
 - Metropolis acceptance probability, 403
 - posterior density, 400
 - potential function, 401
 - proposal distribution, 400–401, 402*f*
 - Steps*Eps, 402–403, 404, 405*f*
- Happiness rating, 671, 682, 691
- HDIs. *See* Highest density intervals (HDIs)
- Head-biased factories
- hierarchical diagram, 269, 269*f*
 - marginal likelihood function, 270
 - posterior distribution, 273
 - posterior probabilities, 270–271
 - prior distribution, 270–271
 - prior probabilities, 271
- Heavy-tailed distribution, 700
- Height, 25, 27, 72, 129–130, 168, 420, 421, 429, 430, 447, 477, 621, 622, 626–629
- Heterogeneous-variance model, 605–606, 605*f*
- Heterogeneous variances and robustness
- contrasts of means, 578
 - difference of scales, 578–579
 - gamma distribution, 573–574
 - groups with different variances, 575, 576*f*, 577*f*
 - hierarchical diagram, 573–574, 574*f*
 - JAGS implementation, 574–575
 - t* distribution, 573–574
 - within-group standard deviation, 575–578
- HGN.csv file, 54
- HGNdf variable, 54
- Hierarchical Bayesian approach
- baseline parameter, 557, 558
 - contrasts, 565–567
 - control types, fruit fly longevity, 561–562
 - data and posterior distribution, 562, 563*f*
 - generic noncommittal prior distributions, 557
 - half-Cauchy/folded-*t* distribution, 558–559
 - heavy-tailed distribution, 559–560
 - hierarchical diagram, 557, 558*f*, 588–589, 588*f*
 - JAGS model, 560–561, 562, 589–590
 - main effect contrast, 595–596, 596*f*
 - MCMC chain diagnostics display, 564
 - multiple comparisons, 567
 - normal distributions with homogeneous variances, 564
 - posterior distributions, 558–559
 - posterior predictive distributions, 563*f*, 564
 - prior beliefs, 558–559
 - salary data, 590–594, 592*f*
 - shrinkage, 559, 567, 568, 598
 - “simple” comparisons, 597, 597*f*
 - smryMCMC function, 594, 595*t*

- thinSteps=10 argument, 562–564
 - top-level distribution, 589
 - two-group case, 568
 - uncertainty, 598
 - Hierarchical linear regression, 573
 - Hierarchical MCMC
 - autocorrelation, 282–283, 285
 - JAGS, 278, 279
 - posterior distribution, 279, 280*f*, 283
 - posterior probabilities, 287
 - prior distributions, 279, 280*f*
 - pseudopriors, 281, 282, 283, 284*f*, 285, 286*f*
 - Hierarchical models, 4, 602, 603*f*
 - coin flip, 223
 - definition, 221
 - dependencies, 223
 - extended hierarchical model, 251–259
 - joint parameter space, 222–223
 - MLE, 247, 249
 - parameterization, 223
 - shrinkage, 245, 247, 248*f*
 - Hierarchical regression
 - dependencies model, 491, 493*f*
 - fictitious data, 491, 492*f*
 - JAGS model, 493–494
 - parameter estimation, 490
 - posterior distribution, 495
 - slope estimation, 491
 - Highest density intervals (HDIs), 6, 27–28, 87–89,
 - 88*f*, 184, 185, 185*f*, 186
 - equal-tailed intervals, 725
 - unimodal probability distribution, 726–727, 726*f*
 - HMC. *See* Hamiltonian Monte Carlo (HMC)
 - Holmes, Sherlock, 16, 17–18, 19, 20, 27–28, 30–31
 - Homogeneity of variance, 479, 510, 556–557
- I**
- ilogit function, 625
 - Income, family, 420, 490–491, 495–498, 497*f*, 501–502, 506, 508
 - initsList function, 201–202
 - Insightful comments and suggestions, 10
 - Interaction factors, 584–586, 585*f*, 587
 - Inverse-link function, 435–436
 - baseline and deflection, 710
 - cell probability, 710–711
 - MCMC process, 709
 - predictor variables, 435–436
 - two-factor ANOVA, 708–709, 709*f*, 710
 - Island-hopping politician, 161
 - position probability, 149, 150–151, 150*f*
 - proposal distribution, 151
 - proposed and current island, 147
 - proposed position, 151
 - random-walk process, 152
 - relative population, 147, 148*f*
 - target distribution, 149–150, 156
 - trajectory, 147–148, 148*f*, 149
- J**
- Jacobian matrix, 730–732
 - JAGS/Stan model, 6, 239–240, 277–278, 279, 280–281, 285, 466–468, 486–487, 488–489, 503, 506–508. *See also* Softmax regression
 - baseball batting ability, 256–258
 - baseline and deflection, 710
 - Bayesian approaches, 351–352
 - Bernoulli, Jacob, 109, 308
 - Bernoulli distribution, 195–196, 196*f*, 408
 - Bernoulli likelihood function, 249
 - beta density, 195
 - beta distribution, 195–197
 - binomial likelihood function, 249, 250–251
 - cell probability, 710–711
 - censored values, 734
 - condensed highlights, 207
 - credible parameter values, 735
 - CSV file, 208
 - data analysis, 732, 733*f*
 - data structure, 208
 - diagram of model, 209–210, 209*f*
 - extended hierarchical model, 253
 - file naming system, 206–207
 - genMCMC function, 207
 - hierarchical Bayesian, 560–561
 - high-level script, 210–211
 - ilogit function, 625
 - independent beta-distributed prior, 209
 - installation, 194, 407
 - JAGS, 408, 409, 414
 - Jags-ExampleScript.R file, 206
 - likelihood function, 195, 414
 - limitations, 415
 - limited duration, 732
 - loading data into R, 197–198
 - logical flow, 196–197

JAGS/Stan model (*Continued*)

- logistic function, 622, 625
 - log probability, 411–412
 - MCMC process, 709
 - model expansion, 218
 - model specification, 198–200, 410–411
 - multivariate model, 734–735
 - normal and interval distribution, 734–735
 - parallel processing, 215–218
 - plotMCMC function result, 211, 211f
 - power analysis, 416
 - prior distribution, 211–213, 412–413
 - probability distributions, 213–215
 - R commands, 735–736
 - relation of R programming language, 194, 194f
 - RStan library, 408, 409
 - RStudio script, 197
 - runjags package, 251
 - sampling command, 408
 - specification, 735
 - standardized parameter values, 624–625
 - string, 407–408
 - transformed parameters, 415–416
 - two-factor ANOVA, 708–709, 709f, 710
 - variable declaration, 407–408, 414–415
- Joint parameter space, 222–223, 224–225, 233–235, 249
- Joint posterior distribution, 228
- Joint prior distribution, 226–228
- Joint probability distribution, 90

L

- Laplace, Pierre-Simon, 100
- Lasso regression method, 532
- Least-squares estimates, 689
- Likelihood distribution function, 228
 - multiple coins flipping scenario, 233
 - single coin flipping scenario, 223–224, 228
- Likelihood function, 125, 266
- Likert scales, 681–682
- Linear model, 506
- Linear regression, 26, 27. *See also* Multiple linear regressions; Simple linear regression
- Load function, 56
- Logarithmic scale, 599–600
- Logistic function
 - gain indicates, 436, 437f
 - orientation, 437
 - position, 437

terminology, 438–439

threshold, 436–437, 438f

- Logistic regression. *See also* Multiple logistic regressions
- correlated predictors, 632–633, 633f
 - dependency diagram, 623–624, 624f
 - gender prediction, 626–629, 627f, 628f
 - JAGS code, 625–626
 - log odds, 629–631
 - metric predictors' interaction, 633–635, 634f
 - 0's and 1's data, 631–632, 632f
- Logit function, 438–440, 621–622, 624–625, 629–630
- Log-linear models, 715
- Long-run relative frequency, 74–76

M

- Marginal likelihood function, 274–278
- head-biased factories, 270
 - tail-biased factories, 270
- Marginal posterior distribution, 273, 627f, 693
- baseball batting ability, 256f, 257f, 260f
 - therapeutic touch, 243–244, 243f
- Marginal prior distribution, 273
- multiple coins flipping scenario, 233
 - single coin flipping scenario, 226–228
 - therapeutic touch, 245, 246f
- Marginal probability distribution, 90
- Margin of error, 498
- Markov, Andrey, 178
- Markov chain Monte Carlo (MCMC), 3, 4, 115–116
- accuracy, 182–187
 - autocorrelation plot, 203–204
 - Bayesian inference, 152
 - beta distribution, 145, 146f
 - convergence diagnostics, 203–204, 204f
 - density plot, 203–204
 - diagMCMC function, 204–205
 - efficiency, 187–188
 - function I, 204–205
 - generation, 202–203
 - goals, 178
 - hierarchical, 278–287
 - initial values, 200–202
 - Metropolis algorithm, 146–156
 - nonhierarchical, 274–278
 - one-parameter models, 144
 - plotPost function, 205–206

- representativeness, 178–182
- sampling, 515–516
- terminology, 177–178
- trace plot, 203–204
- Markov representations of Bayes' rule, 118, 119
- Mathematical model, 31
- Matrix command, 48
- Maximum likelihood estimate (MLE), 200, 247, 249, 459, 460*f*, 622, 623*f*
- MCMC. *See* Markov chain Monte Carlo (MCMC)
- Mean, 84–85
- Mean centering method, 485
- Median of distribution, 87
- Metric-predicted variable
 - dependency, normal distribution, 455, 455*f*
 - gamma distribution, 454
 - GLM, 449–450
 - group identifier, 469
 - JAGS, 456–458, 457*f*, 462–464
 - mean, 452–453
 - model block, 469
 - with multiple metric predictors, 509–552
 - with multiple nominal predictors, 583–620
 - NHST, 470–472
 - noise distribution, 472–473
 - normal distribution, likelihood function, 450, 451*f*
 - with one metric predictor, 477–508
 - with one nominal predictor, 553–582
 - parameter estimation, 450
 - posterior distribution, 470, 471*f*
 - posterior mean, 453
 - posterior precision, 453
 - response time data, 473
 - skewed data, 472
 - standard deviation, 451–452
- Metric predictors
 - additive combination, 425–427, 426*f*
 - baseline, 569–570
 - covariate, 569
 - expected value, 569
 - hierarchical diagram, 569, 570*f*
 - hierarchical linear regression, 573
 - JAGS implementation, 569–570
 - NHST, 445
 - non-additive interaction, 427–428, 428*f*
 - ordinal predicted variable, 685–698
 - posterior distribution, fruit fly longevity, 571, 572*f*
 - single metric predictor, 423–425, 425*f*
 - traditional ANCOVA, 571–572
- Metropolis algorithm
 - Bayes' rule, 161
 - Bernoulli likelihood and beta prior, 157–161
 - bivariate normal distribution, 168–169
 - disadvantage, 170
 - ESS, 170
 - vs.* formal analysis and grid approximation, 169–170, 169*f*
 - infinite and finite random walks, 170
 - matrix arithmetic, 153, 154
 - matrix multiplication, 154–155
 - Metropolis, Nicholas, 156
 - multidimensional continuous parameter space, 157
 - posterior distribution, 168–170
 - proposal distribution, 157
 - θ -hopping coin bias, 161, 162
 - transition matrix, 155–156
 - transition probabilities, 153
- Model averaging, 289
- Model-comparison method
 - alternative-hypothesis prior, 354
 - Bayes' factor, 347–348, 347*f*, 353–354, 618
 - Bernoulli distribution, 348
 - factor-deflection model, 617–618
 - Haldane prior, 344, 345*f*
 - hierarchical diagram, 352–353
 - interaction deflections, 618
 - JAGS, 351–352, 616–617
 - large-scale model, 349–350
 - null hypothesis, 344, 346–347, 354
 - null-value assessment, 354–355
 - omega model, 349, 350*f*
 - omnibus test, 616
 - one-mode model, 349
 - partitioning, 349–350, 356
 - posterior distribution, 346
 - posterior probability, 616–617
 - uninformed distribution, 343
- Mode of distribution, 87
- Monte Carlo simulation
 - complex models, 375
 - goalTally matrix, 373–374
 - HDIOfICDF function, 374–375
 - hypothetical parameter distribution, 373–374
 - JAGS analysis, 372

- Monte Carlo simulation (*Continued*)
 - output, 375
 - R script, 372
 - Movie rating, 694*f*, 695, 696*f*, 702*f*
 - Multinomial logistic regression. *See* Softmax regression
 - Multiple coins flipping scenario
 - full prior distribution, 233
 - grid approximations, 231, 232*f*, 234*f*, 235
 - hierarchical dependencies, 231*f*
 - likelihood function, 233
 - marginal prior distribution, 233
 - posterior distribution, 233–235
 - Multiple group nominal predictors, 641–646, 642*f*, 645*f*
 - Multiple linear regressions
 - correlated predictors, 513
 - credible regression coefficients, 524
 - definition, 509
 - grid representation, 510, 511
 - hierarchical diagram, 514–515, 515*f*, 531–532, 531*f*
 - interpretive perils, 512–513
 - JAGS model specification, 516
 - MCMC sampling, 515–516
 - normal distribution, 510, 511*f*, 512*f*
 - posterior distribution, 517–519, 518*f*
 - prior information, sparse data, 524
 - redundant predictors, 519–523
 - robust, 514–515
 - robustness check, 524–525
 - SAT score, 513, 514*f*
 - Multiple logistic regressions, 623–624, 624*f*
 - Multiplicative interaction
 - additive model, 527
 - different interpretations, 526, 526*f*
 - GPA, 526–527
 - non-additive interaction, 525
 - posterior distribution, 528, 529*f*, 530
 - Spend and PrcntTake, 527–528
 - Must-be-fair model, 290–291, 293, 294
- N**
- Natural frequency representation of Bayes' rule, 118
 - NHST. *See* Null hypothesis significance testing (NHST)
 - Noise, 698
 - distribution, 288
 - parameters, 500–501
 - Noise data
 - dichotomous scale, 444
 - linear regression, 441, 442*f*
 - logistic regression, 441, 443*f*
 - probability density function, 440–441
 - Nominal predicted variable, 649–670
 - Nominal predictors
 - additive combination, 430–432, 432*f*
 - non-additive interaction, 432–435, 434*t*
 - single nominal predictor, 429–430, 431*f*
 - Non-additive effects, 599
 - Non-crossover interaction, 600, 600*f*, 601–602
 - Nonhierarchical MCMC
 - Bayes' rule, 275–276
 - Bernoulli distribution, 276
 - JAGS, 277–278
 - marginal likelihood, 275
 - marginal posterior distribution, 276–277
 - probability distributions, 274
 - Non-normal distribution, 675
 - Nonparametric models, 30
 - Non-trivial probability test, 21–22
 - Normal distribution, 23, 27, 673*f*, 674
 - Normal probability density function, 83–84, 83*f*
 - Null hypothesis rejection, 5
 - Null hypothesis significance testing (NHST), 5, 5*t*, 28
 - Bayesian posterior, 328
 - Bonferonni correction, 326
 - counterproductive incentive structure, 326–327
 - experimentwise false alarm rate, 326
 - false alarm mitigation, Bayesian analysis, 328–329
 - per-comparison false alarm rate, 325–326
 - planned comparison, 326, 327
 - post hoc comparison, 326, 327
 - primary goal of, 318
 - prior knowledge, 315
 - sampling distributions, 329–331
 - soul searching, 327–328
 - space of possible outcomes, 298
 - stopping rule, 299–300
- O**
- Omnibus test, 616, 713–715
 - One-tailed probability, 305
 - OpenBUGS, 193–194

- Ordinal predicted variable
 - examples of, 672, 673*f*
 - metric predictors, 685–698
 - posterior prediction, 698–699
 - single group, 675–682
 - two group, 682–685
- Over-arching distribution, 252
- P**
 - Pairwise posterior parameters, 636, 638*f*
 - Parallel processing, JAGS
 - coda-oriented graphical functions, 217
 - considerations, 218
 - plots argument, 217
 - rjags *vs.* runjags, 216–217
 - runjags package installation, 216
 - running chains in multiple cores, 215–216
 - summarise argument, 217
 - Parameter estimation, 450, 490, 504
 - Parameter values
 - data histogram, 23–24, 23*f*
 - location parameter, 23
 - mathematical form, 24–25
 - mathematical formulas, 22–23
 - normal distribution, 23
 - posterior predictive check, 22
 - scale parameter, 23
 - plotPost function, 340
 - Poisson, Simon-Denis, 707
 - Poisson distribution, 308
 - Poisson exponential model
 - ANOVA model, 705, 706–707
 - data structure, 704, 704*t*
 - inverse-link function (*see* JAGS/Stan model)
 - linear model, 705–706, 706*t*
 - marginal proportions, 704*t*, 705
 - noise distribution, 707–708, 708*f*
 - Poisson noise distribution, 707–708, 708*f*
 - Poisson zeros trick, 214, 215
 - Posterior beta
 - Bayes' rule, 132
 - prior distribution *vs.* likelihood function, 133, 134*f*
 - Posterior distribution, 16–18, 27–28, 228, 489–490, 500, 678*f*, 679, 683, 684*f*, 685, 686*f*, 691, 692*f*
 - Bayes' rule, 111–115, 111*f*, 164–165
 - formal analysis, 165–168
 - head-biased factories, 273
 - hierarchical MCMC, 279, 280*f*, 283
 - with hierarchical shrinkage, 534, 535*f*
 - marginal, regression coefficients, 539, 539*f*, 540–542, 541*f*
 - Metropolis algorithm, 168–170
 - multiple coins flipping scenario, 233–235
 - multiple linear regression, 517–519, 518*f*
 - multiplicative interaction, 528, 529*f*, 530
 - prior distribution, 520, 522*f*
 - redundant predictors, 520, 521*f*
 - single coin flipping scenario, 227*f*, 228–230, 229*f*
 - tail-biased factories, 273
 - without hierarchical shrinkage, 532–534, 533*f*
 - Posterior HDI, 315–316, 316*f*
 - Posterior odds, 268
 - Posterior prediction, 698–699
 - Posterior predictive check, 22, 28–29, 501–502
 - Posterior predictive distribution, 644, 689, 690*f*
 - Power analysis
 - audience-appropriate prior, 366–367
 - average length criterion, 365
 - Bayesian analysis, 378–379
 - BEST, 383
 - beta distribution, 371–372, 371*t*
 - classical approach, 383
 - data consistency, 378
 - data-generating distribution, 370–371
 - data set simulation, 380–381
 - definition, 361
 - entropy, 365–366
 - flow information, 362–364, 363*f*
 - goal achievement, 360–361, 379–380
 - group-level modes, 381–382
 - hierarchical model, 376
 - hypothetical distribution, 366–367
 - idealized hypothesis, 382–383
 - Jags-Ydich-XnomSubj-MbinomBetaOmegaKappa-Power script, 377
 - minimal sample size, 364–365, 367, 367*t*
 - parameter values, 362
 - posterior distribution, 371, 376–377
 - posterior estimation, 362, 363
 - primary methods, 361
 - repetition, 363
 - therapeutic-touch practitioners, 377–378

- Predictor variables. *See also* Metric predictors;
 Nominal predictors
 cumulative normal function, 439–440, 440f
 dichotomous variable, 445
 inverse link function, 435–436
 logit function (*see* Logistic function)
 non-identity link function, 436
 Poisson distribution, 445
 statistical consultant, 445–446
- Preposterior marginal distribution, 368
- Price, Richard, 100
- Prior distribution, 27, 226, 266
 head-biased factories, 270–271
 single coin flipping scenario, 223–224, 227f, 229f, 230
 tail-biased factories, 270–271
- Prior odds, 268
- Prior probability, 266
- Probability
 long-run relative frequency, 74–76
 possibilities and, 77–78
 subjective belief, 76–77
- Probability density function, 79f, 80, 81f, 82–83, 126
- Probability distribution, 78, 274. *See also*
 Continuous probability distribution; Discrete probability distribution
 distribution list, 213–214
 likelihood function, 214–215
- Probability mass function, 78–80
- proc.time function, 66
- Programming languages, 2–3
- Proportion of variance, 517–519
- Pseudopriors, 281, 282, 283, 284f, 285
- Pseudo-random number generators (PRNGs), 74
- p values, NHST
 analysis of variance, 312–313
 Bayesian analysis, 314–315
 Bayesian HDI, 324–325
 cloud of imaginary outcomes, 299, 299f
 cloud of imaginary possibilities, 311, 312f
 data collection duration, 300–301
 definition, 298–299, 299f
 extremeness, 301–302
 fairness hypothesis, 300
 imaginary sample of flips, 311
 intention to fix duration, 308–310
 intention to fix number of flips (N), 302–305
 intention to fix number of heads (z), 305–308
 left- and right-tail p value, 310–311
 likelihood function, 300, 301
 one tailed and two-tailed p values, 301–302
 sample generation process, 300–301
 sampling distribution, 301
 soul searching, 313–314
 space of possible outcomes, 302, 302f
- ## Q
- Quadratic coefficients, 503
- Quadratic component, 496, 498–499
- Quadratic trend, 496
- ## R
- Random-outcome model, 700
- Rat diet, 446, 474
- Rat longevity, 446, 474
- R code
 for coin flipping, 94–95
 grid approximation, 725–726
 ICDF, 728
 MCMC sample, 727–728
 optimize routine, 728–729
 unimodal probability density function, 728
- Realistic data, 3
- Real-world process, 479
- Real-world sampling constraints, 309–310
- Region of practical equivalence (ROPE)
 credible parameter values, 336
 definition, 338–339
 equal-tailed interval, 342–343, 342f
 HDI, 337, 342–343, 342f
 indication, 336
 joint distribution, 341–342, 341f
 marginal distribution, 340–341, 341f
 MCMC, 339
 NHST, 339–340
 null hypothesis testing, 337–338
 null-value assessment, 354–355
 plotPost function, 340
 posterior distribution, 339, 353
 prior probability, 353
 size determination, 338
- Regression. *See also* Logistic regression; Multiple linear regression; Robust linear regression; Softmax regression
 conditional logistic (*see* Conditional logistic regression)
 hierarchical linear, 573

- linear, 26, 27
 - simple linear, 478–479, 478*f*
 - Regression coefficients
 - degree of vagueness, 539–540
 - dichotomous predicted variable, 629–635
 - marginal posterior distribution, 539, 539*f*, 540–542, 541*f*
 - shrinkage, 530–536
 - variable selection, 537–538
 - Relative posterior probabilities, 268
 - Relevant data identification, 25, 26*f*
 - Reparameterization
 - Jacobian matrix, 730–732
 - probability distribution, 729, 730
 - probability mass, 729
 - transformed parameters, 730–732
 - uniform density, 730, 731*f*
 - Replicate (rep) function, 44
 - Resampling method, 31
 - R function
 - analytical derivation, 368
 - beta function, 368
 - data-generating distribution, 370
 - initSampSize argument, 369–370
 - minimum sample sizes, 372
 - minNforHDIpower function, 369–370
 - N* flips, 368
 - preposterior marginal distribution, 368
 - Rhyming couplets, 8
 - Richter scale, 599–600
 - Robust against outliers, 635
 - Robust linear regression
 - data standardization, MCMC sampling, 484–487
 - hierarchical dependencies, 480, 480*f*, 483
 - posterior distribution, 489–490
 - regression lines and *t*-noise distributions, 480–483, 481*f*, 482*f*
 - Stan, 487–489
 - Robust logistic regression, 635–636, 637*f*, 638*f*
 - ROPE. *See* Region of practical equivalence (ROPE)
 - R programming language, 3
 - argument function, 64
 - arithmetic and logical operators, 39–40
 - asqplusb function, 64
 - assignment operator, 40
 - categorical values, 46
 - colon operator, 43
 - combine function, 42
 - command-line interface, 36
 - component-by-component vector operations, 42–43
 - data frame, 51
 - data saving, 55–56
 - debugging, 67–69
 - default value, 64–65
 - double equal sign, 41–42
 - factor function, 46–47
 - for* loop, 65
 - graphical plots, 69
 - help.search function, 39
 - if-else structure, 65
 - installation, 35
 - levels argument, 48
 - levels of factor, 46
 - list structure, 51
 - logical sequence, 45
 - matrix and array, 48–51
 - named components, 51
 - names, elements of vector, 45
 - newbie programmers, 61
 - numerical position, 45
 - plot function, 37, 39
 - processing time measurement, 66–67
 - program/script definition, 36
 - read.csv and read.table functions, 53–55
 - replicate function, 44
 - RStudio, 35, 38
 - sequence function, 43
 - simple graph, 37, 37*f*
 - SimpleGraph.R file, 38
 - source *vs.* load function, 63–64
 - utility functions, 56–61
 - variable names, 61–62
 - vector, 36–37
 - working directory, 62–63
 - RStudio, 35
 - RunJAGS package, 215
- ## S
- Salary
 - professor salary, 591–592, 594, 595*t*, 606
 - teacher salary, 543–546
 - Salary data, 599, 602–603, 604*f*, 605
 - Sample space, 72
 - Sampling distribution, 28
 - with different θ values, 321–322, 322*f*
 - experiment planning, 329–330

- Sampling distribution (*Continued*)
 - with fixed duration, 320, 321*f*
 - with fixed N , 318, 319, 319*f*
 - with fixed z , 319–320, 320*f*
 - model predictions, 330–331
 - Savage-Dickey method, 354, 355
 - save command, 55–56
 - Scholastic aptitude test (SAT), 420, 446, 509, 513, 514*f*, 517, 524, 527, 528, 532–534, 538, 539–540, 543–546
 - Seaweed, 618–619
 - Sequence (Seq) function, 43
 - Sequential testing
 - Bayes' factor, 385, 386*f*, 387*f*, 389–390
 - coins flips, 385, 386*f*, 387*f*
 - HDI, 385–388, 386*f*, 387*f*
 - HDI-with-ROPE rule, 390, 391
 - NHST, 383, 384
 - null hypothesis, 388, 389, 389*f*
 - null value, 384
 - p values, 385, 386*f*, 387*f*
 - ROPE, 386*f*, 387*f*, 388
 - Sherlock Holmes, 16, 17–18, 19, 20, 27–28, 30–31
 - Shrinkage, 245, 247, 248*f*
 - hierarchical Bayesian approach, 559
 - hierarchical, variable selection, 542–544, 545*f*
 - and prediction, posterior distribution, 495
 - regression coefficients, 530–536
 - Simple linear regression, 478–479, 478*f*
 - Single coin flipping scenario
 - Bayes' rule, 224–225
 - grid approximation, 226, 228
 - hierarchical dependencies, 224, 225*f*
 - joint posterior distribution, 228
 - joint prior distribution, 226–228
 - likelihood function, 223–224, 228
 - marginal prior distribution, 226–228
 - parameterization, 226
 - posterior distribution, 227*f*, 228–230, 229*f*
 - prior distribution, 223–224, 227*f*, 229*f*, 230
 - Single group nominal predictors, 638–641, 640*f*
 - Single metric predictor, 423–425
 - Softmax regression
 - dcat distribution, 659–661
 - explambda[k,i] variable, 659–661
 - gain (γ), 650–651
 - hierarchical diagram, 659, 660*f*
 - logistic regression, 653
 - log odds, 651–652
 - posterior parameter estimates, 662–663, 663*f*
 - probability of outcome k , 650, 659
 - property, 654
 - reference category (r), 651
 - reference outcome, 652–653
 - regression coefficients, 652, 663, 664
 - source-ing script, 36
 - Spending per pupil, 514*f*, 524, 527, 530, 538
 - Split-plot design
 - descriptive model, 612–614
 - fertilization method, 611–612, 611*f*
 - interaction contrast, 615, 615*f*
 - JAGS specification, 614
 - main-effect contrasts, 615, 615*f*
 - tilling method, 610, 611–612, 611*f*
 - two-factor between-subject design, 616, 617*f*
 - Standard deviation, 27, 84, 86
 - Structural equation modeling (SEM), 523
 - Student t distribution, 501
 - definition, 458
 - JAGS, 462–464, 463*f*, 464*f*, 465*f*
 - MLEs, 459, 460*f*
 - “normality” parameter, 458–459, 468
 - robust estimation, 461–462
 - “scale” parameter, 458–459
 - Stan codes, 465–468
 - standard deviation, 459–460
 - Subjective belief, 76–77
 - “Substantial” evidence, 268
 - Sum-to-zero deflections, 586, 586*t*
- ## T
- Tail-biased factories, 268–273
 - hierarchical diagram, 269, 269*f*
 - marginal likelihood function, 270
 - posterior distribution, 273
 - posterior probabilities, 270–271
 - prior distribution, 270–271
 - prior probabilities, 271
 - t distribution, 531. *See also* Student t distribution
 - Therapeutic touch
 - data analysis script, 241, 242
 - hierarchical model, 240–241, 241*f*
 - marginal posterior distributions, 243–244, 243*f*
 - marginal prior distributions, 245, 246*f*
 - Three-peaked distribution, 675
 - Thresholded cumulative-normal regression, 685–687, 687*f*
 - Tidal zone, 618

Tilling, 477, 610, 611–612, 611*f*, 615, 615*f*, 616

Traditional ANOVA, 587–588

between-group variance, 556

equality of groups, 556–557

hierarchical Bayesian approach, 557–568

null hypothesis assumption, 556–557

within-group variance, 556

t test, 4, 678*f*, 680–681

Two coin biases

Bernoulli distribution functions, 164, 165

bivariate normal distribution, 168–169

credible differences, 176–177, 177*f*

disadvantage, 170

ESS, 170

vs. formal analysis and grid approximation,
169–170, 169*f*

independent data, 163

infinite and finite random walks, 170

likelihood function, 165

prior beliefs, 163–164

Two-way discrete table

disease diagnosis, 103, 104–105, 104*t*, 106

hair color and eye color proportions, 102, 102*t*

hit rate and false alarm rate, 104–105

joint probability, 101–102, 101*t*, 102–103

low-prior probability, 104

marginal probability, 101–102

posterior probability, 103

Two-way probability distribution

conditional probability, 91–92

independence and, 92–93

joint probability, 90

marginal probability, 90

U

Ulam, Stanislaw, 399

Utility functions

aggregate function, 57, 58

apply function, 59

arguments, 56

formula format, 57

head function, 57

melt command, 59–60

reshape2 package, 59–60

str function, 57

summary function, 56–57

table function, 58

V

Variable selection, 668

a priori, 537

computational methods, 547–548

definition, 536

explanatory model, 544–546, 545*f*

hierarchical shrinkage, 542–544, 545*f*

inclusion probability, 539–542

interaction variables, 548–549

non-negligible posterior probability, 538–539,
539*f*

posterior inclusion probability, 546–547

predicted mean value, 537

regression coefficients, 537–538

Variable types, 42–53

Variance, 86

von Neumann, John, 177

W

Weight, 25, 420, 421, 447, 477, 621, 622,
626–629

Weighted noise, 498–499

Weight prediction, 26, 28–29, 29*f*

WinBUGS, 193–194, 736

Within-subject design

between-subject design, 608–609

carryover effects, 609

definition, 606

observations, 609–610

practice and fatigue effects, 609

“repeated measures”, 606–607

split-plot design, 610–616

two-predictor model, 607

write.csv function, 55