



Early View

State of the art

Understanding the Renin-Angiotensin-Aldosterone-SARS-CoV-Axis: A Comprehensive Review

Nicholas E. Ingraham, Abdo G. Barakat, Ronald Reilkoff, Tamara Bezdicek, Timothy Schacker, Jeffrey G. Chipman, Christopher J. Tignanelli, Michael A. Puskarich

Please cite this article as: Ingraham NE, Barakat AG, Reilkoff R, *et al.* Understanding the Renin-Angiotensin-Aldosterone-SARS-CoV-Axis: A Comprehensive Review. *Eur Respir J* 2020; in press (<https://doi.org/10.1183/13993003.00912-2020>).

This manuscript has recently been accepted for publication in the *European Respiratory Journal*. It is published here in its accepted form prior to copyediting and typesetting by our production team. After these production processes are complete and the authors have approved the resulting proofs, the article will move to the latest issue of the ERJ online.

Copyright ©ERS 2020. This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0.

Understanding the Renin-Angiotensin-Aldosterone-SARS-CoV-Axis: A Comprehensive Review

Nicholas E. Ingraham, MD;¹ Abdo G. Barakat, MD;² Ronald Reilkoff, MD;¹ Tamara Bezdicek, PharmD;³
Timothy Schacker, MD;⁴ Jeffrey G. Chipman, MD;⁵ Christopher J. Tignanelli, MD.^{5,6}; Michael A.
Puskarich MD;^{7,8}

Author Affiliations:

¹ Department of Medicine, University of Minnesota, Division of Pulmonary and Critical Care, Minneapolis, MN

² Department of Anesthesiology, University of Minnesota, Minneapolis, MN

³ Department of Pharmacy, Fairview Pharmacy Services, Minneapolis, MN

⁴ Department of Medicine, University of Minnesota, Division of Medicine and Infectious Disease, Minneapolis, MN

⁵ Department of Surgery, University of Minnesota Division of Acute Care Surgery, Minneapolis, MN

⁶ Institute for Health Informatics, University of Minnesota, Minneapolis, MN

⁷ Department of Emergency Medicine, University of Minnesota, Minneapolis, MN

⁸ Department of Emergency Medicine, Hennepin County Medical Center, Minneapolis, MN

Correspondence:
Nicholas Ingraham, MD

MMC 276
420 Delaware St SE
Minneapolis, MN 55455

Office: (612) 626-1968
Fax: (612) 626-0439
Email: ingra107@umn.edu

NEI ingra107@umn.edu
MAP mike-em@umn.edu
AGB barak041@umn.edu
RR rreilkof@umn.edu
TB tsobask1@fairview.org
TS schac008@umn.edu
JGC chipm001@umn.edu
CTJ ctignane@umn.edu

Conflicts of Interest and Funding Source(s):

1. CJT & MAP: PIs for 2 RCTs investigating ARBs in the treatment of COVID-19 among inpatient and outpatients. CoIs include NEI, TS, JGC.
2. NIH NHLBI T32HL07741 (NEI)

Reprints will not be available from the authors. All authors significantly contributed to developing, writing, and revising this manuscript.

Take Home Message

SARS-CoV-2's interplay with the Renin-Angiotensin-Aldosterone-System likely accounts for much of its unique pathology. Appreciating the degree and mechanism of this interaction highlights potential therapeutic options, including blockade (ARBs).

Abstract

Importance

Coronavirus Disease 19 (COVID-19), the disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has been declared a global pandemic with significant morbidity and mortality since first appearing in Wuhan, China, in late 2019. As many countries are grappling with the onset of their epidemics, pharmacotherapeutics remain lacking. The window of opportunity to mitigate downstream morbidity and mortality is narrow but remains open. The renin-angiotensin-aldosterone system (RAAS) is crucial to the homeostasis of both the cardiovascular and respiratory systems. Importantly, SARS-CoV-2 utilizes and interrupts this pathway directly, which could be described as the renin-angiotensin-aldosterone-SARS-CoV-2-axis (RAAS-SCoV-axis). There exists significant controversy and confusion surrounding how anti-hypertensive agents might function along this pathway. This review explores the current state of knowledge regarding the RAAS-SCoV-axis, informed by prior studies of SARS-CoV, how this relates to our currently evolving pandemic, and how these insights might guide our next steps in an evidence-based manner.

Observations

This review discusses the role of the RAAS-SCoV-axis in acute lung injury and the effects, risks, and benefits of pharmacologic modification of this axis. There may be an opportunity to leverage the different aspects of RAAS inhibitors to mitigate indirect viral-induced lung injury. Concerns have been raised that such modulation might exacerbate the disease. While relevant preclinical, experimental models to date favor a protective effect of RAAS-SCoV-axis inhibition on both lung injury and survival, clinical data related to the role of RAAS modulation in the setting of SARS-CoV-2 remains limited.

Conclusion

Proposed interventions for SARS-CoV-2 predominantly focus on viral microbiology and aim to inhibit viral cellular injury. While these therapies are promising, immediate use may not be

feasible, and the time window of their efficacy remains a major unanswered question. An alternative approach is the modulation of the specific downstream pathophysiologic effects caused by virus that lead to morbidity and mortality. We propose a preponderance of evidence that supports clinical equipoise regarding the efficacy of RAAS-based interventions, and the imminent need for a multisite randomized controlled clinical trial to evaluate the inhibition of the RAAS-SCoV-axis on acute lung injury in COVID-19.

Introduction

COVID-19, the infectious disease caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has left over 180 countries and territories grappling with a devastating pandemic. In December 2019, Wuhan, China, was identified as the epicenter of this outbreak. At the time of this submission, reported COVID-19 cases exceeded 700,000, with more than 30,000 deaths.[1-3] (*At time of article submission*). While early estimates vary and true values remain uncertain, mortality is estimated between 0.4 to 3.4%[4, 5] with initial morbidity and mortality disproportionally affecting older patients.[6] Infectivity (R_0) is estimated at 2.24 to 3.58.[4, 7] COVID-19-related hospitalizations are mostly due to the need for respiratory support and progressively higher levels of care, with respiratory failure being the underlying etiology of COVID-19-related deaths.[8, 9] As many countries are struggling with the onset of their epidemics, pharmacotherapeutics remain lacking.[10, 11] Learning from prior pandemics and related viruses can focus our efforts to control spread and treat those infected.[12]

This novel coronavirus is closely related to the severe acute respiratory syndrome coronavirus (SARS-CoV), which caused an outbreak of disease (SARS) in 2003. Genetic studies found that SARS-CoV-2 shares almost 80% and 50% sequence identity with SARS-CoV and middle eastern respiratory virus coronavirus (MERS-CoV), respectively.[13] The genetic variations between SARS-CoV and SARS-CoV-2 translate into differences in infectivity[14] and immune response[15]. Despite these differences, the shared genome translates into common clinical, microbiologic, and biochemical phenotypes. SARS-CoV-2 mimics SARS-CoV's mechanism of infection which utilizes the

angiotensin-converting enzyme 2 (ACE2), part of the renin-angiotensin-aldosterone system (RAAS). Following binding, ACE2 activity is downregulated through multiple mechanisms, preventing it from performing its usual function in states of health, further summarized below.[13, 16-18] The association between ACE2 and COVID-19 is rooted in two concepts: the mechanism of SARS-CoV-2 infection and the regulatory role of ACE2 during RAAS overactivation. Following the SARS outbreak, extensive research advanced our knowledge of highly morbid coronavirus infections.[16-25] The similar renin-angiotensin-aldosterone-SARS-CoV-2-axis (RAAS-SCoV-axis) interactions shared by the two coronaviruses provide an opportunity to further our understanding of this unique interplay in our pursuit of treatment options.

Early attempts to develop safe and effective vaccines for SARS or MERS have been unsuccessful. It is hypothesized that current efforts will take time to develop, and may or may not be efficacious for this or future coronavirus pandemics.[10, 26-28] Many potential therapies have been proposed with a subset currently undergoing investigation.[29, 30] Meanwhile, extensive efforts to date have been appropriately focused on screening, identification, and containment through the collaboration and coordination between global and local health and governmental agencies.[31] Unfortunately, these public health efforts have had, at best, mixed success curbing the spread of the pandemic.[32] Given the rapid spread of SARS-COV2 and the significant morbidity and mortality associated with infection, it behooves the medical community to evaluate and leverage novel treatments for efficacy, especially if they are already available and have an established safety profile.

RAAS blockade has been proposed as a potential treatment for SARS-CoV-2.[33-36] This hypothesis was initially published by Sun et al.[35, 37] on February 4, 2020, and reinforced in a Drug Development Research publication on March 4, 2020.[33] Other reviews have voiced concern regarding the association between COVID-19 and cardiovascular disease,[38] going too far as to postulate that continued RAAS blockade may cause harm and recommend considering discontinuation.[39] The previous argument is based on the observation that pharmacologic blockers of RAAS can upregulate ACE2 expression, which might increase viral entry into the cell.[39]

Evidence from human subjects to support such an assertion is scant, and as we will see in this review, preclinical and current observational Covid-19 evidence would support the contrary hypothesis – that discontinuation of RAAS blockade may prove harmful. These contrasting hypotheses underscore the dire need to evaluate potential mechanisms, if any, through which RAAS modulation would impact the pathophysiology of COVID-19.[35, 38, 40] In this review, we intend to compile the existing evidence in order to discuss how we might bridge knowledge gaps regarding the interplay between SARS-CoV-2, ACE2, and the RAAS.

RAAS in States of Health

Overview

Renin, angiotensin, and aldosterone represent the core of a complex hormonal axis, referred to as RAAS, which contributes to blood pressure control, sodium reabsorption, inflammation, and fibrosis.[41] RAAS imbalance or modification can cause or treat many diseases including heart failure, hypotension, diabetes, and atherosclerosis, respectively.[42] This review focuses on several physiological and pathological effects of angiotensin II (Ang II) cell signaling. (Figure 1)

Angiotensin II / AT₁ receptor relationship

Angiotensin II (Ang II), the primary physiological product of the RAAS system, is a potent vasoconstrictor. As illustrated in Figure 1, ACE catalyzes the transformation of angiotensin I (Ang I) to Ang II. Ang II elicits its effects by activating two receptors: type 1 angiotensin II (AT₁) receptor and type 2 angiotensin II (AT₂) receptor.[43] Ang II action through AT₁ receptor causes a cascade with resultant inflammation, vasoconstriction, and atherogenesis.[44] These effects also promote insulin resistance and thrombosis.[45] In contrast, AT₂ receptor stimulation causes vasodilation, decreased platelet aggregation, and the promotion of insulin action. However, the expression of AT₂ receptor is low in healthy adults.[45] As such, Ang II's effects in adults are modulated and balanced indirectly by angiotensin II converting enzyme (ACE2), which converts Ang II into lung-protective Angiotensin-(1-7) (Ang-[1-7]), similar to effects seen from AT₂ receptor stimulation.[41, 46] Conceptually

grouping pathways into an Ang II/AT₁ receptor and its opposing pathway ACE2/Ang-(1-7) helps with understanding the countering forces and the sequelae when an imbalance occurs. (Figure 1)

The consequences of excessive Ang II include pulmonary vasoconstriction, inflammatory, and cytokine-induced organ damage[47] secondary to increased membrane permeability,[48] and increased epithelial cell apoptosis.[49] The effects are predominantly mediated through the unopposed AT₁ receptor activation secondary to decreased ACE2 levels.[50] Furthermore, the pro-inflammatory cascade[51] and increased vascular permeability[48] caused by over-activation of the AT₁ receptor in the lungs directly induce acute lung injury (ALI), acute respiratory distress syndrome (ARDS), and lead to death.[10, 33]

Angiotensin-Converting Enzyme 2

ACE2 is expressed in many tissue types, including respiratory epithelial cells.[52] ACE2 is predominantly expressed on the apical surface and converts Ang II into lung-protective Ang-(1-7) and Ang-(1-9) by catalyzing a pathway that prevents unopposed AT₁ receptor activation. Therefore, ACE2 functions as a counterbalance to its structurally similar counterpart ACE[53] through the conversion of Ang II into Ang-(1-7), which in contrast to ACE, promotes vasodilation, reduces proliferation, and prevents apoptosis.[53] ACE2 levels are known to be low in healthy individuals[54] and ACE2 levels decrease with age.[55] Chronically elevated levels of ACE2 are an independent predictor of disease progression; but evidence suggests these changes are compensatory rather than causal.[54]

ACEis and ARBs

Both angiotensin-converting enzyme inhibitors (ACEis) and angiotensin receptor blockers (ARBs) increase tissue and plasma levels of renin and Ang I.(50) ACEis inhibit Ang I to Ang II conversion, decreasing serum Ang II levels to baseline and have little to no affinity for ACE2.[56] At clinical doses, ACEis only partially affect this conversion because 40% of Ang II formation occurs outside the ACE pathway.[57] ACE levels are in turn decreased through a negative feedback in states

of high Ang II levels.[58] Ang II levels increase in response to ARBs at all levels, except in the kidney. The consequence of this increase is a redirection of Ang II activity through the anti-inflammatory and vasodilatory ACE2 and AT₂ receptor pathways.[56] As a result, ACE2, compared to ACE, regulates local levels of both Ang II and Ang-(1-7) to a greater degree.[42] Tikellis et al. suggest that ARBs, therefore, have a higher potential to favorably rebalance the RAAS pathway compared to ACEi.[42] Furthermore, the kallikrein-kinin system is regulated by ACE and ACE2. Increased activity of the kallikrein-kinin system occurs under proinflammatory conditions,[59] specifically bradykinin and des-Arg-BK⁹, which can lead to vascular leakage when unchecked by ACE and ACE2, respectively. This mechanism has been posited to be, at least in part, a cause of cough[60] and pulmonary edema,[61] independent of proposed Ang II induced hydrostatic pressure. The dysregulated kallikrein-kinin system's role remains theoretical in COVID-19; however, if true, AT₁ receptor blockade may be favored (over ACE inhibition) in an attempt to preserve the regulatory mechanisms of this system.[61] While there may still be a key role for ACE inhibition in the context of the current disease that we will elucidate in the following sections, these observations support our decision to narrow the focus of this review to the downstream RAAS modifiers, ARBs.

- ACE2 levels are high in diseased states, likely secondary to an insufficient compensatory response to overactive RAAS activity
- In COVID-19, high rates of pulmonary edema and cough may be due, in part, to reduced breakdown of bradykinin from decreased ACE activity
- ARBs may provide crucial regulation to the overactive RAAS-SCoV-axis

Key Points 1

RAAS in Cardiovascular Disease

Modulation of the RAAS axis is central to the management of cardiovascular disease.[62] Cardiovascular disease, heart failure, atrial fibrillation, and kidney disease are associated with elevated levels of ACE2.[60] It is critical to discern if these elevations are a culprit or sequelae of

disease in order to separate association from causation.[63] More recent opinions suggest ACE2 represents as a biomarker, rather than a culprit, of cardiovascular disease.[64]

Ramchand et al. demonstrated elevated ACE2 levels to be an independent risk factor of major adverse cardiac events in patients with known coronary artery disease, though this data, as noted by the authors, does not differentiate association versus causation.[54] To begin to dissect this interplay, we turn to preclinical models, specifically ACE2 knockout mice that are characterized by severe cardiac defects. Critically, these defects are notably absent in double ACE and ACE2 knockout models which highlights the highly balanced interplay between ACE/Ang II/AT₁ and ACE2/Ang-(1-7).[53, 65] Similar protective effects to prevent cardiovascular events and strokes have been demonstrated in mouse models with overexpressed ACE2.[66] Rachmand et al. concluded that patients with elevated ACE2 might “reflect a persistent albeit insufficient counter-regulatory process to shift the balance away from the deleterious effects of sustained Ang II activation.”[54] As such, ACE2 overexpression may be a *compensatory mechanism* to mitigate the effect of unopposed Ang II stimulation though increased degradation into protective Ang-(1-7).[67]

It is important to note that preclinical animal model treatment with losartan upregulates cardiac ACE2 mRNA expression and increases in ACE2 activity. [68, 69] However, human studies found no significant difference in ACE2 plasma levels in patients treated with ACEi or ARBs have not confirmed such an effect, though local tissue ACE2 levels were not measured.[54]. Through anecdotal reports, COVID-19 may be associated with high rates of myocarditis and heart failure in the critically ill. While beyond the scope of this proposal, we will note that increased Ang II and decreased ACE2 are also associated with impaired cardiac contractility, which can be rescued with losartan.[70] More observational data is needed to assess if other signs of RAAS overdrive (hypokalemia, hypertension) portend poor outcomes and may be associated with arrhythmias that have been noted in the diseases.[71] Nevertheless, this could potentially underscore potential benefits of RAAS modulation that extend beyond the attenuation of lung injury.

- ACEis and ARBs interrupt maladaptive pathophysiologic responses to heightened activation of the RAAS.
- ARBs and ACEis mitigate the deleterious effects of unopposed AT₁ receptor pathway activation, which in turn decreases inflammation, insulin resistance, and lung injury

Key Points 2

RAAS in Pulmonary Disease

Chronic Lung Disease

In chronic obstructive lung disease (COPD), patients treated with ARBs (compared to those on ACEi) had less severe exacerbations, fewer exacerbations overall, lower mortality, lower mechanical ventilation requirements, and fewer hospitalizations.[72] Moreover, veterans over 65 years old that were on ARBs before and during their hospitalization for pneumonia had decreased mortality when compared to patients without such treatment.[73]

Acute Lung Injury

Unopposed RAAS activation via the Ang II/AT₁ receptor causes inflammation[51] increased vascular permeability[48] and severe lung injury,[10, 33] while ARBs significantly attenuate these changes.[48-51] Importantly, the mere presence of high concentrations of Ang II can further regulate the expression of ACE2, leading to dysregulated Ang II/AT₁ receptor activity.[74] In mice, losartan reduced mortality by blunting Ang II-associated increases in soluble epoxide hydrolase, a promoter of lung injury.[75] Animal models of ventilator-associated lung injury have demonstrated benefit from losartan, mitigating Ang II activity and AT₁ receptor expression.[76-78] While most studies include pre-treated animal models, rescue models also demonstrate efficacy, with the restoration of ACE2 levels, blunting of PaO₂ decline, and attenuation of lung injury.[50]

In human patients, genetic cohort studies yield further insights into the relationship between RAAS and acute lung injury. Jerng et al. found that polymorphisms in the ACE gene are

associated with outcomes in ARDS. These findings were corroborated by Adamzik et al., who identified patients with the ACE DD genotype (associated with increased ACE activity) to have the highest risk of ARDS-related death (HR 5.7).[79] Other human studies evaluating the association between RAAS-inhibition and ARDS remain observational. Kim et al. found ARDS patients that were taking ACEi or ARBs had better survival rates when compared to those without RAAS inhibition.[80] A secondary analysis of a 2010 randomized control trial in patients with acute respiratory failure suggested that treatment with ACEi/ARB at discharge following an episode of acute respiratory failure was associated with a 44% reduction in 1-year mortality.[81] More recently, Hsieh et al. observed lower adjusted odds of hospital mortality in patients with sepsis (with and without shock) who were on ARB or ACEi therapy.[82] Mortenson et al. also found a 58% decrease in the odds of hospital mortality in patients taking ARBs before admission.[83] Based on the above, there have been calls to further elucidate the potential benefits of ACEi and ARB in ARDS.[84] However, randomized control trials around this topic were not identified in the peer-reviewed literature or clinicaltrials.gov at the time of this manuscript's preparation.

Pneumonia

While influenza and other types of pneumonia may interact with the RAAS axis, both animal and human studies illustrate clear indirect effects on the RAAS in the setting of certain influenza strains in particular. Previous studies suggest Ang II levels predict mortality in undifferentiated patients with influenza,[85] and continuation of RAAS inhibitor therapy during admission is associated with decreased hospital mortality and odds of intubation in viral cases of pneumonia.[86] RAAS may have implications for other viral pneumonias, as well, as Gu et al. found that children with respiratory syncytial virus tend to have higher Ang II levels compared to healthy children.[87] Based on this observation, they demonstrated the benefit of recombinant ACE2 therapy on RSV infection in a preclinical mouse model.

Critically, both H7N9 and H5N1 influenza have been shown to cause lung injury through ACE2 downregulation, upregulation of Ang II, and AT₁ receptor-induced lung injury.[88, 89] Mouse models of H5N1 and H7N9 demonstrated decreased IL-6, lung edema, lung injury, and mortality if treated with losartan.[89, 90] The mechanism by which losartan prevents lung injury may not reside solely within the RAAS pathway, however.[51] Liu et al. suggest that losartan inhibits the activation of pulmonary dendritic cells.[91] In a study of rats with *pseudomonas* pneumonia, AT₁ receptor blockers suppressed the activation of neutrophils through a mechanism that does not involve the AT₁ receptor down-regulation.[51, 92] Such studies have raised concern that losartan treatment might decrease microbial clearance. In sharp contrast, study results actually demonstrate reduced viral loads[90] and increased bacterial clearance[51] in lung injury models treated with losartan. Given the complexity of these interactions, future investigations remain necessary to further elucidate these relationships.

ACE2 binding viral infection SARS-CoV

In 2003, ACE2 was identified as the binding protein for SARS-CoV. In the years to follow, studies demonstrating the downstream consequences of downregulation of ACE2 with resultant increases in Ang II concentration illustrate the vicious cycle caused by the disruption of the RAAS, we refer to in this manuscript as RAAS-SARS-CoV-2 (RAAS-SCoV-axis).[18] Of relevance to the current pandemic, Wrapp et al. found SARS-CoV-2 to have a 10- to 20-fold higher affinity for ACE2 than SARS-CoV. The authors postulate this may cause increased infectivity, which may help explain the differences in the evolution of the two epidemics.

Both SARS-coronaviruses contain four structural proteins, the nucleocapsid, membrane, envelope, and spike protein.[93] The nucleocapsid, membrane, and envelope proteins have roles in viral replication, structural integrity, and host response among other mechanisms.[93] The spike protein, known as the S-protein, is integral to attaching and infecting host cells.[14, 93] Phan et al. identified specific deletions that increase the potential of the SARS-CoV-2 spike to evade immune responses over time. These changes have implications for both initial and subsequent infection

severity, immunity, and the likelihood of developing an efficacious vaccine. More relevant to the current review, however, is the potential effect of such changes might have on the RAAS-SCoV-axis, which could serve to explain not only their predilection for causing respiratory failure but possibly other commonalities between these infections.

SARS-CoV infection downregulates the surface expression of the binding protein, ACE2, a pivotal component for host cell entry.[17] Notably, low ACE2 expression is associated with increased phenotype severity in *in-vitro* human airway epithelial cell studies.[24] Imai et al. validated these findings in a whole animal model by demonstrating that loss of ACE2 in knockout (KO) mice leads to impairments in oxygenation, increased inflammation, and worsening of edema compared to WT mice after acid-induced lung injury.[23] Pre-treatment with AT₁ receptor inhibitors before acid induced ARDS showed significantly lower lung injury, confirming the AT₁ receptor's role in the physiologic response.[23] Subsequently and most critically, Kuba et al. demonstrated SARS-CoV spike protein augments Ang II increase and ACE2 downregulation, with resultant lung injury.[17] In this model, losartan can attenuate the severity of acute lung injury due to SARS-CoV infection in both a pre-treatment model and a more clinically relevant post-infection (rescue) model.[17]

Ensuing studies have confirmed and built upon these findings.[19] AT₁ receptor blockade upregulates ACE2 through a negative feedback mechanism, and serve as a lung-protective mechanism through the increased conversion of Ang II to Ang-(1-7), effectively blunting pulmonary injury from the virus.[50] These findings have led investigators to posit that the ACE2/Ang-(1-7) pathway may serve as a therapeutic target to combat the pathological effects of Ang II.[74] To that end, there is already at least one trial of recombinant human ACE2 therapy for the treatment of COVID-19 that attempts to leverage this relationship and mitigate downstream lung injury.[23, 35]

The mechanism by which SARS-CoV downregulates ACE2 may be relevant to the development of novel therapeutics, as well. Haga et al. demonstrated that SARS-CoV binding to ACE2 induces ACE2 shedding as a soluble form into the serum,[20] and further studies validated

those findings.[19] In addition, endocytosis of the ACE2 protein occurs after binding to SARS-CoV, further decreasing ACE2 activity,[94] with potential implications for Ang II/Ang1-7 balance described throughout this review. A final critical mechanism that cannot be overlooked includes the role of AT₁ receptor in the downregulation of ACE2. Desholtz et al. found that ACE2 downregulation induced persistent elevation of Ang II through local interaction with the AT₁ receptor, which triggers a vicious cycle where Ang II downregulates ACE2 leading to further increases in local tissue levels of Ang II.[74] These mechanisms promote an unopposed Ang II/AT₁ receptor axis. (Figure 1)

The ACE2-SARS/COVID-19 interplay has led to controversy with clinical implications. The finding that ACE2 KO mice are protected against SARS-CoV infection[17] raises a concern that an upregulated ACE2 is harmful in the setting of SARS-CoV and COVID infections, as the virus cannot enter the cell. Conversely, ACE2 downregulation following infection is concerning considering the preclinical data summarized to this point, which instead suggests a putative causative mechanism of Ang II/Ang (1-7) in the evolution of SARS-CoV-2 associated acute lung injury. Even if abolishing ACE2 were feasible, Han et al. detected alternative receptors, DC/L-SIGN, which mediated SARS-CoV entry indep

ende
ntly
of
ACE2.
[21]

- SARS-CoV decreases surface expression of ACE2 during infection
- Decreased ACE2 activity leads to increased Ang II and further downregulation of ACE2 in a vicious cycle, driving acute lung injury.
- While the main point of entry involves ACE2, other receptors can independently mediate SARS-CoV infection

Key Points 3

Controversies regarding the causative role of ACE2 in COVID-19

Limited mechanistic investigations regarding COVID-19 currently exist.[95] Early Chinese data illustrate a progressive increase in Ang-II levels among patients hospitalized in the setting of confirmed COVID-19,[96] along with concomitant increasing levels of IL-6 levels, with the highest levels in non-survivors.[97] Similarly, a 2014 analysis of H7N9 patients showed progressively

increasing levels of Ang-II, up to 4 weeks following infection, a finding associated with a worse prognosis.[89] In a recent COVID-19 patient study, quantified levels of Ang-II were positively correlated with viral titers and closely associated with PaO₂/FiO₂ measures, suggesting a relationship between infection, RAAS, and lung injury. Accordingly, the study investigators postulated that angiotensin receptor blockade might represent a promising therapeutic target.[96]

As noted previously, ACE2 levels are known to be low in healthy individuals,[54] decrease with age, and are lower in men than in women.[55, 98] Age is an independent risk factor for mortality in patients admitted with COVID-19.[97] The data also suggests a disproportionate rate of disease occurrence, severity, and worse outcomes in males compared to females during this pandemic.[99] These associations might be interpreted to indicate that low ACE2 during times of health are appropriate, but the ability to regulate overactive RAAS or RAAS-SCoV-axis during illness is, at least in part, dependent on ACE2. For instance, one study identified a higher expression of the lung-protective AT₂ receptor among women, which could explain the disproportionately lower mortality in female COVID-19 subjects.[100] Interestingly, ACE2 is located on the X-chromosome, which has been hypothesized to affect the ability of men to maintain adequate ACE2 production in situations of severe illness (i.e., ARDS or COVID-19)[40]

Perhaps most importantly, the most common comorbidities of severely ill patients included hypertension, diabetes mellitus, and coronary heart diseases.[9] These comorbidities are associated with either an overactivation of Ang II/AT₁ or a deficiency of ACE2,[42, 45] and all have shown benefit from RAAS blockade in chronic conditions,[101] including an attenuation in AT₁ receptor expression with losartan.[102] Furthermore, smoking tends to increase AT₁ receptor expression[102] along with ACE2 expression in human and mouse epithelial cells.[103, 104] RAAS activation in states of stress with high circulating Ang II levels may be particularly harmful for such patients, and could explain at least some of the observed clinical outcomes. Namely, data from China found the odds of disease progression in those with COVID-19 are 14 times higher in smokers.[8] However, we

acknowledge that misinterpreting association for causation can be dangerous and that the pathology underlying acute and chronic conditions may not be equivalent.[105] Nevertheless, RAAS manipulation remains an attractive therapeutic target not to treat or prevent infection with COVID-19, but rather mitigate its downstream consequences.

Side effects of RAAS manipulation are understandably a cause for concern, specifically hypotension and acute kidney injury in the setting of acute illness. The most severe risks of ARBs include teratogenicity in the setting of pregnancy, hypersensitivity including angioedema, symptomatic hypotension, worsening of renal function, and electrolyte abnormalities; however, the most common side effects of include fatigue, weakness, diarrhea, chest pain and anemia. Clinical trial and post-marketing surveillance data of losartan demonstrate an excellent safety profile. In over 4,000 patients, there was a low incidence of adverse events (2.1%) comparable to placebo (3.7%).[106] The effects more common in losartan than placebo were clinically minor, including dizziness (3% vs 2%), upper respiratory infection (8% vs 7%), nasal congestion (2% vs 1%) and back pain (2% vs 1%). While losartan has the potential to cause symptomatic hypotension, four studies have identified losartan has a negligible effect on resting blood pressure and heart rate in healthy volunteers.[107-110] These reassuring data cannot be extrapolated to COVID-19 patients; however, when assessing the risk/benefit balance for RAAS modulation, the baseline risk appears low in the healthy population whereas repurposing of other medications may not have the same favorable profile. There are also concerns that the upregulation of ACE2 by ARBs could accelerate SARS-CoV-2 entry in host cells. Such concerns become more relevant when one considers the observation that patients with hypertension tend to have worse outcomes, and many patients with hypertension are on agents that might upregulate ACE2 expression. Consequently, some clinicians have argued for discontinuing ACEi and ARBs and switching their patients to different anti-hypertensive classes. However, to our knowledge, there is no data to suggest such a change is beneficial, nor are there well controlled, risk adjusted analyses demonstrating that these medications are harmful.

On the contrary, data summarized in this review actually suggests that discontinuation of ACEi or ARBs could pose a serious risk of harm through the propagation of excessive Ang II mediated acute lung injury and downregulation of protective ACE2. To this point, in a recent preprint, Liu et al retrospectively studied 511 COVID-19 patients (age > 65 years old) with hypertension. Patients were categorized based on home anti-hypertensive regimen. Patients on ARBs had significantly decreased odds of developing severe COVID-19 disease on univariate analysis (OR 0.34, $p = 0.025$) and multivariable analysis (OR 0.25, $p = 0.046$), while none of the other medications demonstrated improved outcomes. Two subsequent retrospective studies have demonstrated a protective effect of RAS blockade, including significant reductions in viral load.[111, 112] Finally, in addition to the inhibitory effects on AT1R, a pre-print publication from Iran[113] identified losartan as a potential candidate treatment based on molecular docking and dynamic simulation. In addition to the effects on angiotensin homeostasis, therefore, losartan may directly inhibit the viral macrodomain and cell cycle.[113] Even prior to these observational data, based on pre-clinical evidence and non-COVID-19 studies, the International Society of Hypertension, the European Society of Cardiology, and other international committees have strongly recommended that physicians and patients continue treatment with their usual anti-hypertensive therapy as there is no clinical or scientific evidence to suggest that treatment with ACEi or ARBs should be discontinued in the setting of the SARS-CoV-2 pandemic at the current time.[114-116]

Conclusion

Current clinical evidence is insufficient to recommend treating with or withholding RAAS blockade during this pandemic. Despite elevated ACE2 levels being found in chronic disease states, the upregulation of ACE2 seems more likely an adaptive response rather than a culprit.[67] Significant preclinical data and retrospective human data suggest RAAS inhibition decreases lung injury and improves survival, while simultaneously decreasing viral load in animal models with viral infections that utilize the ACE2 receptor. Human clinical data regarding the effects of RAAS inhibitors on outcomes remain modest and an area of active investigation. While the association between

hypertension and poor outcomes might be mediated through ACE2 upregulation and increasing viral susceptibility, these differential outcomes are at least as likely to be a result of dysregulated Ang II-mediated lung injury. We believe this represents a state of clinical equipoise and hypothesize that RAAS inhibition may improve outcomes in patients infected with COVID-19. Furthermore, the wide spectrum of severity during the pandemic highlights the heterogeneity in the patients with COVID-19. As different phenotypes emerge, certain subgroups (i.e. those with overactive RAAS) may differ in their response to therapy. Time will also need to be considered. RAAS modulation may have significantly different effects early in the disease process compared to later given the complexity of the RAAS-SCoV-axis. This underscores the exigency for high-quality evidence to investigate any significant association between RAAS modulation and pulmonary pathophysiology, with a focus on ACE2-mediated viral infections, particularly, SARS-CoV-2.

Acknowledgments:

We gratefully acknowledge the editorial assistance of Michael Lotti, Senior Editor of the University of Minnesota's Institute for Engineering in Medicine.

References

References

1. Dong E, Du H, Gardner L: **An interactive web-based dashboard to track COVID-19 in real time.** *Lancet Infect Dis* 2020.
2. **Coronavirus COVID-19 (2019-nCoV)**
[<https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6>]
3. **Novel Coronavirus (2019-nCoV) situation reports**
[<https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>]
4. Wang Y, Wang Y, Chen Y, Qin Q: **Unique epidemiological and clinical features of the emerging 2019 novel coronavirus pneumonia (COVID-19) implicate special control measures.** *J Med Virol* 2020.
5. Rajgor DD, Lee MH, Archuleta S, Bagdasarian N, Quek SC: **The many estimates of the COVID-19 case fatality rate.** *Lancet Infect Dis* 2020.
6. World Health O: **Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19).** In.; 2020.
7. Zhao S, Lin Q, Ran J, Musa SS, Yang G, Wang W, Lou Y, Gao D, Yang L, He D *et al*: **Preliminary estimation of the basic reproduction number of novel coronavirus (2019-nCoV) in China, from 2019 to 2020: A data-driven analysis in the early phase of the outbreak.** *Int J Infect Dis* 2020, **92**:214-217.
8. Liu W, Tao ZW, Lei W, Ming-Li Y, Kui L, Ling Z, Shuang W, Yan D, Jing L, Liu HG *et al*: **Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease.** *Chin Med J (Engl)* 2020.

9. Guan WJ, Ni ZY, Hu Y, Liang WH, Ou CQ, He JX, Liu L, Shan H, Lei CL, Hui DSC *et al*: **Clinical Characteristics of Coronavirus Disease 2019 in China.** *The New England journal of medicine* 2020.
10. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS: **Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target.** *Intensive Care Med* 2020, **46**(4):586-590.
11. Ingraham NE, Tignanelli CJ: **Fact Versus Science Fiction.** *Crit Care Explor* 2020, **2**(4):e0108.
12. Kuba K, Imai Y, Rao S, Jiang C, Penninger JM: **Lessons from SARS: control of acute lung failure by the SARS receptor ACE2.** *J Mol Med (Berl)* 2006, **84**(10):814-820.
13. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL *et al*: **A pneumonia outbreak associated with a new coronavirus of probable bat origin.** *Nature* 2020, **579**(7798):270-273.
14. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS, McLellan JS: **Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.** *Science* 2020, **367**(6483):1260-1263.
15. Phan T: **Genetic diversity and evolution of SARS-CoV-2.** *Infect Genet Evol* 2020, **81**:104260.
16. Dijkman R, Jebbink MF, Deijns M, Milewska A, Pyrc K, Buelow E, van der Bijl A, van der Hoek L: **Replication-dependent downregulation of cellular angiotensin-converting enzyme 2 protein expression by human coronavirus NL63.** *J Gen Virol* 2012, **93**(Pt 9):1924-1929.
17. Kuba K, Imai Y, Rao S, Gao H, Guo F, Guan B, Huan Y, Yang P, Zhang Y, Deng W *et al*: **A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury.** *Nat Med* 2005, **11**(8):875-879.
18. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC *et al*: **Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus.** *Nature* 2003, **426**(6965):450-454.
19. Glowacka I, Bertram S, Herzog P, Pfefferle S, Steffen I, Muench MO, Simmons G, Hofmann H, Kuri T, Weber F *et al*: **Differential downregulation of ACE2 by the spike proteins of severe acute respiratory syndrome coronavirus and human coronavirus NL63.** *J Virol* 2010, **84**(2):1198-1205.
20. Haga S, Yamamoto N, Nakai-Murakami C, Osawa Y, Tokunaga K, Sata T, Yamamoto N, Sasazuki T, Ishizaka Y: **Modulation of TNF- α -converting enzyme by the spike protein of SARS-CoV and ACE2 induces TNF- α production and facilitates viral entry.** *Proceedings of the National Academy of Sciences* 2008, **105**(22):7809-7814.
21. Han DP, Lohani M, Cho MW: **Specific asparagine-linked glycosylation sites are critical for DC-SIGN- and L-SIGN-mediated severe acute respiratory syndrome coronavirus entry.** *J Virol* 2007, **81**(21):12029-12039.
22. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu N-H, Nitsche A: **SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor.** *Cell* 2020.
23. Imai Y, Kuba K, Rao S, Huan Y, Guo F, Guan B, Yang P, Sarao R, Wada T, Leong-Poi H *et al*: **Angiotensin-converting enzyme 2 protects from severe acute lung failure.** *Nature* 2005, **436**(7047):112-116.
24. Jia HP, Look DC, Shi L, Hickey M, Pewe L, Netland J, Farzan M, Wohlford-Lenane C, Perlman S, McCray PB, Jr.: **ACE2 receptor expression and severe acute respiratory syndrome coronavirus infection depend on differentiation of human airway epithelia.** *J Virol* 2005, **79**(23):14614-14621.
25. Jia HP, Look DC, Tan P, Shi L, Hickey M, Gakhar L, Chappell MC, Wohlford-Lenane C, McCray PB, Jr.: **Ectodomain shedding of angiotensin converting enzyme 2 in human airway epithelia.** *Am J Physiol Lung Cell Mol Physiol* 2009, **297**(1):L84-96.
26. CDC: **From the Centers for Disease Control and Prevention. Severe Acute Respiratory Syndrome (SARS) and coronavirus testing--United States, 2003.** *JAMA* 2003, **289**(17):2203-2206.

27. de Wit E, van Doremalen N, Falzarano D, Munster VJ: **SARS and MERS: recent insights into emerging coronaviruses.** *Nature reviews Microbiology* 2016, **14**(8):523-534.
28. Song Z, Xu Y, Bao L, Zhang L, Yu P, Qu Y, Zhu H, Zhao W, Han Y, Qin C: **From SARS to MERS, Thrusting Coronaviruses into the Spotlight.** *Viruses* 2019, **11**(1):59.
29. Zhang L, Liu Y: **Potential interventions for novel coronavirus in China: A systematic review.** *J Med Virol* 2020, **92**(5):479-490.
30. Li H, Zhou Y, Zhang M, Wang H, Zhao Q, Liu J: **Updated approaches against SARS-CoV-2.** *Antimicrob Agents Chemother* 2020:AAC.00483-00420.
31. Fang Y, Nie Y, Penny M: **Transmission dynamics of the COVID-19 outbreak and effectiveness of government interventions: A data-driven analysis.** *J Med Virol* 2020.
32. Wilder-Smith A, Freedman DO: **Isolation, quarantine, social distancing and community containment: pivotal role for old-style public health measures in the novel coronavirus (2019-nCoV) outbreak.** *J Travel Med* 2020, **27**(2).
33. Gurwitz D: **Angiotensin receptor blockers as tentative SARS-CoV-2 therapeutics.** *Drug Dev Res* 2020.
34. Henry B: **COVID-19 induced Renin–Angiotensin System (RAS) imbalance may drive acute lung injury: the evidence and therapeutic options.** 2020.
35. Sun ML, Yang JM, Sun YP, Su GH: **[Inhibitors of RAS Might Be a Good Choice for the Therapy of COVID-19 Pneumonia].** *Zhonghua Jie He He Hu Xi Za Zhi* 2020, **43**(0):E014.
36. Tignanelli CJ, Ingraham NE, Sparks MA, Reilkoff R, Bezdicek T, Benson B, Schacker T, Chipman JG, Puskas MA: **Antihypertensive drugs and risk of COVID-19?** *Lancet Respir Med* 2020.
37. **BMJ 2020;368:m406.**
38. Zheng YY, Ma YT, Zhang JY, Xie X: **COVID-19 and the cardiovascular system.** *Nat Rev Cardiol* 2020.
39. Sommerstein R: **Rapid Response to Preventing a covid-19 pandemic: ACE inhibitors as a potential risk factor for fatal Covid-19.** 2020.
40. Garami AR: **Rapid response to Preventing a covid-19 pandemic: Is there a magic bullet to save COVID-19 Patients? We can give it a try!** In.: British Medical Journal Publishing Group; 2020.
41. Ghazi L, Drawz P: **Advances in understanding the renin-angiotensin-aldosterone system (RAAS) in blood pressure control and recent pivotal trials of RAAS blockade in heart failure and diabetic nephropathy.** *F1000Research* 2017, **6**:F1000 Faculty Rev-1297.
42. Tikellis C, Thomas MC: **Angiotensin-Converting Enzyme 2 (ACE2) Is a Key Modulator of the Renin Angiotensin System in Health and Disease.** *Int J Pept* 2012, **2012**:256294.
43. Balakumar P, Jagadeesh G: **A century old renin–angiotensin system still grows with endless possibilities: AT1 receptor signaling cascades in cardiovascular physiopathology.** *Cellular signalling* 2014, **26**(10):2147-2160.
44. Strawn WB, Ferrario CM: **Mechanisms linking angiotensin II and atherogenesis.** *Curr Opin Lipidol* 2002, **13**(5):505-512.
45. Dandona P, Dhindsa S, Ghanim H, Chaudhuri A: **Angiotensin II and inflammation: the effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockade.** *J Hum Hypertens* 2007, **21**(1):20-27.
46. Warner FJ, Smith AI, Hooper NM, Turner AJ: **Angiotensin-converting enzyme-2: a molecular and cellular perspective.** *Cell Mol Life Sci* 2004, **61**(21):2704-2713.
47. Zhang H, Baker A: **Recombinant human ACE2: acing out angiotensin II in ARDS therapy.** In.: Springer; 2017.
48. Chen M, Chen C, Yuan X, Chen X, Zheng F, Shao L, Zhang Z: **Angiotensin II aggravates lipopolysaccharide induced human pulmonary microvascular endothelial cells permeability in high glucose status.** *Endocr J* 2018, **65**(7):717-725.
49. Filippatos G, Uhal BD: **Blockade of apoptosis by ACE inhibitors and angiotensin receptor antagonists.** *Curr Pharm Des* 2003, **9**(9):707-714.
50. Wosten-van Asperen RM, Lutter R, Specht PA, Moll GN, van Woensel JB, van der Loos CM, van Goor H, Kamilic J, Florquin S, Bos AP: **Acute respiratory distress syndrome leads to**

- reduced ratio of ACE/ACE2 activities and is prevented by angiotensin-(1-7) or an angiotensin II receptor antagonist.** *J Pathol* 2011, **225**(4):618-627.
51. Raiden S, Nahmod K, Nahmod V, Semeniuk G, Pereira Y, Alvarez C, Giordano M, Geffner JR: **Nonpeptide antagonists of AT1 receptor for angiotensin II delay the onset of acute respiratory distress syndrome.** *J Pharmacol Exp Ther* 2002, **303**(1):45-51.
 52. Mossel EC, Wang J, Jeffers S, Edeen KE, Wang S, Cosgrove GP, Funk CJ, Manzer R, Miura TA, Pearson LD *et al*: **SARS-CoV replicates in primary human alveolar type II cell cultures but not in type I-like cells.** *Virology* 2008, **372**(1):127-135.
 53. Turner AJ, Hiscox JA, Hooper NM: **ACE2: from vasopeptidase to SARS virus receptor.** *Trends in pharmacological sciences* 2004, **25**(6):291-294.
 54. Ramchand J, Patel SK, Srivastava PM, Farouque O, Burrell LM: **Elevated plasma angiotensin converting enzyme 2 activity is an independent predictor of major adverse cardiac events in patients with obstructive coronary artery disease.** *PLoS One* 2018, **13**(6):e0198144.
 55. AlGhatrif M, Cingolani O, Lakatta EG: **The Dilemma of Coronavirus Disease 2019, Aging, and Cardiovascular Disease: Insights From Cardiovascular Aging Science.** *JAMA Cardiol* 2020.
 56. Campbell DJ: **Endogenous angiotensin II levels and the mechanism of action of angiotensin-converting enzyme inhibitors and angiotensin receptor type 1 antagonists.** *Clin Exp Pharmacol Physiol Suppl* 1996, **3**:S125-131.
 57. Hollenberg NK, Fisher ND, Price DA: **Pathways for angiotensin II generation in intact human tissue: evidence from comparative pharmacological interruption of the renin system.** *Hypertension* 1998, **32**(3):387-392.
 58. Schunkert H, Ingelfinger JR, Hirsch AT, Pinto Y, Remme WJ, Jacob H, Dzau VJ: **Feedback regulation of angiotensin converting enzyme activity and mRNA levels by angiotensin II.** *Circ Res* 1993, **72**(2):312-318.
 59. Jurado- Palomo J, Caballero T: **Pathophysiology of Bradykinin-Mediated Angioedema: The Role of the Complement System.** *A Comprehensive Review of Urticaria and Angioedema Selda Pelin Kartal & Zekayi Kutlubay (Eds) Intech Open Access Publisher (Rijeka-Croatia)* 2017:151-176.
 60. Bahat G: **Angiotensin converting enzyme (ACE) inhibition may have a role in the symptoms and progression of COVID-19 infection.** In.; 2020.
 61. van de Veerdonk F, Netea MG, van Deuren M, van der Meer JWM, de Mast Q, Bruggemann RJ, van der Hoeven H: **Kinins and Cytokines in COVID-19: A Comprehensive Pathophysiological Approach (Preprint).** *Preprints* 2020, **2020040023** (doi: **10.20944/preprints202004.0023.v1**).
 62. Braun-Menendez E, Fasciolo JC, Leloir LF, Munoz JM: **The substance causing renal hypertension.** *J Physiol* 1940, **98**(3):283-298.
 63. Walters TE, Kalman JM, Patel SK, Mearns M, Velkoska E, Burrell LM: **Angiotensin converting enzyme 2 activity and human atrial fibrillation: increased plasma angiotensin converting enzyme 2 activity is associated with atrial fibrillation and more advanced left atrial structural remodelling.** *Europace* 2017, **19**(8):1280-1287.
 64. Turner AJ: **ACE2 Cell Biology, Regulation, and Physiological Functions.** In: *The Protective Arm of the Renin Angiotensin System (RAS)*. Edited by Unger T, Steckelings UM, dos Santos RAS. Boston: Academic Press; 2015: 185-189.
 65. Crackower MA, Sarao R, Oudit GY, Yagil C, Kozieradzki I, Scanga SE, Oliveira-dos-Santos AJ, da Costa J, Zhang L, Pei Y: **Angiotensin-converting enzyme 2 is an essential regulator of heart function.** *Nature* 2002, **417**(6891):822-828.
 66. Rentzsch B, Todiras M, Iliescu R, Popova E, Campos LA, Oliveira ML, Baltatu OC, Santos RA, Bader M: **Transgenic angiotensin-converting enzyme 2 overexpression in vessels of SHRSP rats reduces blood pressure and improves endothelial function.** *Hypertension* 2008, **52**(5):967-973.
 67. Turner AJ: **Angiotensin-converting enzyme 2: cardioprotective player in the renin-angiotensin system?** In.: Am Heart Assoc; 2008.

68. Ferrario CM, Jessup J, Chappell MC, Averill DB, Brosnihan KB, Tallant EA, Diz DI, Gallagher PE: **Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2.** *Circulation* 2005, **111**(20):2605-2610.
69. Ocaranza MP, Godoy I, Jalil JE, Varas M, Collantes P, Pinto M, Roman M, Ramirez C, Copaja M, Diaz-Araya G *et al*: **Enalapril attenuates downregulation of Angiotensin-converting enzyme 2 in the late phase of ventricular dysfunction in myocardial infarcted rat.** *Hypertension* 2006, **48**(4):572-578.
70. Perlot T, Penninger JM: **ACE2 - from the renin-angiotensin system to gut microbiota and malnutrition.** *Microbes Infect* 2013, **15**(13):866-873.
71. chen d, Li X, song q, Hu C, Su F, Dai J: **Hypokalemia and Clinical Implications in Patients with Coronavirus Disease 2019 (COVID-19).** *medRxiv* 2020:2020.2002.2027.20028530.
72. Lai CC, Wang YH, Wang CY, Wang HC, Yu CJ, Chen L: **Comparative effects of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers on the risk of pneumonia and severe exacerbations in patients with COPD.** *Int J Chron Obstruct Pulmon Dis* 2018, **13**:867-874.
73. Mortensen EM, Nakashima B, Cornell J, Copeland LA, Pugh MJ, Anzueto A, Good C, Restrepo MI, Downs JR, Frei CR *et al*: **Population-based study of statins, angiotensin II receptor blockers, and angiotensin-converting enzyme inhibitors on pneumonia-related outcomes.** *Clin Infect Dis* 2012, **55**(11):1466-1473.
74. Deshotels MR, Xia H, Sriramula S, Lazartigues E, Filipeanu CM: **Angiotensin II mediates angiotensin converting enzyme type 2 internalization and degradation through an angiotensin ii type i receptor-dependent mechanism.** *Hypertension* 2014, **64**(6):1368-1375.
75. Tao W, Li PS, Xu G, Luo Y, Shu YS, Tao YZ, Yang LQ: **Soluble Epoxide Hydrolase Plays a Vital Role in Angiotensin II-Induced Lung Injury in Mice.** *Shock* 2018, **50**(5):589-594.
76. Chen C, Zhang Z, Li Z, Zhang F, Peng M, Chen Y, Wang Y: **Losartan attenuates microvascular permeability in mechanical ventilator-induced lung injury in diabetic mice.** *Mol Biol Rep* 2014, **41**(2):809-814.
77. Yao S, Feng D, Wu Q, Li K, Wang L: **Losartan attenuates ventilator-induced lung injury.** *J Surg Res* 2008, **145**(1):25-32.
78. Ling X, Lou A, Li Y, Yang R, Ning Z, Li X: **[Losartan regulates oxidative stress via caveolin-1 and NOX4 in mice with ventilator- induced lung injury].** *Nan Fang Yi Ke Da Xue Xue Bao* 2015, **35**(12):1739-1744.
79. Adamzik M, Frey U, Sixt S, Knemeyer L, Beiderlinden M, Peters J, Siffert W: **ACE I/D but not AGT (-6) A/G polymorphism is a risk factor for mortality in ARDS.** *European Respiratory Journal* 2007, **29**(3):482-488.
80. Kim J, Choi SM, Lee J, Park YS, Lee CH, Yim JJ, Yoo CG, Kim YW, Han SK, Lee SM: **Effect of Renin-Angiotensin System Blockage in Patients with Acute Respiratory Distress Syndrome: A Retrospective Case Control Study.** *Korean J Crit Care Med* 2017, **32**(2):154-163.
81. Noveanu M, Breidthardt T, Reichlin T, Gayat E, Potocki M, Pargger H, Heise A, Meissner J, Twerenbold R, Muravitskaya N *et al*: **Effect of oral beta-blocker on short and long-term mortality in patients with acute respiratory failure: results from the BASEL-II-ICU study.** *Crit Care* 2010, **14**(6):R198.
82. Hsieh M-S, How C-K, Hsieh VC-R, Chen P-C: **Preadmission antihypertensive drug use and sepsis outcome: impact of angiotensin-converting enzyme inhibitors (ACEIs) and angiotensin receptor blockers (ARBs).** *Shock* 2020, **53**(4):407-415.
83. Mortensen EM, Restrepo MI, Copeland LA, Pugh JA, Anzueto A, Cornell JE, Pugh MJV: **Impact of previous statin and angiotensin II receptor blocker use on mortality in patients hospitalized with sepsis.** *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* 2007, **27**(12):1619-1626.
84. Boyle AJ, Mac Sweeney R, McAuley DF: **Pharmacological treatments in ARDS; a state-of-the-art update.** *BMC Med* 2013, **11**(1):166.

85. Huang F, Guo J, Zou Z, Liu J, Cao B, Zhang S, Li H, Wang W, Sheng M, Liu S *et al*: **Angiotensin II plasma levels are linked to disease severity and predict fatal outcomes in H7N9-infected patients.** *Nat Commun* 2014, **5**:3595.
86. Henry C, Zaizafoun M, Stock E, Ghamande S, Arroliga AC, White HD: **Impact of angiotensin-converting enzyme inhibitors and statins on viral pneumonia.** In: *Baylor University Medical Center Proceedings: 2018*: Taylor & Francis; 2018: 419-423.
87. Gu H, Xie Z, Li T, Zhang S, Lai C, Zhu P, Wang K, Han L, Duan Y, Zhao Z *et al*: **Angiotensin-converting enzyme 2 inhibits lung injury induced by respiratory syncytial virus.** *Sci Rep* 2016, **6**:19840.
88. Zou Z, Yan Y, Shu Y, Gao R, Sun Y, Li X, Ju X, Liang Z, Liu Q, Zhao Y *et al*: **Angiotensin-converting enzyme 2 protects from lethal avian influenza A H5N1 infections.** *Nat Commun* 2014, **5**:3594.
89. Yang P, Gu H, Zhao Z, Wang W, Cao B, Lai C, Yang X, Zhang L, Duan Y, Zhang S *et al*: **Angiotensin-converting enzyme 2 (ACE2) mediates influenza H7N9 virus-induced acute lung injury.** *Sci Rep* 2014, **4**:7027.
90. Yan Y, Liu Q, Li N, Du J, Li X, Li C, Jin N, Jiang C: **Angiotensin II receptor blocker as a novel therapy in acute lung injury induced by avian influenza A H5N1 virus infection in mouse.** *Sci China Life Sci* 2015, **58**(2):208-211.
91. Liu J, Zhang P, Yu T, Yang Y, Qiu H: **[Effects of losartan on pulmonary dendritic cells in lipopolysaccharide- induced acute lung injury mice].** *Zhonghua Yi Xue Za Zhi* 2014, **94**(41):3216-3219.
92. Raiden S, Pereyra Y, Nahmod V, Alvarez C, Castello L, Giordano M, Geffner J: **Losartan, a selective inhibitor of subtype AT1 receptors for angiotensin II, inhibits neutrophil recruitment in the lung triggered by fMLP.** *J Leukoc Biol* 2000, **68**(5):700-706.
93. Schoeman D, Fielding BC: **Coronavirus envelope protein: current knowledge.** *Virol J* 2019, **16**(1):69.
94. Wang S, Guo F, Liu K, Wang H, Rao S, Yang P, Jiang C: **Endocytosis of the receptor-binding domain of SARS-CoV spike protein together with virus receptor ACE2.** *Virus Res* 2008, **136**(1-2):8-15.
95. Jerng JS, Yu CJ, Wang HC, Chen KY, Cheng SL, Yang PC: **Polymorphism of the angiotensin-converting enzyme gene affects the outcome of acute respiratory distress syndrome.** *Crit Care Med* 2006, **34**(4):1001-1006.
96. Liu Y, Yang Y, Zhang C, Huang F, Wang F, Yuan J, Wang Z, Li J, Li J, Feng C *et al*: **Clinical and biochemical indexes from 2019-nCoV infected patients linked to viral loads and lung injury.** *Sci China Life Sci* 2020, **63**(3):364-374.
97. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, Xiang J, Wang Y, Song B, Gu X *et al*: **Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study.** *Lancet* 2020, **395**(10229):1054-1062.
98. Xudong X, Junzhu C, Xingxiang W, Furong Z, Yanrong L: **Age-and gender-related difference of ACE2 expression in rat lung.** *Life sciences* 2006, **78**(19):2166-2171.
99. Cai H: **Sex difference and smoking predisposition in patients with COVID-19.** *Lancet Respir Med* 2020, **8**(4):e20.
100. Hilliard LM, Jones ES, Steckelings UM, Unger T, Widdop RE, Denton KM: **Sex-specific influence of angiotensin type 2 receptor stimulation on renal function: a novel therapeutic target for hypertension.** *Hypertension* 2012, **59**(2):409-414.
101. Packer M, McMurray JJV: **Importance of endogenous compensatory vasoactive peptides in broadening the effects of inhibitors of the renin-angiotensin system for the treatment of heart failure.** *Lancet* 2017, **389**(10081):1831-1840.
102. Podowski M, Calvi C, Metzger S, Misono K, Poonyagariyagorn H, Lopez-Mercado A, Ku T, Lauer T, McGrath-Morrow S, Berger A *et al*: **Angiotensin receptor blockade attenuates cigarette smoke-induced lung injury and rescues lung architecture in mice.** *J Clin Invest* 2012, **122**(1):229-240.
103. Leung JM, Yang CX, Tam A, Shaipanich T, Hackett TL, Singhera GK, Dorscheid DR, Sin DD: **ACE-2 Expression in the Small Airway Epithelia of Smokers and COPD Patients: Implications for COVID-19.** *Eur Respir J* 2020:2000688.

104. Yilin Z, Yandong N, Faguang J: **Role of angiotensin-converting enzyme (ACE) and ACE2 in a rat model of smoke inhalation induced acute respiratory distress syndrome.** *Burns* 2015, **41**(7):1468-1477.
105. Fang L, Karakiulakis G, Roth M: **Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection?** *Lancet Respir Med* 2020, **8**(4):e21.
106. Burrell LM: **A risk-benefit assessment of losartan potassium in the treatment of hypertension.** *Drug Saf* 1997, **16**(1):56-65.
107. Christen Y, Waeber B, Nussberger J, Porchet M, Borland RM, Lee RJ, Maggon K, Shum L, Timmermans PB, Brunner HR: **Oral administration of DuP 753, a specific angiotensin II receptor antagonist, to normal male volunteers. Inhibition of pressor response to exogenous angiotensin I and II.** *Circulation* 1991, **83**(4):1333-1342.
108. Doig JK, MacFadyen RJ, Sweet CS, Reid JL: **Haemodynamic and renal responses to oral losartan potassium during salt depletion or salt repletion in normal human volunteers.** *J Cardiovasc Pharmacol* 1995, **25**(4):511-517.
109. Ohtawa M, Takayama F, Saitoh K, Yoshinaga T, Nakashima M: **Pharmacokinetics and biochemical efficacy after single and multiple oral administration of losartan, an orally active nonpeptide angiotensin II receptor antagonist, in humans.** *Br J Clin Pharmacol* 1993, **35**(3):290-297.
110. Reid JL: **Inhibitors of the renin-angiotensin system. Clinical pharmacology studies on kinetics, dynamics and concentration-effect relationships.** *Arzneimittelforschung* 1993, **43**(2A):263-264.
111. Meng J, Xiao G, Zhang J, He X, Ou M, Bi J, Yang R, Di W, Wang Z, Li Z *et al*: **Renin-angiotensin system inhibitors improve the clinical outcomes of COVID-19 patients with hypertension.** *Emerg Microbes Infect* 2020, **9**(1):757-760.
112. Bean D, Kraljevic Z, Searle T, Bendayan R, Pickles A, Folarin A, Roguski L, Noor K, Shek A, O'Gallagher K *et al*: **Treatment with ACE-inhibitors is not associated with early severe SARS-Covid-19 infection in a multi-site UK acute Hospital Trust;** 2020.
113. Jazzi AS, Mahnam K, Hejazi SH, Damavandi MS, Sadeghi P, Zeinalian M, Tabesh F, Mirbod SM, Khanahmad H: **Inhibition of Viral Macrodomein of COVID-19 and Human TRPM2 by losartan.** 2020.
114. International Society of Hypertension: **A statement from the International Society of Hypertension on COVID-19 | The International Society of Hypertension.** 2020.
115. **Hypertension Canada's Statement on: Hypertension, ACE-Inhibitors and Angiotensin Receptor Blockers and COVID-19** [<https://hypertension.ca/wp-content/uploads/2020/03/2020-30-15-Hypertension-Canada-Statement-on-COVID-19-ACEi-ARB.pdf>]
116. **COVID-19 and concerns regarding use of ACEi/ARB/ARNi medications for heart failure or hypertension** [<https://www.conferenceharvester.com/Uploads/Documents/9006/CCSCHFSstatementregardingCOVID.pdf>]

Figure Legend

Figure 1: Renin-angiotensin system with COVID-19

ACEi angiotensin converting enzyme inhibitor, ACE1 angiotensin converting enzyme, ACE2 angiotensin converting enzyme 2, ARB angiotensin receptor blocker, AT₁R type 1 angiotensin II receptor, AT₂R type 2 angiotensin II receptor, rhACE2 recombinant human angiotensin converting enzyme 2.

Figure 1: Renin-angiotensin system with COVID-19

Table 1 Summary (in text)

Table 2 Summary (in text)

Table 3 Summary (in text)

