




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Clinical evidence for repurposing chloroquine and hydroxychloroquine as antiviral agents: a systematic review

Chaturaka Rodrigo, Sumadhya Deepika Fernando, Senaka Rajapakse



PII: S1198-743X(20)30293-7

DOI: <https://doi.org/10.1016/j.cmi.2020.05.016>

Reference: CMI 2063

To appear in: *Clinical Microbiology and Infection*

Received Date: 8 April 2020

Revised Date: 5 May 2020

Accepted Date: 16 May 2020

Please cite this article as: Rodrigo C, Fernando SD, Rajapakse S, Clinical evidence for repurposing chloroquine and hydroxychloroquine as antiviral agents: a systematic review, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2020.05.016>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Ltd on behalf of European Society of Clinical Microbiology and Infectious Diseases.

**Clinical evidence for repurposing chloroquine and hydroxychloroquine as
antiviral agents: a systematic review**

Type of article: systematic review

Chaturaka Rodrigo¹, Sumadhya Deepika Fernando², Senaka Rajapakse³

¹Department of Pathology, School of Medical Sciences, UNSW Sydney, Australia

²Department of Parasitology, Faculty of Medicine, University of Colombo, Sri Lanka

³Department of Clinical Medicine, Faculty of Medicine, University of Colombo, Sri Lanka

Corresponding author

Prof. Senaka Rajapakse

Faculty of Medicine, University of Colombo

25, Kynsey Road, Colombo 08, Sri Lanka

Email: senaka@med.cmb.ac.lk

Abstract

Background: Repurposing hydroxychloroquine (HCQ) and chloroquine (CQ) as antiviral agents is a re-emerging topic with new viral epidemics.

Objectives: To summarize evidence from human clinical studies for using HCQ or CQ as antiviral agents for any viral infection

Data sources: PubMed, EMBASE, Scopus, Web of Science for published studies without time or language restrictions. Cochrane Clinical Trial Registry and Chinese Clinical Trials Registry for trials registered after 2015. MedRxiv for pre-prints within the last 12 months.

Study eligibility criteria: Interventional and prospective observational studies (with or without a control group)

Participants: Adults and children with a confirmed viral infection

Intervention: Use of chloroquine or hydroxychloroquine as antiviral agents in one or more groups of the study

Methods: Two authors independently screened abstracts and all authors agreed on eligible studies. A meta-analysis was planned if similar studies were available in terms of participants, intervention, comparator and outcomes.

Results: Nineteen studies were eligible (HIV: 8, HCV: 2, Dengue: 2, Chikungunya: 1, COVID-19: 6) including two pre-prints. Nine and ten studies assessed CQ and HCQ respectively. Benefits of either drug for viral load suppression in HIV is inconsistent. CQ is ineffective in curing dengue (high certainty evidence) and may have little or no benefit in curing chikungunya (low-certainty evidence). The evidence for COVID-19 infection is rapidly evolving but at this stage we are unsure if CQ or HCQ has any benefit in clearing viraemia (very low certainty evidence).

Conclusions: Using HCQ or CQ for HIV / HCV infections is clinically irrelevant now as other effective antivirals are available for viral load suppression (HIV) and cure (HCV). There is no benefit of CQ in dengue and the same conclusion is likely for chikungunya infection. More evidence is needed to confirm if HCQ or CQ is beneficial in COVID-19 infection.

Keywords: Covid-19; pneumonia; hydroxychloroquine; chloroquine; antivirals; Dengue; Chikungunya

Introduction

With the pandemic of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2) or COVID-19, repurposing cheap and accessible drugs for off-label use as antivirals is gaining popularity if they have demonstrated antiviral properties *in vitro* or in animal studies. Chloroquine (CQ), a well-established anti-malarial agent, and hydroxychloroquine (HCQ), a similarly established disease modifying anti-rheumatic (DMARD) agent have received increased attention in recent days for their purported efficacy as antiviral agents in context of COVID-19. Both drugs are out-of-patent, cheap, and widely available in high, middle- and low-income countries.

The antiviral properties of CQ was first explored against viral hepatitis as far back as 1963(1). Since then many observations from *in vitro* and animal experiments have suggested a beneficial role of HCQ and CQ in viral infections(2-13). Yet, the ultimate test for their benefit as antivirals come from human clinical studies and to our knowledge both these drugs are not used as mainstream antiviral agents for any viral infection. This review focusses on the clinical evidence for using both CQ and HCQ as antiviral agents against any viral infection. This reflection and summary of evidence is needed in current context for judicious, evidence-based recommendations for off-label use of these drugs. Even when such evidence may be incomplete or unavailable for emerging infections such as COVID-19, historical attempts for repurposing these agents for other emerging viral infections from time to time, may draw parallels to the current scenario to inform a rational approach to clinical trials and guideline recommendations.

Methods

Study selection criteria

Types of studies: Interventional and observational studies (controlled and non-controlled) including case series were considered, but case reports limited to single patients were excluded. Retrospective studies, animal or *in vitro* experiments were excluded.

Participants: Adults or children with a confirmed viral infection

Intervention and comparator: All participants (non-controlled studies) or one study arm (controlled studies) must have received either CQ or HCQ as antiviral agents (as stand-alone therapy or in combination with other treatments). The comparator for controlled studies were standard treatment, no treatment or placebo.

Outcomes: Primary outcome was viral load suppression for chronic infections and clearance of viraemia for acute infections. Any other significant outcomes (depending on the type of infection) are discussed narratively as reported by authors given the broad scope of this review.

Data sources and search strategy

We searched PubMed, EMBASE, Scopus, Web of Science, Cochrane Clinical Trials Registry (CENTRAL), Chinese Clinical Trials Registry and MedRxiv (for pre-prints) according to the search strategy detailed in table 1. Bibliographies of eligible articles were also searched. The date of the last search was 30th April 2020. Two authors searched and selected abstracts independently and all authors identified studies for full-text review by consensus. The following data items were extracted from each included study; study design and location, viral infection and diagnostic criteria, participant demographics, intervention and control groups (if any), drug doses, primary and secondary outcomes and adverse events attributable to therapy. A meta-analysis was planned if comparable clinical trials were available in terms of participants, interventions, comparators and outcomes. Risk of bias for all randomized controlled studies, regardless of whether published or deposited as pre-prints, were assessed according to Cochrane guidelines(14). Risk of bias in non-randomized trials with more than one intervention was assessed with the ROBINS-I tool(15). All other study designs were considered to have an inherent high risk of bias. Certainty of evidence was assessed for each infection (only from randomized controlled trials - RCTs) according to the GRADE recommendations(16). Uncontrolled studies and RCTs only available as preprints were not considered for assessing certainty of evidence.

Results

A total of 2267 abstract results were found from all databases after removing duplicates (figure 1). The majority of these (n=2026) were published after 1990. After removing retrospective studies, *in vitro* and animal studies, reviews, letters, opinion papers and editorials, only 209 abstracts remained. Further examination identified 26 articles for full-text review revealing a significant disparity with the large number of articles demonstrating *in vitro* evidence for antiviral effects of HCQ and CQ. Only 19 publications (including two preprints) from this subset met the inclusion criteria. One full text paper was in Mandarin but had an English abstract. The studies included randomized and non-randomized controlled trials and uncontrolled prospective studies equivalent to case series. These had tested the efficacy of CQ and HCQ in human immunodeficiency virus (HIV), dengue, chikungunya, Hepatitis C and COVID-19 infections. A meta-analysis was not performed due to unavailability of similar studies. A summary of risk of bias for all controlled clinical trials are given in figure 2. The results are discussed separately for each infection and under each section, randomized clinical trial are discussed first, followed by other study designs. Chronic infections (HIV/HCV) are discussed first as the aim of treatment is viral load suppression rather than cure. Details of study design and setting, participants, interventions, comparators, outcomes and risk of bias for each included study is given in table 2. There are 23 trials in progression, all examining the role of HCQ and CQ in COVID-19 infection (Table 3).

HIV

There were six RCTs (five double blind studies and one open label study)(17-22) and two prospective non-controlled studies(23, 24) on HIV that met the eligibility criteria; Five studies used HCQ (including both non-controlled studies)(17, 18, 20, 23, 24) and three used CQ(19, 21, 22).

HCQ for HIV: Enrolments in these RCTs varied from 40 patients(17) to 83 patients(20). Two RCTs enrolled adults with a CD4 cell count >200/ μ l (17, 18) and the other enrolled adults with a CD4 cell count >400/ μ l (20). HCQ doses used ranged from 400mg/d – 800mg/d and comparators included

placebo(17, 20) or zidovudine(18). Two RCTs reported a significant decrease in viral load with no change in absolute CD4 cell count with HCQ (17, 18) while the third RCT(20) recorded a significant increase in viral load and a decrease in the CD4 cell count with HCQ. The two uncontrolled studies were small (n = 20-22) and used HCQ 400mg/d alone (23) or in combination with hydroxyurea plus didanosine (24). There was a modest reduction in viral load after 48 weeks with combination therapy(24) while markers of HIV immune activation was reduced with HCQ in the other study after 6 months of treatment (23) suggesting an immunomodulatory role of HCQ. HCQ was well tolerated in all studies.

CQ for HIV: Two double blind RCTs enrolled adult patients with HIV (CQ 150-300mg base /d, 8-12 weeks)(21, 22) and one open label trial enrolled paediatric patients (CQ 4mg/kg/d for 24 weeks)(19). The smallest trial had 13 patients(22) while the largest had 70 patients(21). The comparators were either placebo(21, 22) or zidovudine plus didanosine (paediatric trial)(19). Both adult trials enrolled patients with a CD4 cell count of at least 250/ μ l and demonstrated a modestly beneficial immunomodulatory effect of CQ (a significant decline in activated CD8 cells) (21, 22). However, in one trial which had a complex cross-over design, the HIV viral load increased when CQ was administered without any other antiretroviral therapy(21). There was no significant virological or immunological improvement in with CQ in the third (paediatric) trial (19) but significantly more gastrointestinal adverse events were observed with CQ.

Hepatitis C

Use of CQ or HCQ for HCV infection has been evaluated in two RCTs, one a single blinded study (n=120, HCV genotype 4) and another a double blinded study (n=10, HCV genotype 1) (25-27). The first trial used HCQ 400mg/d for 12 weeks added to interferon (IFN) and ribavirin therapy while the control group received IFN and ribavirin only. HCQ was well tolerated and a significantly higher rate of early virological response was noted in the HCQ group (p=0.011). The other trial used CQ 150mg base /d vs. placebo in patients who has already failed IFN and ribavirin therapy and noted a

significant drop in viral RNA levels following 8 weeks of therapy ($p=0.04$), but this effect was transient.

IFN and ribavirin therapy for HCV is largely outdated nowadays as highly effective directly acting antiviral agents are now available. Neither HCQ nor CQ has been evaluated in combination with these agents.

Dengue

Two studies had assessed CQ for dengue infection (none for HCQ). A double-blind RCT ($n=307$) in Vietnam administered CQ 1500mg over 3 days (dosage regimen the same as that used for malaria – table 2) or placebo to clinically suspected dengue patients within first 3 days of fever. A large proportion of the enrolments (84%) were later confirmed to have dengue. CQ failed to demonstrate a clinical benefit in; clearance of viraemia, clearance of NS1 antigen positivity, fever clearance time, preventing dengue haemorrhagic fever, preventing decrease in platelet count or increase in haematocrit (surrogate markers of dengue associated plasma leakage), but significantly increased gastrointestinal adverse events ($p<0.05$) (28). The second study enrolled 129 patients with clinically suspected dengue to a two-arm randomized controlled trial to receive either 600mg CQ base/d for 3 days or placebo. Only 37 participants were later confirmed to have dengue (19 in CQ group; 18 in placebo group)(29). There was no statistically significant difference in the total duration of illness or adverse events.

Chikungunya

Chikungunya is a viral illness endemic in the tropics. Acute infection mimics a viral flu and sometimes lead to a debilitating, persistent inflammatory arthritis. There was only one RCT that had evaluated CQ as an antiviral agent (and none for HCQ) in acute chikungunya infection. This double blind RCT, conducted in French Reunion Islands in 2006, enrolled 54 adult patients with virologically confirmed acute chikungunya infection to receive either CQ (600mg of base/day for 3 days followed by

300mg/d for 2 days) or placebo (30, 31). There was no statistically significant difference in time for viraemia clearance or fever clearance in CQ treated group. The adverse effects were minor but all (n=7) were reported in the CQ group. When followed up at day 200 (since onset of symptoms) by a telephone interview, the CQ treated group was significantly more likely to have persistent arthralgia. However, a later analysis of cytokines and chemokines from plasma samples revealed higher levels of IFN- α , IL-6, IL-8 and MCP-1 in the group that received CQ (more severe disease at baseline and hence, increased likelihood of persistent arthralgia)(32).

Given its established role as a DMARD, HCQ is potentially useful to treat chikungunya induced chronic arthritis. A randomized controlled trial had evaluated HCQ for this purpose after acute chikungunya infection (33). However, as HCQ was not “repurposed” as an antiviral agent, this trial was outside the scope of this review.

Severe Acute Respiratory Syndrome Coronavirus -2 (SARS-CoV-2)

The ongoing epidemic of SARS-CoV-2 (referred to as COVID -19) is rapidly evolving. Potential use of CQ for COVID-19 associated pneumonia first emerged in February 2020 based on several Chinese studies and an expert consensus from Guangdong Province of China (abstract in English) recommends using CQ 500mg (presumably 300mg of CQ base) two times daily for 10 days to treat patients with COVID-19 associated pneumonia(34). We could not verify the evidence for this recommendation independently as the original full-text articles were not available. We have summarized the evidence for using HCQ or CQ for COVID-19 from full-text articles that were available within our search strategy by 30th April 2020 below. Pre-prints from MedRxiv are also discussed but these are not peer reviewed publications and hence not considered in certainty of evidence assessment. Ongoing trials are listed in table 3.

HCQ for COVID-19: There are two peer reviewed publications (one RCT and a non-randomized controlled study)(35, 36) and two non-peer reviewed pre-prints (two RCTs) on this topic(37, 38). A third study (from USA) available on MedRxiv as a pre-print was excluded as it was a retrospective

analysis(39). The only peer reviewed publication of a RCT (in Mandarin, abstract in English) was not indexed in any of the databases searched and was identified from a secondary bibliography search. It describes a randomized clinical trial (NCT04261517) that recruited 15 non-severe, confirmed COVID-19 patients to an HCQ arm (400mg/d for 5 days) and a no HCQ arm(36). Clearance of viraemia (RT-PCR of a throat swab) by day 7, fever clearance time and total duration of hospital stay were similar between the two groups. A non-randomized controlled trial from Marseille, France (full-paper reviewed) enrolled 26 patients with virologically confirmed COVID-19 to receive HCQ 600mg/d(35). Twenty patients completed HCQ treatment (6 patients also received azithromycin) and all had radiological evidence of pneumonia (CT scan). The control group (n=16) was younger (mean age; 51.2 vs. 37.3 years) and had patients treated at another institution or those treated at the same institution but refused HCQ. More patients in the HCQ group were virologically cured by day 6 (70% vs. 12.5%, $p=0.001$) including all patients who received azithromycin.

Results of two more RCTs from China on using HCQ in COVID-19 are available as pre-prints (37, 38). The largest was a multicentre open label clinical trial (ChiCTR2000029868) that enrolled 150 virologically confirmed COVID-19 patients (75 per group) to receive HCQ plus standard care (as defined by Chinese guidelines for COVID-19 management) or standard care only(37). HCQ group received 200mg/d for 3 days followed by 800mg/d for 14-21 days. Only two patients (one in each group) had severe illness. There was no statistically significant difference in the proportion of aviraemic patients by day 28 (85.4% in HCQ group vs 81.3% in control group, $p>0.05$). There was also no difference in time to aviraemia, or symptom resolution by day 28. Many secondary outcomes proposed in methods including all-cause mortality, are not reported in results presumably due to lack of events. The second study (ChiCTR2000029559) enrolled and randomized 62 (31 in each group) virologically confirmed COVID-19 patients with evidence of pneumonia on CT scan to receive HCQ (400mg/d for 5 days) or no HCQ treatment. The respiratory distress was not severe ($\text{PaO}_2/\text{FIO}_2>300$ or $\text{SaO}_2/\text{SPO}_2>93\%$) at enrolment in all recruits. The authors report faster time to clinical recovery (defined as normalization of body temperature and cough relief maintained for 72

hours) and earlier radiological improvement in the HCQ group, but the calculations and radiological reporting standards for these conclusions are unclear. Four patients in the control group vs none in the HCQ group progressed to severe illness.

CQ for COVID-19: There are two published RCTs on using CQ for COVID-19. The first study from China, randomized 22 virologically confirmed COVID-19 patients to receive either Lopinavir/Ritonavir combination or CQ 600mg (base)/day for 10 days (40). There was no statistically significant difference for having a negative RT-PCR by day 14 of illness (10/10 in CQ group vs. 11/12). However, on average, the Lopinavir/Ritonavir group had started treatment later than those in CQ group (2.5 vs. 6.5 days since the onset of illness, $p < 0.001$). The second RCT was a dose ranging study for CQ which did not have a no CQ comparator group (41). It randomized 81 patients clinically suspected of having COVID-19 to receive either CQ 12g over 10 days or 2.7g over 5 days. Diagnosis was later confirmed (virologically) in 31 patients in each group. The recruits had severe infection at time of enrolment as defined by tachycardia, tachypnoea, oxygen saturation $< 90\%$, and hypotension. Recruitment to the high dose arm was terminated early (by day 13) as mortality was significantly higher in this group (15% vs 39%, $p = 0.03$).

Discussion

Despite being researched for several decades and being supported by a large volume of *in vitro* and animal study data for plausible mechanisms for antiviral effects, neither HCQ nor CQ are currently recommended as antiviral agents for any of the infections they had been tested for in clinical trials (with the exception of COVID-19 which is an evolving situation at the time of writing). The most researched infection in this regard is HIV. However, mainstream anti-retroviral treatment in HIV is highly successful in viral load suppression allowing reasonable control of disease though it is not a cure. These newer combinations have not been tested against CQ/HCQ and given the inconsistent evidence from existing trials in this regard, there is no need to. The same can be said of HCV infection where pan-genotypic directly antiviral agents can now achieve over 90% cure rates with

minimum adverse events(42). Repurposing HCQ/CQ for these infections is largely redundant and of historical interest only.

In contrast, for acute viral infections such as dengue, chikungunya or COVID-19, effective antivirals are not available. These infections cause epidemics in vulnerable populations exerting an enormous financial burden on resource limited healthcare systems. For dengue fever, given the evidence presented here we conclude that CQ is of no benefit (high certainty evidence). This conclusion is supported by a single well designed RCT with an adequate sample size and a low risk of bias. CQ may have little or no effect in curing acute chikungunya infection (low certainty evidence) and this conclusion is supported by a small clinical trial which may have inadvertently recruited people with more severe disease to the CQ arm despite randomization.

For COVID-19 we are unsure if either CQ or HCQ is of any benefit as per currently available evidence (very low certainty evidence). The only peer reviewed publication describing a RCT on using HCQ for COVID-19 was a small trial that did not demonstrate any benefit of HCQ. The other peer reviewed publication which did show a benefit was an open label, non-randomized trial with a serious risk of bias in sample selection, confounders and assessing outcomes. The sample size was small, and there was a marked age discrepancy (of approximately 14 years) between test and control groups.

Administration of azithromycin to six subjects who also received HCQ, does not seem to be based on an *a priori* hypothesis or a study protocol. Some control group patients were from a different centre, and it is unclear if management protocols in both institutions were the same. More importantly, it is now established that most patients (>95%) with COVID-19 will recover after a mild infection regardless of antiviral treatment. Therefore, the primary outcome assessed in this study (clearance of viraemia) is less useful for patient management compared to outcomes that demonstrate a benefit in the more severe end of the disease spectrum such as mortality benefit, reduction in intensive care unit admissions or faster discharge from intensive care (or high dependency units), faster recovery from assisted ventilation or prevention of the need for assisted ventilation.

Examining such outcomes require a well-coordinated multi-centre study (preferably with a randomized, double blind, study design) with similar management protocols across all centres. The other two studies from China (available as pre-prints) mentioned in the results section also does not help to resolve this issue due to methodological issues and conflicting conclusions. For example in one of these studies(38), the control group is said to have received “standard” treatment which includes “antivirals, antibiotics and steroids or immunoglobulins”. These may have a serious confounding effect on the interpretation of results.

The role of CQ in COVID-19 is also unclear and not supported by evidence at the moment. One small RCT reported no benefit of CQ in terms of achieving negative viraemia by day 14 compared to treatment with lopinavir and ritonavir(40). The authors report faster radiological improvement and reduced hospital stay with CQ but on average the CQ group received treatment at an earlier stage of the illness than the comparator group. This plus the small sample size make it difficult to interpret the results. The other RCT did not have a no CQ control group which is rather surprising. Instead of proving the efficacy of CQ, the authors investigated if a higher dose of CQ was safe in patients with severe infection. Notably this CQ dose (1200mg/d for 10 days) was much higher than that used for cure of malaria which is the standard indication for CQ. Mortality was significantly higher in the high CQ dose group and patient recruitment was halted prematurely. Thus, this trial also doesn't help to confirm if CQ is beneficial for COVID-19.

Conclusion

CQ and HCQ have been examined for their antiviral properties in many *in vitro* and animal studies for more than five decades and in a limited number of human clinical studies spanning over 25 years. For HIV and HCV infections, the benefit of either drug is doubtful and perhaps no longer relevant as other effective treatments are now available for viral load suppression (HIV) or cure (HCV). There is good evidence that CQ is ineffective for curing dengue infection or preventing dengue haemorrhagic fever. CQ also may not have any benefit in curing acute chikungunya infection. Role of HCQ or CQ in

297 COVID-19 is yet unclear and needs to be assessed by well-designed randomized double-blind clinical
298 trials.

299 **Authors' statements**

300 **Author contributions:** SR conceived the study. CR, SR and SDF independently performed the
301 literature review. CR did the preliminary analysis and wrote the first draft which was double checked
302 by other authors. All authors approved the final version.

303 **Funding:** No grants or funding were received for this study.

304 **Conflict of interest:** None of the authors have any conflicts of interest regarding the content of this
305 manuscript.

306 **Ethical approval:** not required

307

308 References

- 309 1. Pareja-Coronel A. Treatment of viral hepatitis with chloroquine. *Am J Gastroenterol.*
310 1963;39:288-98.
- 311 2. Keyaerts E, Vijgen L, Maes P, Neyts J, Van Ranst M. In vitro inhibition of severe acute
312 respiratory syndrome coronavirus by chloroquine. *Biochem Biophys Res Commun.* 2004;323(1):264-
313 8.
- 314 3. Vincent MJ, Bergeron E, Benjannet S, Erickson BR, Rollin PE, Ksiazek TG, et al. Chloroquine is
315 a potent inhibitor of SARS coronavirus infection and spread. *Virology.* 2005;2.
- 316 4. De Clercq E. Potential antivirals and antiviral strategies against SARS coronavirus infections.
317 *Expert Rev Anti-Infect Ther.* 2006;4(2):291-302.
- 318 5. Rolain JM, Colson P, Raoult D. Recycling of chloroquine and its hydroxyl analogue to face
319 bacterial, fungal and viral infections in the 21st century. *Int J Antimicrob Agents.* 2007;30(4):297-
320 308.
- 321 6. Savarino A, Boelaert JR, Cassone A, Majori G, Cauda R. Effects of chloroquine on viral
322 infections: An old drug against today's diseases? *Lancet Infect Dis.* 2003;3(11):722-7.
- 323 7. Savarino A, Di Trani L, Donatelli I, Cauda R, Cassone A. New insights into the antiviral effects
324 of chloroquine. *Lancet Infect Dis.* 2006;6(2):67-9.
- 325 8. Han Y, Pham HT, Xu H, Quan Y, Mesplède T. Antimalarial drugs and their metabolites are
326 potent Zika virus inhibitors. *J Med Virol.* 2019;91(7):1182-90.
- 327 9. Kumar A, Liang B, Aarthi M, Singh SK, Garg N, Mysorekar IU, et al. Hydroxychloroquine
328 Inhibits Zika Virus NS2B-NS3 Protease. *ACS Omega.* 2018;3(12):18132-41.
- 329 10. Shiryayev SA, Mesci P, Pinto A, Fernandes I, Sheets N, Shresta S, et al. Repurposing of the
330 anti-malaria drug chloroquine for Zika Virus treatment and prophylaxis. *Sci Rep.* 2017;7(1).
- 331 11. Akpovwa H. Chloroquine could be used for the treatment of filoviral infections and other
332 viral infections that emerge or emerged from viruses requiring an acidic pH for infectivity. *Cell*
333 *Biochem Funct.* 2016;34(4):191-6.
- 334 12. Barclay W, Long J, Wright E, Molesti E, Temperton N. Antiviral therapies against Ebola and
335 other emerging viral diseases using existing medicines that block virus entry. *F1000 Res.* 2015;4.
- 336 13. Broder CC. Henipavirus outbreaks to antivirals: The current status of potential therapeutics.
337 *Curr Opin Virol.* 2012;2(2):176-87.
- 338 14. Higgins JPT, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane
339 Collaboration's tool for assessing risk of bias in randomised trials. *BMJ.* 2011;343:d5928.
- 340 15. Sterne JAC, Hernán MA, Reeves BC, Savović J, Berkman ND, Viswanathan M, et al. ROBINS-I:
341 a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ.* 2016;355:i4919.
- 342 16. Guyatt GH, Oxman AD, Vist GE, Kunz R, Falck-Ytter Y, Alonso-Coello P, et al. GRADE: an
343 emerging consensus on rating quality of evidence and strength of recommendations. *BMJ.*
344 2008;336(7650):924.
- 345 17. Sperber K, Louie M, Kraus T, Proner J, Sapira E, Lin S, et al. Hydroxychloroquine treatment of
346 patients with human immunodeficiency virus type 1. *Clin Ther.* 1995;17(4):622-36.
- 347 18. Sperber K, Chiang G, Chen H, Ross W, Chusid E, Gonchar M, et al. Comparison of
348 hydroxychloroquine with zidovudine in asymptomatic patients infected with human
349 immunodeficiency virus type 1. *Clin Ther.* 1997;19(5):913-23.
- 350 19. Engchanil C, Kosalaraksa P, Lumbiganon P, Lulitanond V, Pongjunyakul P, Thuennadee R, et
351 al. Therapeutic potential of chloroquine added to zidovudine plus didanosine for HIV-1 infected
352 children. *J Med Assoc Thai.* 2006;89(8):1229-36.
- 353 20. Paton NI, Goodall RL, Dunn DT, Franzen S, Collaco-Moraes Y, Gazzard BG, et al. Effects of
354 hydroxychloroquine on immune activation and disease progression among HIV-infected patients not
355 receiving antiretroviral therapy: a randomized controlled trial. *JAMA.* 2012;308(4):353-61.

21. Jacobson JM, Bosinger SE, Kang M, Belaunzaran-Zamudio P, Matining RM, Wilson CC, et al. The Effect of Chloroquine on Immune Activation and Interferon Signatures Associated with HIV-1. *AIDS Res Hum Retroviruses*. 2016;32(7):636-47.
22. Murray SM, Down CM, Boulware DR, Stauffer WM, Cavert WP, Schacker TW, et al. Reduction of immune activation with chloroquine therapy during chronic HIV infection. *J Virol*. 2010;84(22):12082-6.
23. Piconi S, Parisotto S, Rizzardini G, Passerini S, Terzi R, Argentero B, et al. Hydroxychloroquine drastically reduces immune activation in HIV-infected, antiretroviral therapy-treated immunologic nonresponders. *Blood*. 2011;118(12):3263-72.
24. Paton NI, Aboulhab J, Karim F. Hydroxychloroquine, hydroxycarbamide, and didanosine as economic treatment for HIV-1. *Lancet*. 2002;359(9318):1667-8.
25. Peymani P, Ghavami S, Yeganeh B, Tabrizi R, Sabour S, Geramizadeh B, et al. Effect of chloroquine on some clinical and biochemical parameters in non-response chronic hepatitis C virus infection patients: pilot clinical trial. *Acta Biomed*. 2016;87(1):46-53.
26. Peymani P, Yeganeh B, Sabour S, Geramizadeh B, Fattahi MR, Keyvani H, et al. New use of an old drug: Chloroquine reduces viral and ALT levels in HCV non-responders (a randomized, triple-blind, placebo-controlled pilot trial). *Can J Physiol Pharmacol*. 2016;94(6):613-9.
27. Helal GK, Gad MA, Abd-Allah MF, Eid MS. Hydroxychloroquine augments early virological response to pegylated interferon plus ribavirin in genotype-4 chronic hepatitis C patients. *J Med Virol*. 2016;88(12):2170-8.
28. Tricou V, Minh NN, van TP, Lee SJ, Farrar J, Wills B, et al. A randomized controlled trial of chloroquine for the treatment of dengue in vietnamese adults. *PLoS Negl Trop Dis*. 2010;4(8).
29. Borges MC, Castro LA, Fonseca BA. Chloroquine use improves dengue-related symptoms. *Mem Inst Oswaldo Cruz*. 2013;108(5):596-9.
30. De Lamballerie X, Boisson V, Reynier JC, Enault S, Charrel RN, Flahault A, et al. On chikungunya acute infection and chloroquine treatment. *Vector Borne Zoonotic Dis*. 2008;8(6):837-9.
31. de Lamballerie X, Ninove L, Charrel RN. Antiviral treatment of chikungunya virus infection. *Infect Disord Drug Targets*. 2009;9(2):101-4.
32. Roques P, Thiberville SD, Dupuis-Maguiraga L, Lum FM, Labadie K, Martinon F, et al. Paradoxical effect of chloroquine treatment in enhancing chikungunya virus infection. *Viruses*. 2018;10(5).
33. Ravindran V, Alias G. Efficacy of combination DMARD therapy vs. hydroxychloroquine monotherapy in chronic persistent chikungunya arthritis: a 24-week randomized controlled open label study. *Clin Rheumatol*. 2017;36(6):1335-40.
34. Multicenter collaboration group of Department of Science Technology of Guangdong Province Health Commission of Guangdong Province for chloroquine in the treatment of novel coronavirus pneumonia. [Expert consensus on chloroquine phosphate for the treatment of novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43(3):185-8.
35. Gautret P, Lagier J-C, Parola P, Hoang VT, Meddeb L, Mailhe M, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents*. 2020:105949-.
36. Chen Jun LD, Liu Li, Liu Ping, Xu Qingnian, Xia Lu, Ling Yun, Huang Dan, Song Shuli, Zhang Dandan, Qian Zhiping, Li Tao, Shen Yinzong, Lu Hongzhou. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *Journal of Zhejiang University*. 2020;49(1):0-.
37. Tang W, Cao Z, Han M, Wang Z, Chen J, Sun W, et al. Hydroxychloroquine in patients with COVID-19: an open-label, randomized, controlled trial. *medRxiv*. 2020:2020.04.10.20060558.
38. Chen Z, Hu J, Zhang Z, Jiang S, Han S, Yan D, et al. Efficacy of hydroxychloroquine in patients with COVID-19: results of a randomized clinical trial. *medRxiv*. 2020:2020.03.22.20040758.

39. Magagnoli J, Narendran S, Pereira F, Cummings T, Hardin JW, Sutton SS, et al. Outcomes of hydroxychloroquine usage in United States veterans hospitalized with Covid-19. medRxiv. 2020:2020.04.16.20065920.
40. Huang M, Tang T, Pang P, Li M, Ma R, Lu J, et al. Treating COVID-19 with Chloroquine. Journal of Molecular Cell Biology. 2020.
41. Borba MGS, Val FFA, Sampaio VS, Alexandre MAA, Melo GC, Brito M, et al. Effect of High vs Low Doses of Chloroquine Diphosphate as Adjunctive Therapy for Patients Hospitalized With Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection: A Randomized Clinical Trial. JAMA Network Open. 2020;3(4):e208857-e.
42. Spengler U. Direct antiviral agents (DAAs) - A new age in the treatment of hepatitis C virus infection. Pharmacol Ther. 2018;183:118-26.

Tables

Table 1. Search strategy (Last date of search - 30th April 2020)

Database	Search terms	Field limits	Language limits	Time limits	Comments
PUBMED	'hydroxychloroquine AND antivir*', 'chloroquine AND antivir*', 'Hydroxychloroquine AND virus' and 'chloroquine AND virus'	None	None	None	
Scopus	As above	Title, Keywords or Abstract	None	None	
Web of Science	As above	None	None	None	
EMBASE	As above	None	None	None	
CENTRAL	"hydroxychloroquine" or "chloroquine"	None	None	2015-2020	Any unpublished/incomplete trial registered prior to 6 years was considered as unlikely to be completed
Chinese Clinical Trials Registry	"hydroxychloroquine" or "chloroquine"	None	None	2015-2020	As above
MedRxiv	"hydroxychloroquine" or "chloroquine"	Title or Abstract	None	Within 12 months	Once a pre-print is deposited, if the study met peer-review standards for publication, it is likely to have been published within this time frame.

Table 2. Summary of included studies

Study and location	Design	Inclusion criteria	Treatment regimens	Results - highlights	Risk of bias* and other comments
<i>HIV</i>					
Sperber et al. 1995, USA(17)	Randomized, double blind, placebo-controlled trial	HIV infected patients with CD4+ cell count between 200-500/ μ l and not taking antiretroviral (ART) therapy	1) HCQ 800mg/d for 8 weeks (n=20) or 2) Placebo for same duration (n=20)	Significant decrease in viral load in HCQ group (p=0.022) No change in absolute CD4+ cell count	Method of sample size calculation or randomization is not given HCQ was well tolerated
Sperber et al. 1997, USA(18)	Randomized, double blind, masked trial	HIV infected patients with CD4+ cell count between 200-500/ μ l	1) HCQ 800mg/d for 16 weeks (n=35) or 2) Zidovudine 500mg/d for same duration (n=37)	Significant decrease in viral load in both HCQ (p=0.02) and Zidovudine groups (p=0.001) No change in absolute CD4+ cell count	Method of sample size calculation or randomization is not given
Paton et al. 2012, United Kingdom(20)	Randomized, double blind, placebo-controlled trial	HIV infected asymptomatic patients, not on ART, CD4+ count > 400 cells/ μ l	HCQ 400mg/d (n=42) or placebo (n=41) for 48 weeks	No significant difference in CD8+ cell activation Greater decline of CD4+ cell count in HCQ group (p=0.03) Greater increase in HIV viral load in HCQ group (p<0.003)	High risk of attrition bias – 31% attrition rate HCQ was well tolerated despite more flu like episodes
Paton et al. 2002, Singapore(24)	Non controlled prospective study	HIV infected patients with a viral load < 10 ⁵ copies/ μ l and CD4+ cell count > 150/ μ l	HCQ 200mg/bid, hydroxyurea 500mg/bid and didanoside 125 – 200mg/bid according to bodyweight for 48 weeks (n=22)	Decline in viral load by an average of 1.3 logs with sustained CD4+ cell level	Neutropenia and elevated amylase levels noted in a majority of patients with two severe adverse events
Piconi et al. 2011, Italy (23)	Non controlled prospective study	HIV infected, adult, immunologic non responders (CD4+ cell count <200/ μ l or < 5% increase in preceding 12 months)	HCQ 400mg/d for 6 months (n=20)	No significant change in absolute CD4+ cell count, Significant reduction in Ki67 expressing (activated) CD4+ cells, Significant reduction in TLR-2, TLR-4 and TLR-5 expressing	Observations supportive of an immunomodulatory effect of HCQ to reduce immune activation in HIV

				CD14+ cells, Significant increase in TLR-2 and TLR-4 expressing regulatory T cells, Plasma IL-6 was significantly decreased Significant reduction in plasma lipopolysaccharide (LPS) levels	
Engchanil et al. 2006, Thailand(19)	Prospective, randomized open label clinical trial	HIV infected children (<14 years of age); CDC clinical category A,B or C; CDC immunological category 2 or 3	1) Zidovudine (ZDV)+ didanosine (ddI) (n=25) 2) ZDV+ddI+ CQ (n=30, includes 9 children where CQ was added to an existing ZDV+ddI therapy) ZDV 150 mg/m ² body surface area/dose every 8 hours; ddI 90-100 mg/m ² body surface area/dose every 12 hours; and chloroquine 4 mg/kg/dose daily Treatment continued for 24 weeks	No significant difference in the drop of viral load or increase in CD4+ cell count between the groups without CQ and with CQ	High risk of performance and detection bias due to open label design. High risk of attrition bias – 22% attrition rate More gastrointestinal adverse events noted in the group with CQ
Jacobson et al. 2016. USA(21)	Randomized, double blind, placebo- controlled cross- over trial	Two cohorts a) off ART for at least 6 months, HIV RNA >1000 copies/ml, CD4+ count >400/ μ l b) on ART for at least 24 months, HIV RNA level undetectable, CD4+ count>350/ μ l	Each cohort had two arms that either received CQ 250mg (150mg base) per day for 12 weeks followed by placebo for 12 weeks or vice versa 37 patients were on ART, 33 patients were off ART	No significant immune activation of CD4+ and CD8+ cells in off-ART cohort Significant reduction in activated CD8+ cells but not in CD4+ cells with CQ in the on- ART cohort. CQ also led to an increase in viral RNA load	High risk of attrition bias – 13% attrition rate No significant increase in adverse events reported for CQ
Murray et al. 2010. USA(22)	Randomized, double blind, placebo- controlled trial	HIV infected adults with CD4+ counts > 250/ μ l and either ART naïve or off ART for 16 months	250mg (150mg base) CQ daily (n=6, CQ 500mg daily (n=3), or placebo (n=4) for 2 months	Percentage of CD38+ and HLA DR+ CD8 cells were significantly reduced with CQ treatment (p=0.016)	Small clinical trial with 3-6 patients per trial arm. Method for sample size calculation is not mentioned CQ may reduce immune activation

in HIV					
<i>HCV</i>					
Helal et al. 2016. Egypt(27)	Randomized, prospective, single blind, controlled study	Patients with chronic active HCV (genotype 4) infection with chronic hepatitis on liver biopsy but without decompensated liver disease	Group 1: Pegylated interferon (IFN) 160µg subcutaneously weekly+ oral ribavirin 1000- 1200mg/d for 12 weeks (n=60) Group 2: as above + 400mg/d HCQ (n=60)	Significantly higher rate of early virological response in group 2 (p=0.011)	High risk of selection bias – method of allocation concealment not clear. High risk of performance or detection bias – single blind study design. No additional adverse events due to HCQ
Peymani et al. 2016. Iran(26)	Randomized, triple-blind, placebo- controlled pilot trial	Patients with HCV (genotype 1) not responding to standard IFN and ribavirin therapy	CQ 150mg (base)/d for 8 weeks or placebo (n=10)	Significant decrease in viral RNA (p=0.04) after 8 weeks	This is a small clinical trial and method of sample size calculation is unclear Rebound of viral RNA noted after stopping CQ
<i>Dengue</i>					
Tricou et al. 2010. Vietnam(28)	Randomized, double-blind, placebo- controlled trial	Patients with clinically suspected dengue, enrolled within first 3 days of fever and later confirmed with diagnostic testing (NS1 antigen test and RT-PCR)	Group 1: CQ 600mg/d (base) for 2 days and 300mg for 1 day (n=153) Group 2: placebo (n=153)	No statistically significant difference in clearance of antigenaemia (NS1), clearance of viraemia or incidence of dengue haemorrhagic fever	Low risk of bias CQ group had significantly more adverse events
Borges et al. 2013. Brazil (29)	Randomized double-blind study	Patients with clinically suspected dengue, later confirmed with diagnostic testing	Group 1: CQ 600mg (base) for 3 days (n=19) Group 2: placebo (n=18)	No statistically significant difference in the duration of fever	High risk of bias as most recruited patients was not confirmed to have dengue. Sample size calculation is unclear. Subjective improvement of symptoms was noted with CQ which reversed upon ceasing the regimen
<i>Chikungunya</i>					

De Lamballerie et al. French Reunion Islands. 2008(30)	Randomized, double-blind, placebo-controlled trial	Patients with confirmed acute chikungunya infection by RT-PCR and seroconversion between day 1 and 16 of illness	Group 1: CQ 600mg/d (base) for 3 days, 300mg/d for 2 days (n=27) Group 2: Placebo (n=27)	No statistically significant difference in fever clearance time or viraemia clearance time	Patients in CQ group were more likely to complain of persistent symptoms at day 200. However, they might have had more severe disease at enrolment as revealed later by an analysis of inflammatory markers of stored blood samples – high risk of selection bias
<i>COVID-19</i>					
Chen et al. China. 2020(36)	Randomized controlled clinical trial	Adult patients with virologically confirmed (RT-PCR) mild COVID-19 infection	Group 1: HCQ 400mg/d for 5 days plus “standard treatment” (n=15) Group 2: “Standard treatment” (n=15)	No statistically significant difference in clearance of viraemia by day 7, fever clearance time or total duration of hospitalization	Article is published in Mandarin. Only English abstract was reviewed Unable to assess risk of bias
Gautret et al. France. 2020(35)	Open label, non-randomized, controlled trial	Adult patients with virologically confirmed (RT-PCR) COVID-19 infection as test group Similar patients treated at a different institution or those refusing HCQ treatment at the same institute	Group 1: HCQ 600mg/d for 10 days (n=26) Group 2: No HCQ treatment (n=16)	Statistically significant rate of clearance of viraemia by day 6 of illness (70% in HCQ group vs. 12.5% in placebo group, p=0.001)	Serious risks of bias due to open label design, small sample size, non-random allocation and confounding effects due to recruiting control patients from a different institution, azithromycin administration in some patients only and baseline age difference in test and control groups
Chen et al. China. 2020(38)	Randomized double blind controlled clinical trial	Adult patients with virologically confirmed (RT-PCR) mild COVID-19 infection	Group 1: HCQ 400mg/d for 5 days plus “standard treatment” (n=31) Group 2: “Standard treatment” (n=31)	Statistically significant faster fever recovery and cough relief in HCQ group	This was a non-peer reviewed preprint High risk of reporting bias- The details of statistical interpretation and effect size are not explained. The “Standard treatment” according to authors included antibiotics, antivirals, steroids and immunoglobulins- Until more details are provided, the results cannot be interpreted

Tang et al. China. 2020(37)	Open label randomized trial	Adult patients with virologically confirmed (RT-PCR) mostly mild COVID-19 infection	Group 1: HCQ 200mg/d for 3 days followed by 800mg/d for 14-21 days + supportive care Group 2: supportive care only	No statistically significant difference in the proportion of aviraemic patients by day 28, time to aviraemia, or symptom resolution by day 28	This was a non-peer reviewed preprint High risk of performance and detection bias
Huang et al. China 2020(40)	Randomized controlled clinical trial (open label?)	Adult patients with virologically confirmed (RT-PCR) COVID-19 infection	Group 1: CQ 600mg base/d for 10 days Group 2: Lopinavir/Ritonavir 400/100 mg orally twice daily for 10 days	No statistically significant difference in viraemia clearance by day 14 Faster radiological resolution and reduced hospital stay in CQ group	It is not clear if this was a double- blind study and if not, there is a high risk of detection and performance bias. Method of sample size calculation is unclear
Borba et al. Brazil 2020(41)	Randomized controlled double-blind study	Adult patients with clinically suspected COVID- 19 infection	Group 1: CQ 1200mg base /d for 10 days Group 2: CQ 900mg base on day 1 and 450mg base/d for 4 days	Significantly higher mortality in high dose group by day 13 (15% vs 39%)	Study terminated due to safety concerns – risk of bias not evaluated

*Risk of bias not assessed for uncontrolled studies as results from these studies were not used to grade evidence. For non-randomized trials with more than one intervention, risk of bias assessed with ROBINS-I tool

Table 3. Clinical trials / studies in progression on using CQ or HCQ as antiviral agents

Name*	Reference ID	Source
Post-exposure Prophylaxis for SARS-Coronavirus-2	NCT04308668	ClinicalTrials.gov
Comparison of Lopinavir/Ritonavir or Hydroxychloroquine in Patients With Mild Coronavirus Disease (COVID-19)	NCT04307693	ClinicalTrials.gov
Chloroquine Prevention of Coronavirus Disease (COVID-19) in the Healthcare Setting	NCT04303507	ClinicalTrials.gov
Treatment of Mild Cases and Chemoprophylaxis of Contacts as Prevention of the COVID-19 Epidemic	NCT04304053	ClinicalTrials.gov
Various Combination of Protease Inhibitors, Oseltamivir, Favipiravir, and Chloroquine for Treatment of COVID19 : a Randomized Control Trial	NCT04303299	ClinicalTrials.gov
New Treatment for Radical Cure of Dengue Fever with Antiviral and Anti-cytokine	CTRI/2017/12/010834	WHO ICTRP
A prospective, open label, randomized, control trial for chloroquine or hydroxychloroquine in patients with mild and common novel coronavirus pulmonary (COVID-19)	ChiCTR2000030054	Chinese Clinical Trials Registry
A prospective, randomized, open label, controlled trial for chloroquine and hydroxychloroquine in patients with severe novel coronavirus pneumonia (COVID-19)	ChiCTR2000029992	Chinese Clinical Trials Registry
Evaluation the Efficacy and Safety of Hydroxychloroquine Sulfate in Comparison with Phosphate Chloroquine in Mild and Common Patients with Novel Coronavirus Pneumonia (COVID-19): a Randomized, Open-label, Parallel, Controlled Trial	ChiCTR2000029899	Chinese Clinical Trials Registry
Evaluation the Efficacy and Safety of Hydroxychloroquine Sulfate in Comparison with Phosphate Chloroquine in Severe Patients with Novel Coronavirus Pneumonia (COVID-19): a Randomized, Open-Label, Parallel, Controlled Trial	ChiCTR2000029898	Chinese Clinical Trials Registry
A prospective, randomized, open-label, controlled clinical study to evaluate the preventive effect of hydroxychloroquine on close contacts after exposure to the Novel Coronavirus Pneumonia (COVID-19)	ChiCTR2000029803	Chinese Clinical Trials Registry
A multicenter, single-blind, randomized controlled clinical trial for chloroquine phosphate in the treatment	ChiCTR2000031204	Chinese Clinical Trials Registry

of novel coronavirus pneumonia (COVID-19)

A Randomized Controlled Trial for Favipiravir Tablets Combine With Chloroquine Phosphate in the Treatment of Novel Coronavirus Pneumonia (COVID-19)	ChiCTR2000030987	Chinese Clinical Trials Registry
Randomized controlled trial for Chloroquine Phosphate in the Treatment of novel coronavirus pneumonia (COVID-19)	ChiCTR2000030718	Chinese Clinical Trials Registry
Clinical Study of Chloroquine Phosphate in the Treatment of Severe Novel Coronavirus Pneumonia (COVID-19)	ChiCTR2000029988	Chinese Clinical Trials Registry
Single arm study for exploration of chloroquine phosphate aerosol inhalation in the treatment of novel coronavirus pneumonia (COVID-19)	ChiCTR2000029975	Chinese Clinical Trials Registry
A Single-blind, Randomized, Controlled Clinical Trial for Chloroquine Phosphate in the treatment of Novel Coronavirus Pneumonia 2019 (COVID-19)	ChiCTR2000029939	Chinese Clinical Trials Registry
A Single-arm Clinical Trial for Chloroquine Phosphate in the treatment of Novel Coronavirus Pneumonia 2019 (COVID-19)	ChiCTR2000029935	Chinese Clinical Trials Registry
Efficacy of Chloroquine and Lopinavir/ Ritonavir in mild/general novel coronavirus (CoVID-19) infections: a prospective, open-label, multicenter randomized controlled clinical study	ChiCTR2000029741	Chinese Clinical Trials Registry
A prospective, open-label, multiple-center study for the efficacy of chloroquine phosphate in patients with novel coronavirus pneumonia (COVID-19)	ChiCTR2000029609	Chinese Clinical Trials Registry
Study for the efficacy of chloroquine in patients with novel coronavirus pneumonia (COVID-19)	ChiCTR2000029542	Chinese Clinical Trials Registry


*The names were extracted as they appeared on relevant databases without any language corrections

Figures

Figure 1. PRISMA flow diagram of study selection process

Figure 2. Risk of bias summary for randomized clinical trials

Journal Pre-proof



	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Borges 2013	+	?	+	+	-	+	-
Chen 2020	+	?	+	+	+	-	-
De Lamballerie 2008	+	?	+	+	+	+	-
Engchanil 2006	+	?	-	-	-	+	?
Helal 2016	+	-	-	-	+	+	
Huang 2020	+	?	?	?	+	+	-
Jacobson 2016	+	?	+	+	-	+	?
Murray 2010	+	?	+	+	+	+	-
Paton 2012	+	+	+	+	-	+	+
Peymani 2016	+	?	+	+		+	-
Sperber 1995	?	?	+	+	+	+	+
Sperber 1997	?	?	+	+	+	+	+
Tang 2020	+	+	-	-	+	+	+
Tricou 2010	+	+	+	+	+	+	+