



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof

Acute respiratory distress syndrome due to SARS-CoV-2 and Influenza A co-infection in an Italian patient: mini-review of the literature

D'Abramo Alessandra, Luciana Lepore, Claudia Palazzolo, Filippo Barreca, Liuzzi Giuseppina, Lalle Eleonora, Nicastri Emanuele



PII: S1201-9712(20)30491-4

DOI: <https://doi.org/10.1016/j.ijid.2020.06.056>

Reference: IJID 4361

To appear in: *International Journal of Infectious Diseases*

Received Date: 23 April 2020

Revised Date: 13 June 2020

Accepted Date: 16 June 2020

Please cite this article as: Alessandra D, Lepore L, Palazzolo C, Barreca F, Giuseppina L, Eleonora L, Emanuele N, Acute respiratory distress syndrome due to SARS-CoV-2 and Influenza A co-infection in an Italian patient: mini-review of the literature, *International Journal of Infectious Diseases* (2020), doi: <https://doi.org/10.1016/j.ijid.2020.06.056>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Title

Acute respiratory distress syndrome due to SARS-CoV-2 and Influenza A co-infection in an Italian patient: mini-review of the literature.

Authors

D'Abramo Alessandra^a, Luciana Lepore^a, Palazzolo Claudia^a, Filippo Barreca^b, Liuzzi Giuseppina^a,
Lalle Eleonora^a, Nicastri Emanuele^a.

Affiliations

^a National Institute for Infectious Diseases 'Lazzaro Spallanzani', IRCCS, via Portuense 292, 00149, Rome, Italy ^b Clinical Infectious Diseases, Department of System Medicine, Tor Vergata University, viale Oxford 81, 00133, Rome, Italy

Drs D'Abramo and Lepore equally contributed

Corresponding Author: Claudia Palazzolo, National Institute for Infectious Diseases IRCCS Diseases 'Lazzaro Spallanzani', IRCCS, Via Portuense 292, 00149, Rome, Italy nr tel/fax email: +3906551702374/+390655170204/claudia.palazzolo@inmi.i

Highlights

- Acute respiratory distress syndrome due to SARS-CoV-2 and Influenza A co-infection
- Challenging in co-infection cases identification
- Improvement of preventive measures and patients' clinical outcome

Abstract

A case of acute respiratory distress syndrome due to SARS-CoV-2 and Influenza A co-infection and a mini-review of the literature is reported. Even in COVID-19 epidemics, the early identification of concurrent respiratory pathogens is important to improve etiological diagnosis, preventive measures and patients' clinical management and outcome.

Keywords

Acute respiratory distress syndrome; SARS-CoV-2; Influenza A, COVID-19 Epidemic

Introduction

In December 2019, a novel coronavirus, the severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) which causes a human disease named coronavirus disease (COVID-19) was identified in the pneumonia outbreaks in Wuhan, China, in December 2019 (Chan et al, 2020). It is currently expanding rapidly to several countries all-around the world, on February 21 the first person-to-person transmission in Italy was reported (Spina et al, 2020). Here, we report a case of SARS-CoV-2 and Influenza A co-infection and a mini-review of the literature.

Case Presentation

A 56 year-old male general surgeon was admitted to the Lazzaro Spallanzani National Institute for

Infectious Diseases in Rome, Italy, on March 6, 2020. He was a smoker, within the overweight range (29-body mass index), with a history of two episodes of acute myocardial infarction treated with coronary angioplasty and stenting. For 4 days, he had been complaining of fever, diarrhea and asthenia after winter ski week in Northern Italy. Two days after the onset of symptoms, a nasopharyngeal swab was positive for SARS-CoV-2 (genes E and S) and Influenza A. On March 6, a chest computed tomography (CT) scan revealed bilateral and multiples peripheral ground glass opacities. Blood tests showed lymphopenia (lymphocyte and monocyte cell count: 0.67 and $0.09 \times 10^9/L$ respectively), C- reactive protein and serum fibrinogen levels were increased (43.3 g/L and 7980 g/L respectively). An arterial oxygen tension (PaO_2)/ fractional inspired oxygen (FiO_2) P/F Ratio was 320. Oral oseltamivir (75 mg twice per day for 5 days) and lopinavir/ritonavir (400/100 mg twice per days for 14 days) were started together with antibiotic therapy (intravenous ceftriaxone 2 gr and oral azithromycin 500 mg per day) and intravenous methylprednisolone (40 mg twice daily for 5 days with tapered discontinuation). On day eight of hospitalization, he developed respiratory failure (P/F Ratio dropped to 202) and a second chest CT scan showed a worsening of the bilateral ground-glass opacities with fibrotic consolidation in both lower pulmonary lobes (Fig1a). At that time, nasopharyngeal swabs were positive for SARS-CoV-2, only. The patient was transferred to the Intensive Care Unit (ICU) and non-invasive ventilation through continuous positive airway pressure (C-PAP) mask was started with positive end-expiratory pressure (PEEP) 7.5 mmHg and FiO_2 40%. After three days, he was re-admitted to the High Isolation Unit and discharged in good clinical conditions with persistently negative nasopharyngeal swabs. A SARS-CoV-2 serology by an in house indirect immunofluorescence assay (IgA 1:1280, IgM 1:320 and IgG 1:80) was positive (Colavita et al, 2019). During the following week, low-grade fever re-occurred with nocturnal sweats, a third CT scan showed a reduction of the ground glass areas with residual interstitial damage (Fig1b) and a further nasopharyngeal swab was negative. Symptoms healed spontaneously few days later.

Discussion

Previous cases of viral pneumonia SARS-CoV-2 and influenza coinfection have been reported in literature. During the new coronavirus epidemic, a total number of 37 cases were described. Table 1 summarizes the characteristics of the coinfecting patients. Fourteen cases belonged to epidemiological studies and clinical data were not available. All patients had a similar clinical presentation (fever, cough and shortness of breath) and 9 of them (9/37, 24.3%) presented a progressive worsening with ARDS. Furthermore, six patients needed ICU monitoring and were subsequently discharged in good clinical conditions with the exception of three patients who died.

Even during a pandemic scenario, several respiratory pathogens should be considered in the diagnostic algorithm, for an early etiological identification and appropriate treatment. SARS-CoV-2 and influenza viruses share common route of transmission, same season occurrence and overlapping clinical features (Lai et al, 2020, Chow et al, 2019). Indeed, SARS-CoV-2 exhibits prevalent human-to-human transmission through close contact with an estimated R_0 of 3.28 and a median of 2.79 with IQR of 1.6 (Liu et al, 2020).

Respiratory symptoms are always the initial manifestations of both SARS-CoV-2 and influenza infections which could progress towards ARDS. Recently, ground-glass opacities and a higher median PaO_2/FIO_2 (198.2 vs 107.0) were observed in COVID-19-induced ARDS rather than H1N1 patients (Tang et al, 2020).

Nevertheless, a timely identification of the two co-infections is needed in relation to difference in treatments and prognosis. Antiviral therapy is currently available for influenza infection (i.e. oseltamivir, zanamivir, and peramivir) while experimental off-label drugs (i.e. lopinavir/ritonavir, chloroquine, and hydroxychloroquine) have been commonly used in COVID-19 treatment. In particular, the boosted

protease inhibitor lopinavir/ritonavir has been previously associated with significantly fewer adverse clinical outcomes for the treatment of SARS and furthermore, in association with the Interferon Beta-1b, it has been demonstrated to be beneficial in animal studies against the Middle East Respiratory Syndrome (Chu et al, 2004, Chan et al 2015). Considering the severity of the clinical picture and the prompt availability of lopinavir/ritonavir in our Institute, clinicians opted for this treatment although the antiviral effects remain to be determined (Cao et al, 2020).

Despite a recent report on beneficial effect of steroids treatment in COVID-19 patients who develop ARDS, its routinely use remains controversial with lack of an accurate assessment of the harm/benefit balance (Wu et al, 2020).

The epidemiological situation in Italy in February 2020 and the recommendations in use at that time allowed a timely identification of our patient with a prompt hospitalization and subsequent diagnostic investigations. Therefore, the early antiviral treatments of both influenza and SARS-CoV-2, together with a brief steroid course and oxygen supplementation, had an impact on the patient's outcome avoiding the progressive worsening and the evolution towards the severe ARDS phase. In conclusion, even in epidemic setting, the early and prompt identification of concurrent respiratory pathogens is important in order to improve etiological diagnosis, preventive measures and patients' clinical management and outcome. Further studies are needed to better understand the pathogenic role of viral coinfections in respiratory diseases.

Declaration of interests

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Funding

This paper was funded by Ricerca Corrente Line1 on emerging and re-emerging infections funded by the Italian Ministry of Health.

Ethical approval

This study was approved by the Spallanzani Institute Ethical Board and patient's written informed consent for publication was collected

Conflict of interest

All authors have no conflict of interest to declare.

References

1. Azekawa S, Namkoong H, Mitamura K, Kawaoka Y, Saito F (2020). Co-infection with SARS-CoV-2 and influenza A virus. *IDCases*. 2020 Apr 21;20:e00775. doi: 10.1016/j.idcr.2020.e00775.
2. Blasco ML, Buesa J, Colomina J, Forner MJ, Galindo MJ, Navarro J, et al. Co-detection of Respiratory Pathogens in Patients Hospitalized With Coronavirus Viral disease-2019 Pneumonia. *J Med Virol*. 2020 Apr 22. doi: 10.1002/jmv.25922.
3. Cao B, Wang Y, Wen D, Liu W, Wang J, Fan G, et al. A trial of lopinavir–ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med*. 2020 May 7;382(19):1787-1799. doi: 10.1056/NEJMoa2001282.
4. Chan JF, Yao Y, Yeung ML, Deng W, Bao L, Jia L, et al. Treatment with lopinavir/ritonavir or

- interferon-beta1b improves outcome of MERS-CoV infection in a nonhuman primate model of common marmoset. *J Infect Dis.* 2015;212(12):1904–13.
5. Chan JF, Kok KH, Zhu Z, Chu H, To KK, Yuan S, et al. Genomic characterization of the 2019 novel human-pathogenic coronavirus isolated from a patient with atypical pneumonia after visiting Wuhan. *Emerg Microbes Infect.* 2020 Jan 28;9(1):221-236. doi: 10.1080/22221751.2020.1719902.
 6. Chow EJ, Doyle JD, Uyeki TM. Influenza virus-related critical illness: prevention, diagnosis, treatment. *Crit Care.* 2019 Jun 12;23(1):214. doi: 10.1186/s13054-019-2491-9.
 7. Chu CM, Cheng VC, Hung IF, Wong MM, Chan KH, Chan KS, et al. Role of lopinavir/ritonavir in the treatment of SARS: initial virological and clinical findings. *Thorax.* 2004;59(3):252–6
 8. Colavita F, Biava M, Castilletti C, et al. Inflammatory and Humoral Immune Response during Ebola Virus Infection in Survivor and Fatal Cases Occurred in Sierra Leone during the 2014–2016 Outbreak in West Africa. *Viruses* **2019**; 11:373. Available at: <https://www.mdpi.com/1999-4915/11/4/373>
 9. Cuadrado-Payán E, Montagud-Marrahi E, Torres-Elorza M, Bodro M, Blasco M, Poch E, et al. (2020). SARS-CoV-2 and influenza virus co-infection. *Lancet.* 2020 May 16;395(10236):e84. doi: 10.1016/S0140-6736(20)31052-7.
 10. de Souza Luna LK, Perosa DAH, Conte DD, Carvalho JMA, Alves VRG, Cruz JS, et al. Different Patterns of Influenza A and B Detected During Early Stages of COVID-19 in a University Hospital in São Paulo, Brazil. *J Infect.* 2020 May 21;S0163-4453(20)30313-3. doi: 10.1016/j.jinf.2020.05.036.
 11. Ding Q, Lu P, Fan Y, Xia Y, Liu M. The clinical characteristics of pneumonia patients coinfecting with 2019 novel coronavirus and influenza virus in Wuhan, China. *J Med Virol.* 2020 Mar 20. doi: 10.1002/jmv.25781.

12. Garazzino S, Montagnani C, Donà D, Meini A, Felici E, Vergine G, et al. Multicentre Italian study of SARS-CoV-2 infection in children and adolescents, preliminary data as at 10 April 2020. *Euro Surveill.* 2020 May;25(18):2000600. doi: 10.2807/1560-7917.ES.2020.25.18.2000600.
13. Hashemi SA, Safamanesh S, Ghafouri M, Taghavi MR, Mohajer Zadeh Heydari MS, Namdar Ahmadabad H, et al. Co-infection with COVID-19 and influenza A virus in two died patients with acute respiratory syndrome, Bojnurd, Iran. *J Med Virol.* 2020 May 15;10.1002/jmv.26014. doi: 10.1002/jmv.26014.
14. Khodamoradi Z, Moghadami M, Lotfi M. Co-infection of Coronavirus Disease 2019 and Influenza A: A Report from Iran. *Arch Iran Med.* 2020 Apr 1;23(4):239-243. doi: 10.34172/aim.2020.04.
15. Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of Co-infection Between SARS-CoV-2 and Other Respiratory Pathogens. *JAMA.* 2020 Apr 15;323(20):2085-2086. doi: 10.1001/jama.2020.6266
16. Konala VM, Adapa S, Gayam V, Naramala S, Daggubati SR, Kammari CB, et al. Co-infection with Influenza A and COVID-19. *Eur J Case Rep Intern Med.* 2020 Apr 20;7(5):001656. doi: 10.12890/2020_001656.
17. Konala VM, Adapa S, Naramala S, Chenna A, Lamichhane S, Garlapati PR, et al. A Case Series of Patients Coinfected With Influenza and COVID-19. *J Investig Med High Impact Case Rep.* Jan-Dec 2020;8:2324709620934674. doi: 10.1177/2324709620934674.
18. Lai CC, Liu YH, Wang CY, Wang YH, Hsueh SC, Yen MY, et al. Asymptomatic carrier state, acute respiratory disease, and pneumonia due to severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2): Facts and myths. *J Microbiol Immunol Infect.* 2020 Mar 4. pii: S1684-1182(20)30040-2. doi: 10.1016/j.jmii.2020.02.012.

19. Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al. Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus-Infected Pneumonia. *N Engl J Med*. 2020 Mar 26;382(13):1199-1207. doi: 10.1056/NEJMoa2001316.
20. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *J Travel Med*. 2020. 27:taaa02. doi: 10.1093/jtm/taaa021
21. Nowak MD, Sordillo EM, Gitman MR, Paniz Mondolfi AE. Co-infection in SARS-CoV-2 Infected Patients: Where Are Influenza Virus and Rhinovirus/Enterovirus? *J Med Virol*. 2020 Apr 30;10.1002/jmv.25953. doi: 10.1002/jmv.25953.
22. Pongpirul WA, Mott JA, Woodring JV, Uyeki TM, MacArthur JR, Vachiraphan A, et al. Clinical Characteristics of Patients Hospitalized with Coronavirus Disease, Thailand. *Emerg Infect Dis*. 2020 Apr 8;26(7). doi: 10.3201/eid2607.200598.
23. Richardson S, Hirsch JS, Narasimhan M, Crawford JM, McGinn T, Davidson KW, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA*. 2020 Apr 22;323(20):2052-2059. doi: 10.1001/jama.2020.6775
24. Spina S, Marrazzo F, Migliari M, Stucchi R, Sforza A, Fumagalli R. Lancet. The response of Milan's Emergency Medical System to the COVID-19 outbreak in Italy. 2020 Mar 14;395(10227):e49-e50. doi: 10.1016/S0140-6736(20)30493-1.
25. Tang X, Du R, Wang R, Cao T, Guan L, Yang C, et al. Comparison of Hospitalized Patients With ARDS Caused by COVID-19 and H1N. *Chest*. 2020 Mar 26. pii: S0012-3692(20)30558-4. doi: 10.1016/j.chest.2020.03.032.
26. Wehl G, Laible M, Rauchenzauner M. Co-infection of SARS CoV-2 and Influenza A in a Pediatric Patient in Germany *Klin Padiatr* 2020 May 11. doi: 10.1055/a-1163-7385. Online ahead of print.

27. Wu C, Chen X, Cai Y, Xia J, Zhou X, Xu S, et al. Risk Factors Associated With Acute Respiratory Distress Syndrome and Death in Patients With Coronavirus Disease 2019 Pneumonia in Wuhan, China. *JAMA Intern Med.* 2020 Mar 13. doi: 10.1001/jamainternmed.2020.0994.
28. Wu Q, Xing Y, Shi L, Li W, Gao Y, Pan S, et al. Coinfection and Other Clinical Characteristics of COVID-19 in Children. *Pediatrics.* 2020 May 6:e20200961. doi: 10.1542/peds.2020-0961
29. Wu X, Cai Y, Huang X, Yu X, Zhao L, Wang F, et al. Co-infection with SARS-CoV-2 and Influenza A Virus in Patient with Pneumonia, China. *Emerg Infect Dis.* 2020 Jun;26(6):1324-1326. doi: 10.3201/eid2606.200299.
30. Zhu X, Ge Y, Wu T, Zhao K, Chen Y, Wu B, et al. Co-infection with respiratory pathogens among COVID-2019 cases. *Virus Res.* 2020 May 11;285:198005. doi: 10.1016/j.virusres.2020.198005.

Figure Legend

Title

Chest computer tomography scan at patient's worsening and recovery

Legend

- (a) bilateral ground-glass opacities with fibrotic consolidation in both lower pulmonary lobes (b) reduction of the ground glass areas with residual interstitial damage.




Table. Clinical Characteristics of SARS-CoV-2 and Influenza Coinfection, COVID-19 Epidemic

References	Sex	Age	Comorbidities	Coinfection	Oseltamivir	Antivirals	Glucorticoids	ARDS	ICU	NIV	IMV	Outcome
Azekawa S et al, 2020	F	78	Dyslipidemia, hypothyroidism	Influenza A	Yes	No	No	No	No	No	No	Survived
Blasco ML et al, 2020	NA	NA	NA	Influenza A and RSV	NA	NA	NA	NA	NA	NA	NA	NA
Cuadrado-Payán et al, 2020	M	53	ERSD	Influenza A	Yes	Yes	NA	Yes	Yes	No	Yes	Survived
	M	78	T2DM	Influenza A	Yes	Yes	NA	Yes	Yes	No	Yes	Survived
	M	56	T2DM	Influenza A and B	No	No	NA	No	No	No	No	Survived
	F	81	ERSD	Influenza B	Yes	Yes	NA	Yes	Yes	No	Yes	Survived
De Souza Luna et al, 2020	NA	36	NA	Influenza B	NA	NA	NA	NA	NA	NA	NA	NA
Ding et al., 2020	F	47	None	Influenza A	Yes	Yes	Yes	No	No	No	No	Survived
	M	50	Hypertension, cancer	Influenza A	Yes	Yes	Yes	Yes	No	Yes	No	Survived
	F	66	Hypertension, CVD, HBV	Influenza B	Yes	Yes	No	No	No	No	No	Survived
	M	39	HBV	Influenza B	Yes	Yes	Yes	No	No	No	No	Survived
	F	49	None	Influenza A	Yes	Yes	No	No	No	No	No	Survived
Garazzino S et al, 2020	NA	<17	NA	Influenza A	NA	NA	NA	NA	NA	NA	NA	NA
Hashemi SA et al, 2020	F	78	NA	Influenza A	Yes	Yes	NA	Yes	NA	NA	NA	Dead
	M	75	NA	Influenza A	Yes	Yes	NA	Yes	NA	NA	NA	Dead
Khodamoradi et al., 2020	F	74	Hypertension, CVD	Influenza A	Yes	Yes	No	No	No	No	No	Survived
	M	40	None	Influenza A	Yes	Yes	No	No	No	No	No	Survived
	M	64	None	Influenza A	Yes	Yes	No	No	No	No	No	Survived
	M	50	None	Influenza A	Yes	Yes	No	No	No	No	No	Survived
Kim D et al, 2020	NA	NA	NA	Influenza A	NA	NA	NA	NA	NA	NA	NA	NA
Konala VM et al, 2020	M	57	hypertension, T2DM, CVD, AICD	Influenza A	Yes	Yes	No	No	No	No	No	Survived
	F	35	sickle cell trait	Influenza A	Yes	Yes	No	No	No	No	No	Survived
	F	68	T2DM, hypertension, GERD	Influenza B	Yes	Yes	NA	Yes	Yes	No	Yes	Dead
Konala VM et al, 2020	F	66	T2DM, CVD, hypertension, CKD	Influenza A	Yes	Yes	NA	Yes	Yes	No	Yes	Survived
Nowak MD et al, 2020	NA	NA	NA	Influenza A	NA	NA	NA	NA	NA	NA	NA	NA
Pongpirul WA et al, 2020	M	61	None	Influenza A	Yes	No	No	No	No	No	No	Survived

Richardson S et al, 2020	NA	NA	NA	Influenza A	NA	NA	NA	NA	NA	NA	NA	NA
Wehl G et al, 2020	NA	4 months	None	Influenza A	Yes	No	No	No	No	No	No	Survived
Wu Q et al, 2020	NA	<17	NA	MP, Influenza A&B, RSV	NA	NA	NA	NA	NA	NA	NA	NA
Wu X et al, 2020	M	69	None	Influenza A	Yes	NA	NA	Yes	Yes	No	Yes	Survived
Zhu X et al, 2020	NA	NA	NA	2 pts Influenza A 5 pts Influenza B	NA	NA	NA	NA	NA	NA	NA	NA

Legend: SARS-CoV-2: severe acute respiratory syndrome coronavirus-2, COVID-19: coronavirus disease-19, CVD: cardiovascular disease, HBV: hepatitis B virus, NA: not available, ARDS: Acute Respiratory Distress Syndrome, ICU: Intensive Care Unit, NIV: non-invasive ventilation, IMV: invasive mechanical ventilation, T2DM: type 2 diabetes mellitus. AICD: automatic implantable cardioverter defibrillator. RSV: Respiratory syncytial virus. MP: Mycoplasma pneumonia. ESRD: end-stage kidney disease. CKD: chronic kidney disease. GERD: gastroesophageal reflux disease