Universität Bonn

Notes for the lecture

Functional Analysis and PDE

held by

Prof. Dr. Stefan Müller

TEXed by

Daniel Emse

Winter 2024/25

Corrections and improvements

If you have corrections or improvements, contact me via (`daniel.emse@uni-bonn.de`).

Contents

Lecture **1**

0 Overview/Teaser **2**

1 Structures **3**
1.1 Topological spaces 3
1.2 Metric spaces 6
1.2.1 Definition and examples 6
1.2.2 Convergence and continuity in metric spaces 8
1.2.3 Completeness 9
1.3 Normed spaces 11
1.4 Hilbert spaces 12

2 Function spaces **14**
2.1 Spaces of bounded, continuous and differentiable functions 14

Appendix **A**

List of definitions **B**

List of statements **C**

Index **D**

Lecture

Chapter 0

Overview/Teaser

[09.10.2024, Lecture 1]

1. What is FA?
   - Analysis on infinite-dimensional spaces

     examples: $C^0(X), C^\infty(U \subseteq \mathbb{R}^n), \ell_p = \{a : \mathbb{N} \to \mathbb{R} \mid \sum_{n=0}^\infty |a_n|^p < \infty\}$ $(1 \leq p \leq \infty)$
   - Combination of geometric, analytic and (linear) algebraic ideas
2. What is FA good for?
   - Geometry/Topology with manifolds, bundles, loop space
   - PDEs, scalar products
   - Algorithms/Numerics (approxmiation theory; errors)
   - infinite-dimensional representations
   - Mathematical Physics / Quantum Mechanics; spectral theory
   - Probability Theory (infinite products)
3. Why combine PA & PDE?
   - $u : U \subseteq \mathbb{R}^n \to \mathbb{R}$ as a C^2 function:

     $$\Delta u(x) = \sum_{i=1}^n \frac{\partial^2 u}{\partial x_i^2}. \tag{0.1}$$

     with $-\Delta u = f$.

     Classical approach: Given a potential $f \in C^2$, does there exist a solution and is that solution unique?

     Modern approach: $\Delta : X \to Y$ linear map between function spaces. e.g. $X = C^2(U), Y = C^0(U)$. Existance of solutions for all $f \in C^0$ is equiv to surjectivity of $f\Delta$ and uniqueness is equivalent to injectivity.

     Fun fact: C^2 is a bad space for Δ. But Δ is a good operator on many function spaces, e.g. $C^{2,\alpha}$ with $0 < \alpha < 1$, so D^2u is α-Hölder continuous, or for $W^{2,p}$ as Sobolev spaces (where $D^2u \in L^p$ for $1 < p < \infty$), where D^2 is the weak derivative
4. „Big things“
   - Spaces (of founctions) and convergence (norm, scalarproduct)
   - (Bounded) Linear operators, Hahn-Banach theorem, uniform boundedness, ...
   - Hilbert spaces, projections
   - Spectral theory (LA: $A : \mathbb{R}^n \to \mathbb{R}^n$ symmetric, so $\langle Ax, y\rangle = \langle x, Ay\rangle \Rightarrow \exists$ orthogonal basis of eigenvectors; we want something similar in infinite dimensions which makes diagonazing easier, and we can easily compute the inverse or powers of the operator)

Chapter 1

Structures

Generally, all vector spaces will be with $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$.

- on „infinite-dimensional“ spaces
- Convergence and neighbourhoods, topological space
- Metric space (metric, triangle inequality, gives a *distance*)
- Normed space (vector space, $\|\lambda x\| = |\lambda| \cdot \|x\|$ for $\lambda \in \mathbb{R}$, gives a *length*)
- Scalar product space ($\|x\| = \sqrt{\langle x, x\rangle}$, gives a *length and an angle*)

1.1 Topological spaces

Let X a set and 2^X the power set.

Definition 1.1: Topological space, Topology

A *topological space* $(X, \mathcal{T})$ is a pair of a set X and subset $\mathcal{T}$ of 2^X with the following properties:

(*i*) $\emptyset \in \mathcal{T}, X \in \mathcal{T}$

(*ii*) finite intersections: $U, V \in \mathcal{T} \Rightarrow U \cap V \in \mathcal{T}$

(*iii*) arbitrary unions: $U_\alpha \in \mathcal{T}\ \forall \alpha \in A \Rightarrow \bigcup_{\alpha \in A} U_\alpha \in \mathcal{T}$.

The set $\mathcal{T}$ is called a *Topology on* X and the elements in $\mathcal{T}$ are often called *open sets*. A set C is called *closed* if $X \setminus C$ is open.
If $\mathcal{T}_1$ and $\mathcal{T}$ are topologies on X, then $\mathcal{T}_1$ is called *finer* than $\mathcal{T}_2$ or $\mathcal{T}_2$ is called *coarser* than $\mathcal{T}_1$ if $\mathcal{T}_1 \supset \mathcal{T}_2$.

Example 1.2. (*i*) The finest topology is $\mathcal{T} = 2^X$.

(*ii*) The coarsest topology is $\mathcal{T} = \{\emptyset, X\}$.

(*iii*) Standard topology on $\mathbb{R}$: $U \in \mathcal{T}_{st} \iff \forall x \in U \exists \varepsilon > 0 : (x - \varepsilon, x + \varepsilon) \subseteq U$. We will usually use this topology.

(*iv*) Let $X = \mathbb{R}$ and $\mathcal{T} = \{U \subseteq \mathbb{R} \mid U = \emptyset \text{ or } \mathbb{R} \setminus U \text{ is countable}\}$. Then, the closed sets are countable[1] or $\mathbb{R}$.

(*v*) Let $(X, \mathcal{T})$ be a topological space. Then, for $A \subseteq X$, the *relative topology* $\mathcal{T}_A$ on A is defined by $\mathcal{T}_A = \{U \cap A \mid U \in \mathcal{T}\}$.

(*vi*) Let B be a set and $\mathcal{T}_\beta$ a topology on X for all $\beta \in B$. Then, $\mathcal{T} := \bigcap_{\beta \in B} \beta$ defines a topology on X.

[1] We call a set *countable* if it is empty, finite or has an bijection to $\mathbb{N}$.

Given any $\mathcal{S} \subseteq 2^X$,

$$\bigcap_{\substack{\mathcal{T} \text{ top.} \\ \mathcal{T} \supseteq \mathcal{S}}} \mathcal{T} \tag{1.1}$$

is a topology and is called the *coarsest topology containing* $\mathcal{S}$.

(*vii*) *Product topology* of $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ as topological spaces is definied with

$$\mathcal{S} := \{U \times Y \mid U \in \mathcal{T}_X\} \cup \{X \times V \mid V \in \mathcal{T}_Y\}. \tag{1.2}$$

The product topology is $\mathcal{T}_{\mathcal{S}}$, the coarsest topology containing $\mathcal{S}$.

The product topology is the coarsest topology so that the projections $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$ are „continuous“.

Definition 1.3: Interior, Closure, Boundary

Let $(X, \mathcal{T})$ be a topological space and $A \subseteq X$. We define the *interior* A°, the *closure* $\overline{A}$ and the *boundary* ∂A by

$$A^\circ = \bigcup_{\substack{U \subseteq A \\ U \in \mathcal{T}}} U, \quad \overline{A} = \bigcap_{\substack{C \supseteq A \\ X \setminus C \in \mathcal{T}}} C, \quad \partial A = \overline{A} \setminus A^\circ. \tag{1.3}$$

Definition 1.4: Dense subsets, separable topological space

$A \subseteq X$ is *dense* if $\overline{A} = X$. The space X is *separable* if there exists a countable, dense subset.

Definition 1.5: Open neighbourhood

Let $x \in X$. Then, $U \subseteq X$ is an *open neighbourhood of* x if $U \in \mathcal{T}$ and $x \in U$.

Definition 1.6: Continuous maps, Homeomorphisms

Let $(X, \mathcal{T}_X), (Y, \mathcal{T}_Y)$ topological spaces. Then, $f : X \to Y$ is called *continuous* if

$$\forall V \supseteq \{x\} \in \mathcal{T}_Y : \quad f^{-1}(V) = \{x \in X \mid f(x) \in V\} \in \mathcal{T}_X. \tag{1.4}$$

We call f a *homeomorphism* if f is bijective and both f and f^{-1} are continuous.

If $\mathcal{T} = 2^X$, and $f : X \to Y$ is continuous.

If $\mathcal{T} = \{\emptyset, \mathcal{T}\}$, then every constant map is continuous.

[09.10.2024, Lecture 1]
[11.10.2024, Lecture 2]

Definition 1.7: Convergence

Let $(X, \mathcal{T})$ be a topological space and $x : \mathbb{N} \to X$ a sequence. Let $x^* \in X$. We say that x *converges to* x^* if for **every** neighbourhood U of x, the set $\{j \in \mathbb{N} \mid x_j \notin U\}$ is finite.

Example. (*i*) If $\mathcal{T} = \{\emptyset, X\}$, every sequence converges to every point $x^* \in X$.

(*ii*) If $\mathcal{T} = 2^X$, take $U = \{x^*\}$, so only sequences which are constant with up to finitely many exceptions converge.

Definition 1.8: Hausdorff space

A topological space is called a *Hausdorff space* if

(*i*) sets consisting of a single point are closed,

(*ii*) for different points $x \neq y$, there exist disjoint open neighbourhoods: $\exists U_x, U_y \in \mathcal{T}$ with $x \in U_x, y \in U_y$ and $U_x \cap U_y = \emptyset$.

Proposition 1.9. *If $(X, \mathcal{T})$ is Hausdorff, then every sequence has at most one limit point.* □

Definition 1.10: Compactness

A subset K of a topological space $(X, \mathcal{T})$ is *compact* if every open cover of K contains a **finite** subcover: If A is an index set and $U_\alpha \in \mathcal{T}$ for every $\alpha \in A$ and

$$\bigcup_{\alpha \in A} U_\alpha \supseteq K, \tag{1.5}$$

then there exists a **finite** set $A' \subseteq A$ so that

$$\bigcup_{\alpha \in A'} U_\alpha \supseteq K. \tag{1.6}$$

Lemma 1.11: Continuous maps compact to compact

If $f : (X, \mathcal{T}_X) \to (Y, \mathcal{T}_Y)$ is continuous and $K \subseteq X$ is compact in X, then $f(X)$ is compact in Y.

Proof. Follows directly from the definitions. □

Theorem 1.12: Continuous maps with compact domain attain max and min

If $f : (X, \mathcal{T}_X) \to \mathbb{R}$ is continuous and $K \subseteq X$ is compact, then f attains its minimum and maximum on K, i.e. $\exists \underline{x}, \overline{x} \in K$ so that $f(\underline{x}) = \inf_{x \in K} f(x), f(\overline{x}) = \sup_{x \in K} f(x)$.

Proof. $f(K) \subseteq \mathbb{R}$ is compact, the rest is easy. □

Definition 1.13: Connectedness

Let $(X, \mathcal{T})$ be a topological space. We call X *connected* if it cannot be written as a disjoint union of two open subsets. A subset $A \subseteq X$ is connected if $(A, \mathcal{T}_A)$ is connected.

Exercise 1. Show that intervals in $\mathbb{R}$ are connected.

Theorem 1.14: Intermediate value theorem

If $f : (X, \mathcal{T}_X) \to \mathbb{R}$ is continuous and $(X, \mathcal{T}_X)$ is connected, and there exist $a, b \in X$ so that $f(a) < 0, f(b) > 0$. Then, there exists an $x \in X$ with $f(x) = 0$.

Proof. Assume $f(x) \neq 0$ for all $x \in X$. We know that $U := f^{-1}((0, \infty))$ is open, and $X \setminus U = f^{-1}((-\infty, 0]) = f^{-1}((-\infty, 0))$ is also open. Then, $X = U \cup (X \setminus U) \Rightarrow X$ not connected. ↯ □

1.2 Metric spaces

1.2.1 Definition and examples

Let X be a set and $d : X \times X \to [0, \infty)$.

Definition 1.15: Metric space, metric

(X, d) is called a *metric space* and d is called *metric* if

(*i*) $d(x, y) = 0 \Leftrightarrow x = y$,

(*ii*) $\forall x, y : d(x, y) = d(y, x)$,

(*iii*) (triangle inequality) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in X$.[1]

If only (*ii*) and (*iii*) are fulfilled, we call d a *semimetric* or *pseudometric*, and with $x \sim y :\Leftrightarrow d(x, y) = 0$, we get a metric $\overline{d}$ on the equivalence classes.

[1] „Where the triangle inequality is not true is on airline flights. Sometimes, it is cheaper to fly from Köln/Bonn to Mallorca first before flying to mainland Spain later.“ – Stefan Müller, 11.10.2024

Lemma 1.16: Composition of concave map and metric is metric

Assume $\psi : [0, \infty) \to [0, \infty)$ is continuous, non-decreasing, ***concave*** *and $\psi(0) = 0, \psi(x) > 0$ for $x > 0$. If d is a metric on X, so is $\psi \circ d$.*

Proof. It is enough and easy to check that $\psi(a + b) \leq \psi(a) + \psi(b)$. □

Example. Define $\psi(t) := \frac{t}{1+t} = 1 - \frac{1}{1+t}$. Then, $\psi \in C^0([0, \infty)) \cap C^2((0, \infty))$. Non-decreasing and concave are equivalent to $\psi' \geq 0$ and $\psi'' \leq 0$.

Example. (*i*) Let $X = \mathbb{R}^n$ and $d_\infty(x, y) = \max_{1 \leq i \leq n} |x_i - y_i|$.

(*ii*) $X = \mathbb{R}^n, d_p(x, y) := (\sum_{i=1}^n |x_i - y_i|^p)^{\frac{1}{p}}$ for $1 \leq p < \infty$,

(*iii*) $\ell_p = \left\{ a : \mathbb{N} \to \mathbb{R} \;\middle|\; (\sum_{i=1}^\infty |x_i|^p)^{\frac{1}{p}} < \infty \right\}$ and $d_p(a, b) := (\sum_{i=1}^\infty |a_i - b_i|^p)^{\frac{1}{p}}$ for $1 \leq p < \infty$

(*iv*) $\ell_\infty := \{a : \mathbb{N} \to \mathbb{R} \mid \sup_{i \in \mathbb{N}} |a_i| < \infty\}$ with $d_\infty(a, b) := \sup_{i \in \mathbb{N}} |a_i - b_i|$

(*v*) $X = \mathbb{R}$ with $d(x, y) = \frac{|x-y|}{1+|x-y|} = \psi(|x - y|)$ for $\psi(t) = \frac{t}{1+t}$. By Lemma 1.16, d is a metric on $\mathbb{R}$. Fun fact: $0 \leq d(x, y) < 1$.

(*vi*) Metric on the space of all sequences:

$$d(a, b) = \sum_{i=0}^{\infty} 2^{-i} \frac{|a_i - b_i|}{1 + |a_i - b_i|} \leq \sum_{i=0}^{\infty} 2^{-i} \leq 2. \tag{1.7}$$

(*vii*) Let X be a set and (Y, d_Y) a metric space and $f : X \to Y$ **injective**. Then, we can define the *pullback metric* $d_X(x, y) := d_Y(f(x), f(y))$. d_X is a metric on X.

(*viii*) $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, \infty\}$. Then,

$$f : \overline{\mathbb{R}} \to [-1, 1], x \mapsto \begin{cases} -1 & x = -\infty \\ \frac{x}{1+|x|} & x \in \mathbb{R} \\ 1 & x = \infty \end{cases} \tag{1.8}$$

is bijective. Then, $d_{\overline{\mathbb{R}}}(x, y) = |f(x) - f(y)|$ is a pullback metric.

(*ix*) Shortest path on a graph

Remark. Let (X,d) be a metric space, let $A,B \subseteq X$, let $x \in X$ and let $r > 0$. Further distances:

Diameter of a subset

$$\sup\{d(x,y) \mid x,y \in A\} \tag{1.9}$$

(with the convention $\operatorname{diam}(A) = \infty$ if $\{d(x,y) \mid x,y \in A\}$ is unbounded)

Distance between two sets

$$\operatorname{dist}(A,B) = \inf\{d(x,y) \mid x \in A, y \in B\} \tag{1.10}$$

Distance between a point and a set

$$\operatorname{dist}(x,A) = \inf\{d(x,y) \mid y \in B\} = \operatorname{dist}(\{x\},A) \tag{1.11}$$

r-neighbourhood of a set

$$B_r(A) := \{x \in X \mid \operatorname{dist}(x,A) < r\} \tag{1.12}$$

r-ball of a point

$$B_r(x) = B(x,r) = \{y \in X \mid d(y,x) < r\} \tag{1.13}$$

Hausdorff distance of two sets

$$d_H(A,B) := \inf\{r > 0 \mid A \subseteq B_r(B) \text{ and } B \subseteq B_r(A)\} \tag{1.14}$$

and $d_H(\{x\},\{y\}) = d(x,y)$.

Remark. It is easy to show that

$$d_H(A,B) = \max\left(\sup_{x\in A}\operatorname{dist}(x,B), \sup_{y\in B}\operatorname{dist}(y,A)\right). \tag{1.15}$$

Proof. Exercise. □

Proposition 1.17 (Topology induced by a metric)**.** *Let (X,d) be a metric space. Let $\mathcal{T}_d$ consist of all sets U with the following property:*

$$\forall x \in U \exists \varepsilon > 0 : B_\varepsilon(x) \subseteq U. \tag{1.16}$$

Then, $\mathcal{T}_d$ is a topology on X.

Proof. Clear. For the intersection, take the ball with the smaller radius. □

Lemma 1.18: Hausdorff distance is metric on closed and bounded sets

Let (X,d) be a Hausdorff space. Then, d_H defines a metric on

$$\mathcal{C} := \{A \subseteq X \mid \emptyset \neq A \text{ is closed and bounded}\}. \tag{1.17}$$

Proof. Exercise. □

[11.10.2024, Lecture 2]
[16.10.2024, Lecture 3]

1.2.2 Convergence and continuity in metric spaces

Proposition 1.19. *Let (X,d) be a metric space and $x:\mathbb{N}\to X$ a sequence, and $x^*\in X$. Then, the following two statements are equivalent:*

(i) *x converges to x^* in $(X,\mathcal{T}_d)$,*

(ii) $\lim_{n\to\infty} d(x_n,x^*)=0$.

Moral. „In metric spaces, anything can be done by sequences“

Proposition 1.20. *Let (X,d) be a metric space and let $A\subseteq X$. Then, the following two statements are equivalent:*

(i) *A is closed, so $X\setminus A\in\mathcal{T}_d$,*

(ii) *A is* sequentially closed*, so for every sequence $x:\mathbb{N}\to A$ which converges to $x^*\in X$, one has $x^*\in A$.*

Moreover, the closure of any set A is given by the sequential closure*, this means*

$$\overline{A}=\{x^*\in X\mid \exists x:\mathbb{N}\to A, x \text{ converges to } x^*\}. \tag{1.18}$$

Proof. Easy exercise. □

Proposition 1.21. *Let $(X,d_X),(Y,d_Y)$ be two metric spaces and $f:X\to Y$. Then, the following three statements are equivalent:*

(i) *f is continuous in as a map $f:(X,\mathcal{T}_{d_X})\to(Y,\mathcal{T}_{d_Y})$,*

(ii) *For every $x\in X,\varepsilon>0$ there exists a $\delta>0$ so that $d_X(z,x)<\delta\Rightarrow d_Y(f(z),f(x))<\varepsilon$,*

(iii) *For every sequence x convergent to x^*, also $f(x)$ converges to $f(x^*)$.*

We know that every metric space is Hausdorff. *Question:* If $(X,\mathcal{T})$ is a Hausdorff space, does there exist a metric $d:X\times X\to[0,\infty)$ such that $\mathcal{T}=\mathcal{T}_d$? Or shorter, is X metrizable?

If X is second countable (so there exist countable neighbourhoods), X is metrizable, and then, X ist first countable.

Definition 1.22: Equivalent metrics, stronger metrics

Let d_1 and d_2 be metrics on X.
We say *d_1 is stronger than d_2* if $\mathcal{T}_{d_1}\supset\mathcal{T}_{d_2}$.
We say *d_1 is equivalent to d_2* if $\mathcal{T}_{d_1}=\mathcal{T}_{d_2}$.

Proposition 1.23. *The following statements are equivalent:*

(i) *d_1 is stronger than d_2,*

(ii) *The identity map $x\mapsto x$ for $(X,d_1)\to(X,d_2)$ is continuous,*

(iii) *Every sequence that converges in d_1 also converges in d_2,*

(iv) *$\forall x\in X\forall\varepsilon>0\exists\delta>0$ so that $d_1(x,y)<\delta\Rightarrow d_2(x,y)<\varepsilon$.*

Example.

(i) Let $X=\mathbb{R}$ and $d_1(x,y)=|x-y|, d_2=|\arctan(x)-\arctan(y)|$. Then, d_1 and d_2 are equivalent (Exercise 1.4).

(*ii*) Let $\ell_1 := \{x : \mathbb{N} \to \mathbb{R} \mid \sum_{k=0}^{\infty} |x_k| < \infty\}$ with $d_1(x,y) := \sum_{k=0}^{\infty} |x_k - y_k|$ and $d_\infty(x,y) = \sup_{k\in\mathbb{N}} |x_k - y_k| \le d_1(x,y)$. We see that d_1 is stronger than d_∞, but not equivalent: See the map

$$\mathbb{N} \to (\mathbb{N} \to \mathbb{R}), \quad j \mapsto x_k^{(j)} := \begin{cases} 0 & k < j-1 \text{ or } k > 2j \\ \frac{1}{k} & j \le k \le 2j. \end{cases} \tag{1.19}$$

Then, $\left|x_k^{(j)}\right| \le \frac{1}{j}$ with $d_\infty(x^{(j)}, 0) \le \frac{1}{j} \to 0$ as $j \to \infty$, but $d_1(x_k^{(j)}, 0) = \sum_{k=0}^{\infty} x_k^{(j)} = \sum_{k=j}^{2j} \frac{1}{k} \ge j\frac{1}{2j} = \frac{1}{2}$.

Definition 1.24: Separable metric space

A metric space (X,d) is called *separable* if $(X, \mathcal{T}_d)$ is separable.

1.2.3 Completeness

Definition 1.25: Cauchy sequence, completeness

Let (X,d) be a metric space.

(*i*) A sequence $x : \mathbb{N} \to X$ is a *Cauchy sequence* if $\forall \varepsilon > 0 \exists k_0$ so that for all $k, \ell \ge k_0$: $d(x_k, x_\ell) < \varepsilon$,

(*ii*) (X,d) is called *complete* if every Cauchy sequence converges.

Example.

(*i*) $\mathbb{Q}$ equipped with the standard metric is not complete.

(*ii*) (ℓ_p, d_p) are complete for all $1 \le p \le \infty$.

(*iii*) $\mathbb{R}$ equipped with the standard metric is complete, but not with the metric $\tilde{d}(x,y) := |\arctan(x) - \arctan(y)|$, although they induce the same topology (Exercise 1.4).

(*iv*) $\ell_1 = \{x : \mathbb{N} \to \mathbb{R} \mid \sum_{k=0}^{\infty} |x_k| < \infty\}$ with d_1 is complete (see (*ii*)), but (ℓ_1, d_∞) is not complete:

Take $j \mapsto x_k^{(j)}$ and set $x_k^{(j)} := \begin{cases} \frac{1}{k} & 1 \le j \le k \\ 0 & \text{else} \end{cases}$. For any $m > j$, we get $\left|x_k^{(j)} - x_k^{(m)}\right| \le \frac{1}{j}$, so $d_\infty(x^{(j)}, x^{(m)}) \le \frac{1}{j}$ for $m > j$, so the sequence is Cauchy but has no limit.

Proposition 1.26. *Let (X,d) be a complete metric space and $A \subseteq X$ closed. Let $d_A := d|_{A\times A}$. Then, (A, d_A) is complete.*

Proof. If $x : \mathbb{N} \to A$ is a Cauchy sequence in X, by completeness, $x_k \to x^*$ in X. As A is (sequentially) closed, $x^* \in A$. □

Definition 1.27: Isometric immersion, isometry

Let (X, d_X) and (Y, d_Y) be metric spaces. A map $f : X \to Y$ is called an *isometric immersion* if

$$d_Y(f(x), f(y)) = d_X(x,y) \tag{1.20}$$

for all $x, y \in X$. The map f is called an *isometry* if it is in addition bijective

Remark. An isometric immersion is automatically bijective. Thus, if f is an isometric immersion, than f is an isometry from X to $f(X)$. The spaces X and $f(X)$ (equipped with the

restriction of the metric d_Y) are indistinguishable as metric spaces. In particular, a sequence $f : \mathbb{N} \to \mathbb{R}$ is convergent in X if and only if $f \circ x$ is convergent in $f(X)$, and X is a Cauchy sequence in X if and only if $f \circ x$ is a Cauchy sequence in $f(X)$.

Theorem 1.28: Completion

Let (X, d) be a metric space. Then, there exists a complete, metric space $(\tilde{X}, \tilde{d})$ and an isometric immersion $j : X \to \tilde{X}$ such that $j(X)$ is dense in $\tilde{X}$.

The space $\tilde{X}$ is called the *completion of* X (and is unique up to isometries).

Proof. We will see a short proof later. Here we just sketch the standard proof which is based on considering the space of Cauchy sequences in X modulo converging sequences. This proof is modelled on the construction of the real numbers $\mathbb{R}$ from the rational numbers $\mathbb{Q}$ (see e.g. Alt's book for details).

Let

$$\hat{X} := \{x : \mathbb{N} \to X \mid x \text{ Cauchy sequence}\}. \tag{1.21}$$

Define

$$x \sim y \quad :\Longleftrightarrow \quad \lim_{k\to\infty} d(x_k, y_k) = 0. \tag{1.22}$$

It is easy to see that $\sim$ is an equivalence relation. The equivalence class of $x \in \hat{X}$ is defined by

$$[x] := \left\{ z \in \hat{X} \;\middle|\; z \sim x \right\}, \tag{1.23}$$

and the space of equivalences is defined by $\hat{X} := {}^{X}/_{\sim}$. On $\hat{X}$, we define

$$\tilde{d}([x], [y]) := \lim_{j\to\infty} d(x_j, y_j). \tag{1.24}$$

It follows from the definition of the equivalence classes and the triangle inequality that $\tilde{d}$ is well-defined, i.e., the right hand side only depends on the equivalence classes of the Cauchy sequences x and y. Since x and y are Cauchy sequences, it is also easy to see that $j \mapsto d(x_j, y_j)$ is a Cauchy sequence in $\mathbb{R}$ and that therefore $\lim_{j\to\infty} d(x_j, y_j)$ exists. It is easy to see that $\tilde{d}$ is a metric on $\tilde{X}$. Then, one shows that

- $(\tilde{X}, \tilde{d})$ is complete,
- the map $j : X \to \tilde{X}$ which maps a to the (equivalence class of the) constant sequence with value a is an isometry
- $j(X)$ is dense in $(\tilde{X}, \tilde{d})$

□

Lemma 1.29: Hausdorff metric space is complete

Let (X, d) be a complete, metric space. Then, the space $(\mathcal{C}, d_H)$ of closed, bounded, non-empty set with the Hausdorff metric (introduced in Lemma 1.18) is complete.

Proof. Exercise. *Hint:* First show that a decreasing Cauchy sequence of sets B_j converges to $B^* := \bigcap_{j\in\mathbb{N}} B_j$ (and in particular, $B^* \neq \emptyset$). By passage to a subsequence, you can assume without loss of generality that $d_H(B_j, B_{j+1}) < 2^{-j}$ (explain why). For a general Cauchy sequence A_j define $B_j := \overline{\bigcup_{k\geq j} A_j}$ and show that $d_H(A_j, B_j) \to 0$. □

One can also introduce a metric on set of (equivalence classes of) bounded and complete metric spaces, the so called *Gromov-Hausdorff metric*. Here we say that $(X, d_X) \sim (Y, d_Y)$ if there

exist an isometry $I : X \to Y$. Then, one defines

$$\begin{aligned} &d_{GH}((X, d_X), (Y, d_Y)) \\ &:= \inf \{ d_{H,Z}(I(X), J(Y)) \mid (Z, d_Z) \text{ metric space, } I : X \to Z, J : Y \to Z \text{ isom. immers.} \} \end{aligned} \tag{1.25}$$

where $d_{H,Z}$ denotes the Hausdorff distance in the metric space (Z, d_Z). The GH metric plays an important role in geometry as well as in application to image processing.

[16.10.2024, Lecture 3]
[18.10.2024, Lecture 4]

1.3 Normed spaces

Let X be a $\mathbb{K}$-vector space. Recall our convention that $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$.

Definition 1.30: Norm

A pair $(X, \|\cdot\|)$ is called *normed space* if X is a $\mathbb{K}$-vector space and $\|\cdot\| : X \to [0, \infty)$ with the following properties:

(*i*) Definiteness: $\|x\| = 0 \Rightarrow x = 0$

(*ii*) Homogeneity: $\|\lambda x\| = |\lambda| \cdot \|x\|$ for all $\lambda \in \mathbb{K}, x \in X$

(*iii*) Triangle inequality: $\|x + y\| \leq \|x\| + \|y\|$ for all $x, y \in X$.

If only (*ii*) and (*iii*) hold, we call $\|\cdot\|$ a *seminorm*.

If (*i*)$-$(*iii*) hold, the map d defined by $d(x, y) = \|x - y\|$ is a metric that is translation invariant.

It follows from the triangle inequality that

$$|\|x\| - \|y\|| \leq \|x - y\| \tag{1.26}$$

That implies, that the map $X \to \mathbb{R}, x \mapsto \|x\|$ is continuous (actually, Lipschits continuous with Lipschitz constant 1).

Definition 1.31: Banach space

A normed space which is complete is called *Banach space*.

Example. For $1 \leq p \leq \infty$, the sequence spaces ℓ_p are Banach spaces if equipped with norms $\|x\|_p := (\sum_{k=0}^{\infty} |x_k|^p)^{\frac{1}{p}}$ for $p < \infty$ and $\|x\|_\infty := \sup_{k \in \mathbb{N}} |x_k|$ respectively.

Definition 1.32: Stronger norms, equivalent norms

Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be norms on a $\mathbb{K}$-vector space X. We say that $\|\cdot\|_1$ is *stronger than* $\|\cdot\|_2$ if the corresponding metric d_1 is stronger than d_2.
We say that $\|\cdot\|_1$ and $\|\cdot\|_2$ *are equivalent* if d_1 and d_2 are equivalent.

Proposition 1.33. *Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be norms on a $\mathbb{K}$-vector space X. Then, $\|\cdot\|_1$ is stronger that $\|\cdot\|_2$ iff there exists a $C > 0$ such that*

$$\forall x \in X : \quad \|x\|_2 \leq C \cdot \|x\|_1 . \tag{1.27}$$

Moreover, $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent iff three exist $c, C > 0$ such that

$$\forall x \in X : \quad c \|x\|_1 \leq \|x\|_2 \leq C \|x\|_1 . \tag{1.28}$$

Proof. The second statement follows immediately from the first statement.

Recall that d_1 is stronger than d_2 iff the identity map from (X, d_1) to (X, d_2) is continuous. If (1.27) holds, the identity map is continuous.

Now assume that $\mathrm{id} : (X, d_1) \to (X, d_2)$ is continuous. Apply the ε-δ definition of continuity at the point $x = 0$:

$$\exists \delta > 0 : \quad \forall y : \quad \|y\|_1 < \delta \Rightarrow \|y\|_2 < 1. \tag{1.29}$$

Let $C > 0$ be any number such that $C > \frac{1}{\delta}$. Let $X \in X \setminus \{0\}$. Define $y := \frac{x}{C\|x\|_1} \Rightarrow \|y\|_1 = \frac{1}{C} < \delta$. This implies that $\|y\|_2 < 1$. Now, $x = C\|x\|_1 y$ and thus $\|x\|_2) C\|x\|_1 \|y\|_2 \leq C\|x\|_1$. □

Example. (*i*) On $\mathbb{R}^n$, all norms are equivalent. (Analysis II)

(*ii*) See $\ell_1 := \{x : \mathbb{N} \to \mathbb{R} \mid \sum_{k=1}^{\infty} |x_k| < \infty\}$ with $\|x\|_1 = \sum_{k=1}^{\infty}, |x_k|$ and $\|x\|_\infty = \sup_{k\in\mathbb{N}} |x_k|$. We have $\|x\|_\infty \leq \|x\|$, so $\|\cdot\|_1$ is stronger than $\|\cdot\|_\infty$ but $\|\cdot\|_\infty$ is not equivalent to $\|\cdot\|_1$.

1.4 Hilbert spaces

For $\mathbb{C} \ni \alpha = a + bi$ with $a, b \in \mathbb{R}$, we say $\overline{\alpha} := a - bi$ is the *complex conjugate*. We get $\alpha\overline{\alpha} = a^2 + b^2$.

Definition 1.34: Sesquilinear form

Let X be a $\mathbb{K}$-vector space. A map $X \times X \to \mathbb{K}, (x_1, x_2) \mapsto \langle x_1, x_2 \rangle_X$ is called a *sesquilinear form* if for all $x, y, x_1, x_2, y_1, y_2 \in X$ and for alle $\alpha \in \mathbb{K}$, we have

(*i*) $\langle \alpha x, y \rangle_X = \alpha \langle x, y \rangle_X$ and $\langle x, \alpha y \rangle_X = \overline{\alpha} \langle x, y \rangle_X$,

(*ii*) $\langle x_1 + x_2, y \rangle_X = \langle x_1, y \rangle_X + \langle x_2, y \rangle_X$, $\langle x, y_1 + y_2 \rangle_X = \langle x, y_1 \rangle_X + \langle x, y_2 \rangle_X$.

A sesquilinear form is *symmetric* if

$$\langle y, x \rangle_X = \langle x, y \rangle_X \qquad \forall x, y \in X. \tag{1.30}$$

A sesquilinear form is *positive semidefinite* if

$$\langle x, x \rangle_X \geq 0 \qquad \forall x \in X, \tag{1.31}$$

and *positive definite* if $\langle x, x \rangle_X \geq 0$ and $\langle x, x \rangle_= 0 \Rightarrow X = 0$ for all $x \in X$.

Example. (*i*) $X = \mathbb{C}$ and $\langle x, y \rangle = x\overline{y}$,

(*ii*) $X = \mathbb{C}^n$ and $\langle x, y \rangle = \sum_{i=1}^{n} x_i \overline{y_i}$,

(*iii*) $X = (\ell_2, \mathbb{C})$ and $\langle x, y \rangle = \sum_{i\in\mathbb{N}} x_i \overline{y_i}$.

Definition 1.35: Scalar product

A positive definite, symmetric sesquilinear form on a $\mathbb{K}$-vector space is a *Scalar product.*. Then, the space $(X, \langle \cdot, \cdot \rangle_X)$ is called a *Pre-Hilbert space*.

What are the key features of a scalar product?

- Geometrically: Orthogonal if $\langle x, y \rangle_X = 0$; orthogonal projections; Pythagoras
- Analytically: Cauchy-Schwarz inequality

Lemma 1.36: Scalar product traits

Let (X) be a $\mathbb{K}$-vector space and $\langle\cdot,\cdot\rangle_X$ a positive semidefinite, symmetric sesquilinear form and define

$$\|x\| := \sqrt{\langle x,x\rangle_X}. \tag{1.32}$$

Then,

(i) Homogeneity: $\|\alpha x\| = |\alpha| \cdot \|x\|$,

(ii) Cauchy-Schwarz inequality: $|\langle x,y\rangle| \leq \|x\| \cdot \|y\|$,

(iii) Triangle inequality: $\|x+y\| \leq \|x\| + \|y\|$,

(iv) Parallelogram identity: $\|x+y\|^2 + \|x-y\|^2 = 2(\|x\|^2 + \|x\|^2)$.

In particular, $\|\cdot\|$ is a seminorm. If, in addition, $\langle\cdot,\cdot\rangle_X$ is positive definite, then $\|\cdot\|$ is a norm.

Proof. *(i)* clear by definition.

(ii) Without loss of generality, assume $\langle x,y\rangle \neq 0$.

Special case: $\langle x,y\rangle \in \mathbb{R}_+$. Assume in addition that $\|x\| \neq 0, \|y\| \neq 0$. Set $\xi := \frac{x}{\|x\|}$ and $\eta := \frac{y}{\|y\|}$, so $\|\xi\| = \|\eta\| = 1$. We have

$$\begin{aligned} &0 \leq \langle \xi-\eta, \xi-\eta\rangle = \|\xi\|^2 + \|\eta\|^2 - \langle\xi,\eta\rangle - \langle\eta,\xi\rangle = \|\xi\|^2 + \|\eta\|^2 - 2\mathrm{Re}\,\langle\xi,\eta\rangle \\ \iff\ &1 = \frac{1}{2}\left(\|\xi\|^2 + \|\eta\|^2\right) \geq \mathrm{Re}\,\langle\xi,\eta\rangle = \frac{\mathrm{Re}\,\langle x,y\rangle}{\|x\|\,\|y\|} \cdot |\langle x,y\rangle| = \mathrm{Re}\,\langle x,y\rangle \leq \|x\|\,\|y\|\,. \end{aligned} \tag{1.33}$$

If $\|x\| = 0$, apply the calculation with $\xi = kx$ and $\eta = \frac{1}{k}y$. Then, $|\langle x,y\rangle| = |\langle\xi,\eta\rangle| \leq \frac{\|y\|^2}{k^2}$ for all $k > 0$ which implies $\langle x,y\rangle = 0$. A similar argument applies if $\|y\| = 0$.

From special case to general case: Assume $\langle x,y\rangle \neq 0$. Define $\alpha := \frac{\langle x,y\rangle}{|\langle x,y\rangle|}$, so $\|\alpha\| = 1$, and define $\tilde{x} := \overline{\alpha}\cdot x$, so $\langle\tilde{x},y\rangle = \overline{\alpha}\cdot\langle x,y\rangle = \overline{\alpha}\alpha\,|\langle x,y\rangle| = |\langle x,y\rangle| > 0$.

By the special case,

$$|\langle x,y\rangle| = |\langle\tilde{x},y\rangle| \leq \|\tilde{x}\| \cdot \|y\| = \|x\| \cdot \|y\|\,. \tag{1.34}$$

(iii) Exercise.

(iv) Exercise.

□

Lemma 1.37: Scalar product from norm

Let $(X, \|\cdot\|)$ be a normed space. Then, there exists a scalar product $\langle\cdot,\cdot\rangle$ on X such that $\|x\| = \sqrt{\langle x,x\rangle}$ iff $\|x+y\|^2 + \|x-y\|^2 = 2\left(\|x^2\| + \|y\|^2\right)$.

Proof. Exercise. The *only if* part is Lemma 1.36.*(iv)*. □

Definition 1.38: Hilbert space

A pre-Hilbert space which is complete in the norm induced by the scalar product is called a *Hilbert space*.

Chapter 2

Function spaces

2.1 Spaces of bounded, continuous and differentiable functions

Definition 2.1: $B(X;Y)$: Space of bounded functions

Let X be a set and $(Y, \|\cdot\|_Y)$ be a normed space. The *space of bounded functions* $B(X;Y)$ is definied by

$$B(X;Y) := \left\{ f : X \to Y \;\middle|\; \sup_{x\in X} \|f(x)\|_Y < \infty \right\}. \tag{2.1}$$

We define

$$\|f\|_{B(X;Y)} := \sup_{x\in X} \|f(x)\|_Y . \tag{2.2}$$

Proposition 2.2. *The Space $(B(X;Y), \|\cdot\|_{B(X;Y)})$ is a normed space. If Y is a Banach space, then, $B(X;Y)$ is a Banach space.*

Proof. Exercise. □

If $Y = \mathbb{R}$ (or $\mathbb{C}$), we write $B(X) := B(X, \mathbb{R})$.

Example. (*i*) If $X = \{1, \dots, n\}$, we have $B(X;\mathbb{R}) = \mathbb{R}^n$ and $B(X;Y) = \underbrace{Y \times \cdots \times Y}_{n \text{ times}}$.

(*ii*) If $X = \mathbb{N}$ and $Y = \mathbb{R}$, we have $B(\mathbb{N};\mathbb{R}) = \ell_\infty$, the space of bounded sequences.

[18.10.2024, Lecture 4]
[23.10.2024, Lecture 5]

Proposition 2.3. *The space $B(X) = B(X;\mathbb{R})$ is separable if and only if X is finite.*

Example. If $X = \mathbb{N}$, then $B(X) = \ell_\infty$, the space of bounded sequences, is not separable.

Proof. If X is finite, wlog $X = \{1, \dots, n\}$. Then,

$$B(X) = \{f : \{1, \dots, n\} \to \mathbb{R}\} = \{(f_1, \dots, f_n) \mid f_i \in \mathbb{R}\} \cong \mathbb{R}^n \tag{2.3}$$

As $\mathbb{R}$ is separable because $\mathbb{Q}$ is dense and countable, and finite products of separable spaces are separable, we have that $B(X)$ is separable.

Assume X is not finite. Idea: Find a set $A \subseteq B(X)$ which is not countable and such that

$$\|a - b\| \geq \delta > 0 \text{ for all } a \neq b \in A. \tag{2.4}$$

If $B(X)$ was separable, there exists a $D \subseteq B(X)$ which is dense and countable.

Claim: (2.4) implies that there exists an injective map $j : A \to D$.

Proof of the claim: Let $a \in A$. By density of D, there exists $a' \in D$ such that $\|a - a'\| < \frac{\delta}{2}$. Similar, take $\|b - b'\| < \frac{\delta}{2}$. If $a' = b'$, by the triangle inequality, we get $\|a - b\| < \delta$, so we get $a = b$.

Take $A = \{f : X \to \{0,1\}\} = \{0,1\}^X$; this set is uncountable by Cantor's second diagonalization argument (see Analysis I). If $f \neq g \in A$, then there exists an A such that $f(x) - g(x) \in \{-1, 1\}$, so we get that $\|f - g\| \geq 1$. □

Definition 2.4: $C(X;Y), C_b(X;Y)$: Space of continuous (bounded) functions

Let $(X, \mathcal{T})$ be a topological space and $(Y, \|\cdot\|)$ a normed space. Then, we define

$$\begin{aligned} C(X;Y) &:= \{f : X \to Y \mid f \text{ is continuous}\} \\ C_b(X;Y) &:= \{f : X \to Y \mid f \text{ is continuous and bounded}\}. \end{aligned} \tag{2.5}$$

In general, $C \neq C_b$, e.g. $X = (0,1), Y = \mathbb{R}, f = \frac{1}{x}$.

Proposition 2.5. *Let $(Y, \|\cdot\|)$ be a Banach space and for $f \in C_b(X;Y)$ define*

$$\|f\|_{C_b} := \sup_{x \in X} \|f(x)\|. \tag{2.6}$$

Then, $(C_b(X;Y), \|\cdot\|_{C_b})$ is a Banach space.

Proof. We have $C_b(X;Y) \subseteq B(X;Y)$. Hence, it suffices to show that if $f_k \in C_b(X;Y)$ and if $\|f_k - f_*\| \to 0$ for $f_* \in B(X;Y)$, then $f_* \in C_b(X;Y)$. The key for that is the triangle inequality:

$$\|f_*(z) - f_*(x)\| \leq \underbrace{\|f_*(z) - f_k(z)\|}_{\leq \|f_k - f_*\|} + \underbrace{\|f_k(z) - f(k)_x\|}_{\substack{\text{use continuity} \\ \text{of } f_k}} + \underbrace{\|f_*(x) - f_k(x)\|}_{\leq \|f_k - f_*\|}. \tag{2.7}$$

Let $\varepsilon > 0, x \in X$. There exists a $k \in \mathbb{N}$ such that $\|f_k - f_*\| < \frac{\varepsilon}{3}$.

By the continuity of f_k there exists an open neighbourhood U of x such that for all $z \in U$ we have $\|f_k(z) - f_k(x)\| < \frac{\varepsilon}{3}$. Hence, if $z \in U$, we get

$$\|f_*(z) - f_*(x)\| < \frac{\varepsilon}{3} + \frac{\varepsilon}{3} + \frac{\varepsilon}{3} = \varepsilon. \tag{2.8}$$

□

Theorem 2.6: $C(X;Y) = C_b(X;Y)$ if X is compact

*If $(X, \mathcal{T})$ is a **compact** topological space and $(Y, \|\cdot\|)$ is a normed space, then we have $C(X;Y) = C_b(X;Y)$.*

Proof. The map $Y \to \mathbb{R}, y \mapsto \|y\|$ is continuous and we have $|\|y'\| - \|y\|| \leq \|y' - y\|$.

If $f \in C(X;Y)$, we also get that $g : x \mapsto \|f(x)\|$ is continuous with $g : X \to \mathbb{R}$, and by Theorem 1.12 we know that $\max_{x \in X} g(x) < \infty$. □

Theorem 2.7: $C(X;Y)$ is separable if domain is compact

*Let (X, d) be a **compact** metric space and $(Y, \|\cdot\|)$ be a separable, normed space. Then, $C(X;Y)$ is separable.*

Example. Let $Y = \mathbb{R}$. If (X, d) is compact, we get that $C(X, \mathbb{R}) = C(X)$ is separable. By proposition 2.3 $B(X)$ is separable iff X is finite. Thus, there are more bounded functions than there are continuous functions.

Proof. Exercise. The key idea is that continuous on a compact set $\Rightarrow$ uniformly continuous, e.g. if $f \in C(X; Y)$,

$$\forall \varepsilon > 0 \exists \delta_{f,\varepsilon} > 0 : \quad d(x, y) < \delta \Rightarrow \|f(x) - f(y)\| < \varepsilon. \tag{2.9}$$

Cover X by finitely many δ-balls $B(x_i, \delta)$ for $= 1, \dots, N$. □

Theorem 2.8: $C(U; Y)$ for $U \subseteq \mathbb{R}^n$

Let $U \subseteq \mathbb{R}^n$ open, $(Y, \|\cdot\|)$ normed space. Let $K_i \subseteq U$ be compact with $K_i \subseteq K_{i+1}$ and $\bigcup_{i=1}^{\infty} K_i = U$, and assume that for every $x \in U$, there exists an $r > 0$ and an i such that $B(x, r) \subset K_i$. For $f \in C(U; Y)$ define

$$[f]_i := \sup_{x \in K_i} \|f(x)\|, \qquad d(f, g) := \sum_{i=1}^{\infty} 2^{-i} \frac{[f - g]_i}{1 + [f - g]_i}. \tag{2.10}$$

Then, $[\cdot]_i$ is a seminorm on $C(U; Y)$ and d is a metric on $C(U; Y)$.
If Y is a Banach space, then $(C(U; Y), d)$ is complete. If Y is separable, then $(C(U; Y), d)$ is separable.

Proof. $[\cdot]_i$ is a seminorm, see the triangle inequality above. A function could be 0 on K_i but not 0 on the entire space.

We get that

$$d_i(f, g) = \frac{[f - g]_i}{1 + [f - g]_i} \tag{2.11}$$

is a semimetric, so also

$$d(f, g) = \sum_{i=1}^{\infty} 2^{-i} d_i(f, g) \tag{2.12}$$

is a semimetric. So we need to show that $d(f, g) = 0 \Rightarrow f = g$: If $d(f, g) = 0 \Rightarrow [f - g]_i = 0$, so $f - g = 0$ on K_i. As $U = \bigcup_{i=1}^{\infty} K_i$, we get $f - g = 0$ on U.

Completeness: Let $j \mapsto f_j$ be a Cauchy sequence for d, so f_j is Cauchy for each $[\cdot]_i$, so also $f_j|_{K_i}$ is a Cauchy sequence in $C(K_i; Y) = C_b(K_i; Y)$ as K_i is compact. Hence, there exists a $f_{*,i} : K_i \to Y$ such that $f_j|_{K_i} \to f_{*,i}$. As $K_i \subseteq K_{i+1}$, we get that $f_{*,i+1}|_{K_i} = f_{*,i}$. With that, we can define $f_* : U \to Y$ such that $f_{*,i} = f_*|_{K_i}$. With that, we get that $\|f_j|_{K_i} - f_*|_{K_i}\| \to 0$ and with that, we get that $d(f_j, f_*) \to 0$. □

Remark. (*i*) $\sum_{i=1}^{\infty} 2^{-i} \frac{d_i}{1+d_i}$ is used in various examples e.g. $\mathbb{R}^{\mathbb{N}}$.

(*ii*) There exists no norm on $C(U; Y)$ which induces the same topology as d (similar to the homework for $\mathbb{R}^{\mathbb{N}}$).

(*iii*) For each $U \subseteq \mathbb{R}^n$ open, there exists a sequence K_i of compact sets, e.g.

$$K_i := \left\{ x \in U \;\middle|\; \operatorname{dist}(x, \mathbb{R}^n \setminus U) \geq 2^{-i}, |x| \leq 2^i \right\}. \tag{2.13}$$

(*iv*) Different choices of the sequence K_i lead to the same topology on $C(U; Y)$ (or to equivalent metrics)

Proposition 2.9. *Let $U \subseteq \mathbb{R}^n$ open, $C(U; Y)$ as in Theorem 2.8. Let $f : \mathbb{N} \to C(U; Y)$ and $g \in C(U; Y)$. Then, the following statements are equivalent:*

(*i*) $\lim_{j\to\infty} d(f_j, g) = 0$,

(*ii*) $\lim_{j\to\infty} d_i(f_j, g) = 0$ *for all* i *(where* $d_i(f,g) = [f-g]_i$*),*

(*iii*) ***for all*** *compact set* $K \subseteq U$ *we have*

$$\lim_{j\to\infty} \sup_{x\in K} \|f_j(x) - g(x)\| = 0, \tag{2.14}$$

we call this locally uniform convergence.

Proof. „$(iii) \Rightarrow (ii)$“: obvious.

„$(ii) \Rightarrow (i)$“: We have $\lim_{j\to\infty} d_i(f_j, g) = 0$, so also

$$\lim_{j\to\infty} \frac{d_i(f_j,g)}{1+d_i(f_j,g)} \to 0 \quad \Rightarrow \quad \sum_{i=1}^{N} 2^{-i} \frac{d_i(f_j,g)}{1+d_i(f_j,g)} \to 0 \tag{2.15}$$

and with that also

$$\limsup_{j\to\infty} d(f_j,g) \le \sum_{i=1}^{N} \frac{d_i(f_j,g)}{1+d_i(f_j,g)} + \sum_{i=N+1}^{\infty} 2^{-i} \le 2^{-N} \tag{2.16}$$

and as N was arbitary, this goes to 0.

„$(i) \Rightarrow (iii)$“: For each $x \in K$ there exist $i_x, r_x > 0$ such that $B(x, r_x) \subseteq K_{i_x}$. Then, we have $\bigcup_{x\in K} B(x, r_x) \supseteq K$. As K is compact, there exist $x_1, \dots, x_N, r_1, \dots, r_n$ such that

$$K \subseteq \bigcup_{i=1}^{N} B(x_k, r_k) \qquad \text{where} \quad B(x_k, r_k) \subseteq K_{i_k}. \tag{2.17}$$

□

[23.10.2024, Lecture 5]
[25.10.2024, Lecture 6]

Definition 2.10: $C^k, C^\infty, [f]_{C^l}, \|f\|_{C^k}, d(f,g)$

Let $U \subset \mathbb{R}^n$ be open and bounded. We define

$$C^k(\overline{U};\mathbb{R}^m) := \left\{ f \in C^k(U;\mathbb{R}^m) \;\middle|\; D^l f \text{ has a continuous extension to } \overline{U} \;\forall l \le k \right\} \tag{2.18}$$

and, for $0 \le l \le k$,

$$[f]_{C^l(U)} := \sup_{x\in\overline{U}} \left| D^l f(x) \right|, \quad \|f\|_{C^k} := \sum_{l=0}^{k} [f]_{C^l}. \tag{2.19}$$

Moreover, we set $C^\infty(\overline{U};\mathbb{R}^m) := \bigcup C^k(\overline{U};\mathbb{R}^m)$ and

$$d(f,g) := \sum_{l=0}^{\infty} 2^{-l} \frac{\|f-g\|_{C^l}}{1+\|f-g\|_{C^l}}. \tag{2.20}$$

Theorem 2.11: $\|\cdot\|_{C^k}$ **is norm;** C^∞ **traits**

$[\cdot]_{C^l}$ is a seminorm on $C^k(\overline{U};\mathbb{R}^m)$ for $0 \le l \le k$, and $\|\cdot\|_{C^k}$ is a norm.
Moreover, if $C^k(\overline{U};\mathbb{R}^m)$ is a ***separable*** *Banach space, then, $C^\infty(\overline{U};\mathbb{R}^m)$ is a complete, separable metric space.*

Proof. Exercise. Hints in the notes. □

Definition 2.12: Functions with compact support

(*i*) Let $(X, \mathcal{T})$ be a topological space, $(Y, \|\cdot\|)$ a normed space and $f : X \to Y$. Then, we define the *support* of f

$$\operatorname{supp}(f) := \overline{\{x \in X \mid f(x) \neq 0\}}. \tag{2.21}$$

(*ii*) The space of continuous functions with compact support is defined as

$$C_c(X;Y) := \{f \in C(X;Y) \mid \operatorname{supp}(f) \text{ is compact}\}. \tag{2.22}$$

(*iii*) Let $U \subseteq \mathbb{R}^n$ oben, $Y = \mathbb{R}^m$. Then, we set

$$C_c^k(U;\mathbb{R}^m) := C^k(U;\mathbb{R}^m) \cap C_c(U;\mathbb{R}^m) \subseteq C^k(U;\mathbb{R}^m) \tag{2.23}$$

and

$$\mathcal{D}(U;\mathbb{R}^m) := C_c^\infty(U;\mathbb{R}^m) = C^\infty(U;\mathbb{R}^m) \cap C_c(U;\mathbb{R}^m). \tag{2.24}$$

Notation. Some people write $C_0(\dots)$ instead of $C_c(\dots)$. People wbo use $C_c(\dots)$ sometimes define $C_0(U;\mathbb{R}^m) := \overline{C_c(U;\mathbb{R}^m)}$ in $C(\overline{U};\mathbb{R}^m)$. Those are not the same as for $U = (-1,1)$ and $f(x) = 1 - |x|$, we have $f \in C_0(U)$ but $f \notin C_c(U)$.

Interesting subspaces of the space of continuous functions arise when we consider functions with a given modulus of continuity $\rho : [0,\infty) \to [0,\infty)$, i.e., functions for which $\|f(x) - f(y)\| \leq C_f \rho(d(x,y))$ where $\lim_{t \downarrow 0} \rho(t) = 0$. The most important examples are Hölder continuous functions, which correspond to $\rho(t) = t^\alpha$. For simplicity we focus on functions on $A \subseteq \mathbb{R}^n$.

Definition 2.13: Hölder continuity

Let $\alpha \in (0,1]$, $A \subseteq \mathbb{R}^n$ and $f : A \to \mathbb{R}^n$. Then, we say that f is *Hölder continuous* if

$$[f]_{\alpha,A} := \sup_{\substack{x,y \in A \\ x \neq y}} \frac{|f(x) - f(y)|}{|x - y|^\alpha}. \tag{2.25}$$

Then, we define

$$C^{0,\alpha} := \{f \in C_b(A;\mathbb{R}^m) \mid [f]_{\alpha,A} < \infty\} \tag{2.26}$$

and

$$\|f\|_{\alpha,A} := \sup_{x \in A} |f(x)| + [f]_{\alpha,A}. \tag{2.27}$$

If $U \subseteq \mathbb{R}^n$ is open and bounded, we define

$$C^{k,\alpha} := \left\{ f \in C^k(\overline{U};\mathbb{R}^m) \;\middle|\; \left[D^k f\right]_{0,\alpha,U} < \infty \right\} \tag{2.28}$$

and

$$\|f\|_{k,\alpha,U} := \|f\|_{C^k(U)} + \left[D^k f\right]_{0,\alpha,U}. \tag{2.29}$$

Proposition 2.14. *$[\cdot]_{\alpha,A}$ is a seminorm on $C^{0,\alpha}(A;\mathbb{R}^m)$ and the spaces $C^{0,\alpha}(A;\mathbb{R}^m)$ and $C^{k,\alpha}(A;\mathbb{R}^m)$ are Banach spaces with norms $\|\cdot\|_{\alpha,U}$ and $\|\cdot\|_{k,\alpha,U}$.*

[25.10.2024, Lecture 6]
[30.10.2024, Lecture 7]

Appendix

List of definitions

1.1 Topological space, Topology . . . 3
1.3 Interior, Closure, Boundary . . . 4
1.4 Dense subsets, separable topological space . . . 4
1.5 Open neighbourhood . . . 4
1.6 Continuous maps, Homeomorphisms . . . 4
1.7 Convergence . . . 4
1.8 Hausdorff space . . . 5
1.10 Compactness . . . 5
1.13 Connectedness . . . 5
1.15 Metric space, metric . . . 6
1.22 Equivalent metrics, stronger metrics . . . 8
1.24 Separable metric space . . . 9
1.25 Cauchy sequence, completeness . . . 9
1.27 Isometric immersion, isometry . . . 9
1.30 Norm . . . 11
1.31 Banach space . . . 11
1.32 Stronger norms, equivalent norms . . . 11
1.34 Sesquilinear form . . . 12
1.35 Scalar product . . . 12
1.38 Hilbert space . . . 13
2.1 $B(X;Y)$: Space of bounded functions . . . 14
2.4 $C(X;Y), C_b(X;Y)$: Space of continuous (bounded) functions . . . 15
2.10 $C^k, C^\infty, [f]_{C^l}, \|f\|_{C^k}, d(f,g)$ . . . 17
2.12 Functions with compact support . . . 18
2.13 Hölder continuity . . . 18

List of statements

1.11 Continuous maps compact to compact 5
1.12 Continuous maps with compact domain attain max and min 5
1.14 Intermediate value theorem 5
1.16 Composition of concave map and metric is metric 6
1.18 Hausdorff distance is metric on closed and bounded sets 7
1.28 Completion 10
1.29 Hausdorff metric space is complete 10
1.36 Scalar product traits 13
1.37 Scalar product from norm 13
2.6 $C(X;Y) = C_b(X;Y)$ if X is compact 15
2.7 $C(X;Y)$ is separable if domain is compact 15
2.8 $C(U;Y)$ for $U \subseteq \mathbb{R}^n$ 16
2.11 $\|\cdot\|_{C^k}$ is norm; C^∞ traits 17

Index

$C(X;Y)$, 15
$C_b(X;Y)$, 15

Banach space, 11
bounded functions space, 14

Cauchy sequence, 9
Cauchy-Schwarz inequality, 13
closed set, 3
compact, 5
complete, 9
completion, 10
connected, 5
continuity
    Hölder -, 18
continuous, 4
convergence, 4
    locally uniform -, 17
countable, 3

definiteness, 11

Gromov-Hausdorff metric, 10

Hausdorff, 5
    - space, 5
Hausdorff distance, 7
Hilbert space, 13
    Pre-, 12
homeomorphism, 4
homogeneity, 11
Hölder
    - continuous, 18

isometric immersion, 9
isometry, 9

locally uniform convergence, 17

metric, 6
    equivalent -, 8
    Gromov-Hausdorff -, 10
    pseudo-, 6
    pullback -, 6
    semi-, 6
    stronger -, 8
metric space, 6
    complete -, 9
    separable -, 9

norm, 11
    equivalent -, 11
    semi-, 11
    stronger -, 11
normed space, 11

open neighbourhood, 4
open set, 3

parallelogram identity, 13
pseudometric, 6
pullback metric, 6

scalar product, 12
semimetric, 6
seminorm, 11
separable, 4, 9
sequence
    Cauchy -, 9
sequential closure, 8
sesquilinear form, 12
    positive definite -, 12
    positive semidefinite -, 12
    symmetric -, 12
set
    boundary, 4
    closed -, 3
      sequentially -, 8
    closure, 4
    compact -, 5
    countable, 3
    dense, 4
    Hausdorff distance between -s, 7
    interior, 4
    open -, 3
    open neighbourhood, 4
    sequentially closed -, 8
space
    - of continuous functions, 15
      - (bounded), 15
    Banach -, 11
    bounded function -, 14
    completion (of -), 10
    connected -, 5
    Hausdorff -, 5
    Hilbert -, 13
      Pre-, 12

metric -, 6
normed -, 11
topological -, 3
support, 18
topological space, 3
connected -, 5
separable -, 4
Topology, 3
topology
relative -, 3
triangle inequality, 6, 11, 13
reversed -, 11