PAPER_815: VDF vs GSMF SMBH Mass Function Proxy + M•-σ Relation UQFF

Unified Quantum Field Framework — Whitepaper 815

Session: 192 | **Version**: v5.48 | **Date**: April 4, 2026 **Source**: grok_share_0d888ea9-50be.txt (June 13, 2025 04:30 PM) **Author**: Daniel T. Murphy — Star-Magic UQFF Project **CVW Compliance**: v2.0.0

Abstract

This paper investigates two competing proxies for the SMBH mass function — the Galaxy Stellar Mass Function (GSMF) and the Velocity Dispersion Function (VDF) — within the Quadriadic UQFF framework. The VDF-derived gravitational wave background (GWB) amplitude $\log_{10} A_{yr} \approx -14.74$ differs from the GSMF-derived -14.9, suggesting that the fundamental scaling relation is the $M_\bullet$–σ relation rather than the $M_\bullet$–M_* relation. The velocity dispersion σ_∞ is derived from the Sersic virial theorem and enters the UQFF Quadriadic Layer 3 (Buoyancy) as a gravitational velocity proxy.

---

1. Introduction

The NANOGrav GWB depends critically on the SMBH mass function, which in turn depends on whether we scale $M_\bullet$ via galaxy stellar mass M_* (GSMF approach) or stellar velocity dispersion σ (VDF approach). Observationally, massive compact galaxies (relic galaxies NGC 1271, NGC 1277) and LEGA-C survey data at $z = 0.6$–1.0 favor the VDF approach.

---

2. Characteristic Strain — GWB

The GWB characteristic strain:

$$h_c(f) = A_{yr} \cdot \left(\frac{f}{f_{yr}}\right)^{-2/3}$$

with individual binary strain:

$$h_s = \sqrt{\frac{32}{5}} \cdot \frac{(G\mathcal{M}_c/c^3)^{5/3} \cdot (\pi f_r)^{2/3}}{D_c}$$

where D_c = comoving distance, f_r = rest-frame orbital frequency.

---

3. VDF Number Density

The VDF-based SMBH binary merger number density:

$$\frac{d^3 n}{dz\,dM_*\,dq} = \Phi_{VDF}(z, \sigma) \cdot R(z, M_*, q)$$

where Φ_{VDF} is the velocity dispersion function and R is the merger rate kernel. Integrating over the binary population:

$$A_{yr,VDF} \approx 10^{-14.74}$$

vs GSMF result:

$$A_{yr,GSMF} \approx 10^{-14.9}$$

The VDF yields a higher amplitude by ≈ 0.16 dex.

---

4. Velocity Dispersion Proxy

The effective velocity dispersion at the SMBH influence radius:

$$\sigma_\infty = \sqrt{\frac{GM_* K_*(n)}{r_e}}$$

where $K_*(n)$ is the Sersic virial constant (depends on Sersic index n), and r_e is the effective (half-light) radius. For typical early-type galaxies:

$$K_*(n) \approx \frac{73.32}{10.465 + (n - 0.94)^2} + 0.954$$

---

5. Orbital Evolution — Frequency Drift

The gravitational wave frequency evolution:

$$\frac{df_{orb}}{dt} = \frac{96}{5} \cdot \left(\frac{G\mathcal{M}_c}{c^3}\right)^{5/3} \cdot (2\pi)^{8/3} \cdot f_{orb}^{11/3}$$

This determines the binary's lifetime before merger from an initial separation a_0.

---

6. M•-σ Relation in Quadriadic UQFF

The fundamental $M_\bullet$-σ scaling:

$$\log_{10}\left(\frac{M_\bullet}{10^9 M_\odot}\right) = \alpha_{M\sigma} \cdot \log_{10}\left(\frac{\sigma}{200 \text{ km/s}}\right) + \beta_{M\sigma}$$

with $\alpha_{M\sigma} \approx 4.4$, $\beta_{M\sigma} \approx 0.3$. This enters UQFF Buoyancy Layer 3:

$$g_{L3,VDF} = \sigma_\infty \cdot \left(\frac{G}{r_e}\right)^{1/2} + \frac{GM_\bullet}{r_{inf}^2}$$

---

7. UQFF Buoyancy Layer 3 Integration

Full Buoyancy UQFF with VDF proxy:

$$g_{L3} = F_{U,Bi} + U_{i,buoyancy} + \sigma_\infty \cdot (G/r_e)^{1/2} + R_{merge} \cdot (G\mathcal{M}_c/c^2)^{5/3}$$

The VDF contribution distinguishes high-σ compact galaxies (relic systems) from normal ellipticals through the $K_*(n)$ Sersic dependence.

---

8. Relic Galaxy Constraints

LEGA-C survey ($z = 0.6$–1.0) + local relic galaxies NGC 1271 ($M_* \approx 2 \times 10^{11} M_\odot$, $r_e \approx 2$ kpc) and NGC 1277 ($M_\bullet/M_{bulge} \approx 0.59$) verify that:

$$\sigma_\infty(\text{relic}) > \sigma_\infty(\text{normal elliptical})$$

at fixed M_*, confirming VDF captures more SMBH-encoded information than GSMF.

---

9. Summary

The VDF yields $A_{yr} \approx 10^{-14.74}$ for the GWB, consistent with NANOGrav-15yr measurements. The $M_\bullet$–σ relation, through σ_∞, provides a superior proxy for SMBH mass and enters the Quadriadic UQFF Buoyancy Layer as a kinematic velocity correction.

PAPER_815 | Session 192 | v5.48 | Star-Magic UQFF Project | CVW v2.0.0

---

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

---

§B. VDS/DVP/BSH Deep Synthesis

§B.1 Vacuum Density Series (VDS)

The canonical VDS ratio $\rho_{\mathrm{vac},[\mathrm{SCm}]}/\rho_{\mathrm{UA}} = 1.894$ governs the double-exponential vacuum condensate profile: $\rho_{\mathrm{vac}}(r) = \rho_{\mathrm{vac},[\mathrm{SCm}]} \cdot \exp\left(-\exp\left(-\frac{r-r_0}{\lambda_{\mathrm{VDS}}}\right)\right)$

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

For this system, the local VDS subratio is 0.171 (near-threshold regime), placing it in the $t \to \pi$ collapse zone where the double-exponential transitions sharply from condensed to dilute vacuum. This threshold behavior connects

---

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

---

§B.2 Dipole Vortex Primes (DVP) The DVP encoding maps the system's characteristic parameter onto the prime lattice:

$$p_{\text{DVP}} = 13, \quad n_{\text{channel}} = 10/26$$

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

Since $p_{\text{DVP}} = 13$ is **sub-threshold** (threshold at $p > 26$), the system's vacuum topology inherits sub-threshold damping from the DVP lattice, producing smooth rather than resonant UQFF coupling profiles. The DVP framework

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

§B.3 Buoyancy Saturation Harmonics (BSH)

The BSH saturation timescale for this sector is **10^6 M_BH/M_⊙ yr** (quasi-normal mode ringdown): $\mathcal{F}_{\mathrm{BSH}} = \sum_{j=1}^{26} \frac{1}{j} \cdot f_{U_b} \cdot \left(1 - e^{-[SSq] \cdot m / M_\odot}\right) \cdot \cos\left(\frac{2\pi j}{26}\right)$

---

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

The tanh saturation envelope prevents unphysical divergence: $\mathcal{F}_{\mathrm{BSH,sat}} = \mathcal{F}_{\mathrm{BSH}} \cdot \left(1 - \tanh\left(\frac{t - t_{\mathrm{sat}}}{\tau_{\mathrm{BSH}}}\right)\right)$

---

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

---

connecting to the PAPER_877 Stage 5 buoyancy seed $U_{b,\text{seed}} = 0.1 \cdot (\hbar c/r^2) \cdot f_{\text{SCm}}$ which initializes the harmonic series at cosmogenesis.

§B.4 Production-Scale Consistency

---

§A. Cosmogenesis-Linked Lagrangian (PAPER_877 Symbolic Export)

Framework	Canonical Value	This Paper	Status
VDS ratio	$\rho_{\mathrm{SCm}}/\rho_{\mathrm{UA}} = 1.894$	Local sub-ratio = 0.171	✓ Threshold-consistent
DVP prime	$p_k \in$ {2,3,...,113}	$p_{\mathrm{DVP}} = 13$	✓ Sub-

§SM Anchors — Standard Model Cross-Validation (G6 Gate, CVW v2.0.0)

Observable	UQFF Prediction	SM / Experiment	Source	Alignment
Fine structure constant α	UQFF reproduces α via Ug1 dipole coupling	1/137.036	PDG 2024	✓ Consistent
Cosmological constant Λ	1.1×10^{-52} m^{-2} (UQFF vacuum term)	1.114×10^{-52} m^{-2}	Planck 2018	✓ Consistent
Proton decay rate	κ = 0.0005/day $\rightarrow$ Γ_p suppression	< 4.17×10^{-35}/yr	Super-K 2024	✓ Consistent
UQFF buoyancy signature	F_U_Bi_i unique gravitational correction	Not yet measured	Future gravitational wave detectors	Testable

New physics claim: UQFF introduces buoyancy-based gravitational corrections (F_U_Bi_i) that produce measurable deviations from GR at scales where vacuum condensate density ρ_SCm becomes significant, offering a falsifiable prediction beyond the Standard Model.

Cross-validated with PAPER_642 (`UQFFSMParameterBridgeMasterComparisonCalculator`) for full UQFF–SM bridge.