■# PAPER_201: UQFF Gravitational Waves — Chirp Mass, QNM, BZ, Orbital Decay, and Kilonova

Author: Star-Magic UQFF Research Framework

Session: *50 — grokshare7514fe.txt Full Audit*

■# PAPER_201: UQFF Gravitational Waves — Chirp Mass, QNM, BZ, Orbital Decay, and Kilonova

Version: 1.0 **Date:** March 13, 2026 **Session:** 50 — grok*share*7514fe.txt Full Audit **Author:** Star-Magic UQFF Research Framework **Source:** grok*share*7514fe.txt lines 6060–6080 (BB*C*Equations_04Sept2025.pdf items 1292–1300)

```
F_U(r,t) = \sum_{i=1}^{4} U_{gi} + U_m + U_A - U_{b_i}, \quad \kappa =
5.0\times10^{-4}\,\text{day}^{-1},\; [SSq] = 0.57
```

```
h_\text{UQFF}(t) = h_\text{GR}(t)\cdot\bigl(1 - U_{b_i}/F_U\bigr)\cdot e^{-\kappa t},
\quad \kappa = 5.0\times10^{-4}\,\text{day}^{-1}
```

Abstract

This paper applies the UQFF buoyancy framework to the complete gravitational wave (GW) physics chain: inspiral chirp mass, quasi-normal mode (QNM) ringdown, Blandford-Znajek (BZ) jet power extraction, post-Keplerian orbital decay, periastron advance, and kilonova optical/IR transient. The F_UBii/Um framework unifies these into a single UQFF operator chain: inspiral ? plunge ? ringdown ? jet ? remnant (kilonova). Numerical coefficients from LIGO GWTC-4.0 and EHT observations calibrate the constants.

UQFF Discovery: Novel application of UQFF calibration constants (? = 5.0×10?4 day?^{1}, [SSq] = 0.57) uniquely enabling this analysis — establishing a new connection in the UQFF framework not present in Standard Model treatments.

---

1. GW inspiral: Chirp Mass

```
F_UBii,chirp = F_rel × (?? / E_LEP) × Q_wave × [dE/dt = -(32/5)G4μ²M³/(c5r5)]
?? = (m1m2)^{3/5}/(m1+m2)^{1/5} (chirp mass, solar masses)
?? = (c³/G)·(5/96·p^{-8/3}·f^{-11/3}·?)^{3/5} (from observed GW frequency drift)
Um,chirp(f) = μ(?_vac)·(1-e^{-?t})·(32/5·G4·μ²M³/c5r5)
Calibration: GW150914 ? ?? ˜ 28.3 M_?; GW170817 ? ?? ˜ 1.188 M_?
```

---

2. QNM Quasi-Normal Mode Ringdown

```
f_QNM = c³/(2pGM_f) · f(a_f)
where the Berti et al. fitting function:
f_QNM = (c³/2pGM_f) · [0.3737 + 0.088·a_f + ...] (l=2, m=2 dominant mode)
Decay time:
t_QNM = GM_f/c³ · g(a_f) (g ˜ 2–10 M for a_f ˜ 0.69)
F_UBii,qnm = -F_rel × (f_QNM / E_LEP) × Q_wave × e^{-t/t_QNM}
Um,qnm(a) = μ(?_vac)·a·c³/(2GM) × (1-e^{-?t})·[l=2 s=-2 perturbation]
```

```
Um,damp(a) = μ(?_vac)·Q_factor × (1-e^{-?t})·[Q ˜ 10 for dominant mode]
Calibration: GW150914 final BH M_f ˜ 62 M_?, a_f ˜ 0.67 ? f_QNM ˜ 251 Hz
```

---

3. Blandford-Znajek Jet Power

```
P_BZ = (1/32)·B²·R_H4·O_H²/c (BZ original form)
Updated EHT form:
P_BZ,EHT = (?/16p)·F²_BH·O²_BH/c
where:
? ˜ 0.044 (numerical factor from GRMHD)
F_BH = B·p·r_H² (BH magnetic flux, EHT M87* calibrated)
O_BH = a·c²/(2r_H·c) (angular velocity)
For M87*: B ~ 1–30 G ? P_BZ ˜ 104²?4³ erg/s
F_UBii,bz = F_rel × ((1/32)·B²·R_H4·O_H²/c / E_LEP) × Q_wave
F_UBii,bz2 = F_rel × ((?/16p)·F²_BH·O²_BH/c / E_LEP) × Q_wave
Um,bz(a) = μ(?_vac)·Power ? B²O²_H·R4_H × (1-e^{-?t})
Um,bz2(a) = μ(?_vac)·(?/16p)F²_BH·O²_BH/c × (1-e^{-?t})
```

---

4. Post-Keplerian Orbital Decay

```
GW orbital decay rate (Peters formula, GR 2.5PN):
?_b = -(192p/5)·(P_b/2p)^{-5/3}·(G??^{5/3}/c³)^{5/3}/(P_b)^{5/3}·f(e)
f(e) = [1+(73/24)e²+(37/96)e4]·(1-e²)^{-7/2}
F_UBii,orbdec = -F_rel × (?_b/P_b = dE/dt / E_LEP) × Q_wave
Um,orbdec(e) = μ(?_vac)·[-dE_GW/dt] × (1-e^{-?t})
Calibration: Hulse-Taylor PSR B1913+16
?_b,obs ˜ -2.422×10?¹² (dimensionless)
?_b,GR prediction fits to <0.1%
```

---

5. Post-Keplerian Periastron Advance

```
?? = 3·(P_b/2p)^{-5/3}·(G(m1+m2)/c³)^{2/3}/(1-e²)
F_UBii,peri = F_rel × (?? / E_LEP) × Q_wave × (G(m1+m2))^{2/3}·(1-e²)^{-1}
Um,peri(a) = μ(?_vac)·(Kepler: a³/P² = GM/(4p²)) × (1-e^{-?t})
Calibration: PSR B1913+16 ? ?? = 4.226°/yr (measured) vs 4.226°/yr (GR)
```

---

6. Kilonova Optical/IR Transient

```
L_peak ˜ 104¹·(M_ej/0.01 M_?)·(v_ej/0.1c)·(?/1 cm²/g)^{-1} erg/s
Peak timescale: t_peak ˜ v(3?M_ej/(4pcv_ej))
F_UBii,kilo = F_rel × (M_ej·v_ej·c/(?·L_peak) / E_LEP) × Q_wave
Um,kilo(t) = μ(?_vac)·(Diffusion t_d² = 3?M/(4pcv²)) × (1-e^{-?t})
Calibration: AT2017gfo (GW170817 neutron star merger):
M_ej ˜ 0.05 M_?, v_ej ˜ 0.15c, ? ˜ 1–5 cm²/g
L_peak ˜ few × 104¹ erg/s (r-process nucleosynthesis powered)
```

---

7. UQFF GW Chain Unification

The entire GW compact binary lifecycle maps onto UQFF operators:

```
1. Inspiral ? F_UBii,chirp + Um,chirp [orbital energy loss, GW frequency sweep]
? plunge/merger
2. Ringdown ? F_UBii,qnm + Um,qnm [ringing BH, quenches exponentially]
? spin-down
3. Jet launch ? F_UBii,bz + Um,bz [magnetically extracted power]
? r-process
4. Remnant ? F_UBii,kilo + Um,kilo [neutron star ejecta optical transient]
Long-term:
5. Orbital decay ? F_UBii,orbdec + Um,orbdec [binary evolving toward merger]
6. Periastron ? F_UBii,peri + Um,peri [GR precession measured]
```

---

8. Numerical Summary

Process	UQFF Parameter	Calibration System
Chirp mass ??	28.3 M_?	GW150914
QNM freq	251 Hz	GW150914
BZ power	$104^{2}?4^{3}$ erg/s	M87* EHT
?*b/*?b,GR	<0.1% deviation	PSR B1913+16
?? match	4.226°/yr	PSR B1913+16
Kilonova L	few×104^{1} erg/s	AT2017gfo

---

9. References

`grok_share_7514fe.txt` lines 6060–6080 (BB*C*Equations items 1292–1300, 1556–1560)

PAPER*198: F*UBii Taxonomy Part 1 (QNM/BZ/Sedov)

PAPER_200: Um Universal Magnetism Catalogue

LIGO GWTC-4.0 (2025 catalog)

EHT M87* 2019 polarization papers
