

PAPER_018: Aether Noise Spectrum Characterization for LISA

Authors: Daniel Murphy & UQFF Research Collective

Date: 2026-03-07

Domain: 1.2 — Gravitational Waves — LISA Future Detector

Primary Validation File: `validate_lisa_extended.py`

Calibration constants: $\kappa = 0.0005/\text{day}$, $[\text{SSq}] = 0.57$

Abstract

We characterize the spectral signature of UQFF aether fields in the LISA frequency band (0.1–100 mHz). UQFF predicts that U_m vacuum field oscillations imprint narrow spectral lines at harmonics of $f_U \approx 1$ mHz onto the stochastic gravitational wave background (SGWB), with an integrated aether power fraction of 222.93% relative to the GR SGWB. The peak aether feature at $f_{\text{peak}} = 0.99$ mHz yields an integrated detection SNR of 12,695,834—trivially detectable by LISA over a 4-year mission. A broad TRZ suppression dip ($\sim 10\%$) appears near 5 mHz. These spectral features have no astrophysical analogue and constitute a smoking-gun test of UQFF vacuum structure.

UQFF Discovery: Novel application of UQFF calibration constants ($\kappa = 5.0 \times 10^{-4} \text{ day}^{-1}$, $[\text{SSq}] = 0.57$) uniquely enabling this analysis establishing a new connection in the UQFF framework not present in Standard Model treatments.

1. Physical Motivation

UQFF predicts that the vacuum supports actively coupled aether fields characterized by: - **U_m** : Magnetic energy parameter (= 1.0 in calibrated UQFF) - **β_m** : Modulation parameter (= 0.01) - **f_{TRZ}** : Trans-zero frequency factor (= 0.1)

These fields interact with propagating GWs to create: 1. Spectral **power excess** at fundamental and harmonic frequencies 2. **Sideband pairs** from β_m modulation 3. A broad **suppression dip** from TRZ near 5 mHz

2. Aether Noise Spectrum Model

The total stochastic power spectral density (PSD) in the LISA band:

$$S_{\text{UQFF}}(f) = S_{\text{GR}}(f) \times [1 + P_{\text{aether}}(f)] \times F_{\text{TRZ}}(f)$$

where the GR background follows $S_{\text{GR}}(f) \propto \Omega_{\text{GW}}(f) \propto f^{2/3}$ (inspiral-dominated), and:

$$S_{\text{UQFF}}(f) = S_{\text{GR}}(f) [1 + P_{\text{aether}}(f)] F_{\text{TRZ}}(f)$$

$$P_{\text{aether}}(f) = U_m \sum_n e^{-n/2} \delta(f - n f_U) W(\Delta f)$$

$$F_{\text{TRZ}}(f) = 1 - 0.1 \exp\left[-\frac{(f - f_{\text{TRZ}})^2}{2\sigma_{\text{TRZ}}^2}\right]$$

Key numerical results: $U_m = 1.0e-4$, $f_U(\text{harmonic } 1) = 9.9e-1 \text{ mHz} = 9.9e-4 \text{ Hz}$, $\Omega_{\text{GW}} \sim 1.0e-9$, F_{TRZ} peak suppression = $1.0e-1$ (10%)

$$P_{\text{aether}}(f) = U_m \times \sum_n \exp(-n/2) \times \delta(f - n f_U) \times W(\Delta f)$$

with spectral line width $W(\Delta f) \approx 10\%$ fractional bandwidth, and:

$$F_{\text{TRZ}}(f) = 1 - 0.1 \times \exp[-(f - f_{\text{TRZ,peak}})^2 / (2\sigma_{\text{TRZ}}^2)]$$

3. Spectral Components

Component	Frequency	Amplitude	Origin
GR SGWB	0.1-100 mHz	$\Omega_{\text{GW}} \sim 10^{-9}$	Inspiral population
U_m harmonic $n=1$	0.99 mHz	$U_m \times 1.0$	Fundamental aether line
U_m harmonic $n=2$	$\sim 2 \text{ mHz}$	$U_m \times e^{-1}$	1st harmonic
U_m harmonic $n=3$	$\sim 3 \text{ mHz}$	$U_m \times e^{-1.5}$	2nd harmonic
U_m harmonics $n=4,5$	$\sim 4, 5 \text{ mHz}$	$U_m \times e^{-n/2}$	Higher harmonics
β_m sidebands	$f_n \pm 0.01 \text{ mHz}$	$\beta_m \times \text{amplitude}$	Modulation pairs
TRZ suppression	$\sim 5 \text{ mHz}$	-10%	Trans-zero dip

4. Quantitative Results

Metric	Value
Frequency range	0.10 - 100 mHz
Spectral bins	200
Spectral resolution Δf	$8 \times 10^{-6} \text{ mHz}$
Observation time	4 years
Aether power fraction $P_{\text{aether}}/P_{\text{GR}}$	222.93%
Peak aether frequency f_{peak}	0.99 mHz
Integrated detection SNR	12,695,834
Detectable (SNR > 5)	□ True

The 222.93% aether power fraction means the UQFF aether signal is more than **twice** the GR SGWB in the LISA band—an unmistakably strong signature.

5. Comparison With Known Noise Sources

Source	Spectrum Shape	Prediction	UQFF Distinguishable?
GR SGWB (inspiral)	$\propto f^{(2/3)}$	$\Omega_{\text{GW}} \sim 10^{-9}$	Baseline

Source	Spectrum Shape	Prediction	UQFF Distinguishable?
Galactic WD foreground	Peaked at ~ 3 mHz	Reduced by UQFF factor	Yes — WD band modified
LISA instrument noise	$\propto f^4$ (low f)	Not modified	Yes — different spectral shape
Cosmological SGWB	Flat (inflation)	Not modified	Yes — UQFF adds lines
UQFF aether lines	Harmonic comb	222.93%	Unique signature

6. Validation

Run command (Windows-safe):

```
set PYTHONUTF8=1
python validate_lisa_extended.py
```

Test status from `compute_aether_noise_spectrum()`:

Check	Result
P_aether/P_GR > 100%	☐ PASS (222.93%)
f_peak identified	☐ PASS (0.99 mHz)
Integrated SNR > 5	☐ PASS (12,695,834)
Detectable = True	☐ PASS

TEST STATUS: PASSED: `compute_aether_noise_spectrum`

7. Observational Strategy for LISA

1. **Targeted narrow-band search at 1, 2, 3 mHz:** Comb filter for U_m harmonic signature with 10% bandwidth windows
2. **Cross-correlation test:** Compare LISA data-stream with predicted sideband pattern at $f_n \pm f_{\text{mod}}$ ($f_{\text{mod}} = 0.01$ mHz)
3. **TRZ notch identification:** Low-pass filter around 5 mHz to identify broad suppression against WD foreground
4. **Time-series modulation:** Look for $\sim 10\%$ amplitude oscillations at ~ 1 -hour period in GW background estimation

8. Conclusion

UQFF aether fields create a distinctive spectral fingerprint in the LISA band: a harmonic comb at multiples of ~ 1 mHz with 222.93% integrated power relative to the GR background, and a $\sim 10\%$ TRZ suppression near 5 mHz. With detection $\text{SNR} > 12$ million,

these signatures are trivially observable by LISA and have no counterpart in standard astrophysics. A single 4-year LISA dataset is sufficient to confirm or rule out UQFF vacuum field coupling at high significance.

Validator: `validate_lisa_extended.py` — PASSED (`compute_aether_noise_spectrum`)

- PASSED: `compute_aether_noise_spectrum`
- ALL TESTS PASSED - LISA extended methods validated

References

- `validate_lisa_extended.py`
- `paper18_validate_lisa_extended.out`

Appendix: UQFF Production Framework Reference (v4.75+)

Added by `upgrade_early_whitepapers.py` (v4.75). This appendix cross-references the production physics constants and master equations to enable reproducibility against the current codebase state.

A.1 Calibration Constants

Symbol	Value	Description
κ	$5.0 \times 10^{-4} \text{ day}^{-1}$	UQFF exponential decay rate
[SSq]	0.57	Universal Quantized Factor
β_i	0.60-0.61	Buoyancy coupling coefficient
k_1	1.5	Ug1 DPM-dipole coupling
k_2	1.2	Ug2 outer-bubble charge coupling
k_3	1.8	Ug3 string-rotation coupling
k_4	2.0	Ug4 vacuum-concentration coupling
η	10^{-22}	Inertia tensor scale
$E_{\text{react}}(0)$	10^{46} J	Reference reactive energy

A.2 F_U Master Equation (Complete — 4 terms)

$$F_U = U_{g1} + U_{g2} + U_{g3} + U_{g4} + U_{bi} + U_m - \sum_{i=1}^4 [\lambda_i \cdot U_i(r, t) \cdot E_{\text{react}}]$$

Term	Description	Implementation
Ug1	DPM magnetic dipole	<code>compute_Ug1_SOURCE4 / compute_Ug1()</code>
Ug2	Outer-field bubble (charge-reactivity)	<code>compute_Ug2_SOURCE4 / compute_Ug2()</code>
Ug3	Magnetic string rotation	<code>compute_Ug3_SOURCE4 / compute_Ug3()</code>
Ug4	Vacuum concentration (star-BH)	<code>compute_Ug4_SOURCE4 / compute_Ug4()</code>
Ubi	Buoyancy force	<code>compute_Ubi_SOURCE4 / compute_Ubi()</code>
Um	Universal Magnetism (Heaviside-amplified)	<code>compute_Um_SOURCE4 / compute_Um()</code>
$-\sum \lambda_i \cdot U_i \cdot E_{\text{react}}$	4th dissipation term (PAPER_420)	<code>compute_FU_SOURCE4 / full pipeline</code>

4th dissipation term parameters (PAPER_420):

$\lambda_1=10^{-10}$, $\lambda_2=10^{-12}$, $\lambda_3=10^{-11}$, $\lambda_4=10^{-13}$ (free parameters, not yet empirically calibrated)

A.3 Um Heaviside Phase-Transition Amplifier (PAPER_421)

$$U_m^{\text{full}} = U_m^{\text{base}} \times (1 + 10^{13} \Theta(\rho_{\text{SCm}} - \rho_c)) \times (1 + A_q \cos(\Delta\omega t))$$

Symbol	Value	Description
ρ_c	10^{15} kg/m^3	SCm critical superconducting density
A_q	0.1	Quasi-periodic beating amplitude (10%)
$\Delta\omega$	$2\pi/(434 \cdot 365.25) \text{ rad/day}$	434-year Gleisberg supercycle

A.4 UQFF Four Operational Modes

Mode	Dominant Term	Primary Use Case
Compressed	Ug_sum + Newtonian base	Isolated stellar/BH systems
Resonant	5 resonance frequencies (aDPM, aTHz, ...)	Multi-scale field interactions
Buoyant	$\beta_i \times \text{Ubi}$	Expanding nebulae, stellar winds
Superconductive	$\text{Um} \times (1 + 10^{13} \cdot f_H)$	Magnetars, SCm critical-density regime

Implementation status: all 4 modes operational in MAIN_1_CoAnQi.cpp, CondensedPhysics.py, and CondensedPhysics2.py.